

Draft Genome Sequence of *Helicobacter fennelliae* Strain MRY12-0050, Isolated from a Bacteremia Patient

Emiko Rimbara,^a Mari Matsui,^a Shigetarou Mori,^a Satowa Suzuki,^a Masato Suzuki,^a Hyun Kim,^a Tsuyoshi Sekizuka,^b Makoto Kuroda,^b Keigo Shibayama^a

Department of Bacteriology II, National Institute of Infectious Diseases, Musashimurayama-shi, Tokyo, Japan^a; Laboratory of Bacterial Genomics, Pathogen Genomics Center, National Institute of Infectious Diseases, Shinjuku-ku, Tokyo, Japan^b

***Helicobacter fennelliae*, a human enterohepatic pathogen, causes bacteremia and colitis. We isolated *H. fennelliae* strain MRY12-0050 from a female patient; this strain was isolated from 2 other patients from the same hospital during the same period, suggesting human-to-human transmission. This is the first report of an *H. fennelliae* genome sequence.**

Received 3 July 2013 Accepted 11 July 2013 Published 8 August 2013

Citation Rimbara E, Matsui M, Mori S, Suzuki S, Suzuki M, Kim H, Sekizuka T, Kuroda M, Shibayama K. 2013. Draft genome sequence of *Helicobacter fennelliae* strain MRY12-0050, isolated from a bacteremia patient. *Genome Announc.* 1(4):e00512-13. doi:10.1128/genomeA.00512-13.

Copyright © 2013 Rimbara et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](http://creativecommons.org/licenses/by/3.0/).

Address correspondence to Emiko Rimbara, rimbara@nih.go.jp.

Helicobacter species are Gram-negative, spiral bacteria that are categorized into 2 groups, (i) gastric helicobacters and (ii) enterohepatic helicobacters. *Helicobacter pylori* is a major gastric *Helicobacter*, and the genome sequences of several gastric *Helicobacter* strains have been analyzed so far. On the other hand, analyses of genomic sequences for enterohepatic helicobacters have been limited. *H. fennelliae* is an enterohepatic *Helicobacter* that causes bacteremia mainly in immunocompromised hosts (1–3). In this study, *H. fennelliae* strain MRY12-0050 was isolated from the blood of a female patient with non-Hodgkin lymphoma. At first, the strain was misidentified as *H. cinaedi* because the strains have similar morphologies. It was later identified as *H. fennelliae* by sequencing of both the 16S rRNA and 23S rRNA genes. *H. fennelliae* was isolated from 2 other patients from the same hospital ward. Pulsed-field gel electrophoresis (PFGE) of 3 isolates showed the same PFGE pattern between isolates, suggesting human-to-human transmission (4). Since genome-sequencing data for *H. fennelliae* have not yet been reported, we herein describe a draft genome sequence for *H. fennelliae* MRY 12-0050, which causes nosocomial infection.

H. fennelliae was cultured in brucella agar (Becton Dickinson, NJ) containing 5% horse blood under microaerobic conditions with hydrogen, which was provided by the gas replacement method by using an anaerobic gas mixture (i.e., H₂, 10%; CO₂, 10%; and N₂, 80%). Genomic DNA was prepared, and a Roche 454 Life Sciences genome sequencer junior system was used to generate DNA sequences at 32× coverage. The same genomic DNA was analyzed by the Illumina MiSeq system paired-end sequences, and the resulting sequence represents 127× coverage. Sequences obtained by using the Roche 454 were assembled using Newbler Assembler v2.7, whereas sequences obtained by using the Illumina MiSeq were assembled with Geneious. The assembled sequences were merged by using Geneious to generate 49 contigs. The contig N₅₀ was approximately 157 kb in length, and the largest contig assembled was approximately 300 kb. Finally, the DNA

sequences were annotated by using the RAST (Rapid Annotation using Subsystem Technology) server (5).

The whole genome is 2.15 Mb in size, has a G+C content of 37.9%, and contains 2,507 genes (2,467 protein-coding genes and 40 structural RNAs). RAST annotation showed that *H. cinaedi* strain CCUG 18818 (score 502), *H. hepaticus* strain ATCC 51449 (score 456), and *H. canadensis* strain MIT 98-5491 (score 444) are the closest neighbors of *H. fennelliae* MRY12-0050. Although *H. cinaedi* and *H. hepaticus* have been known to express cytolethal distending toxin (CDT), which causes DNA damage to target cells, a CDT cluster was not identified in *H. fennelliae* MRY12-0050. Because both *H. cinaedi* and *H. fennelliae* are likely to cause a nosocomial infection and bacteremia in humans, further analyses are needed to elucidate the pathogenic factor in *H. fennelliae*.

Nucleotide sequence accession numbers. This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession no. [BASD000000000](https://www.ncbi.nlm.nih.gov/nuccore/BASD000000000). The version described in this paper is the first version, BASD01000000.

ACKNOWLEDGMENTS

We thank Yuko Sasaki, Kumiko Kai, Yoshie Taki, and Yumiko Yoshimura for their technical assistance. We also thank Shunji Takahashi, Satoshi Yamamoto, and Masaya Mukai for providing bacterial isolates and clinical information.

This study was supported by a Grant-in-Aid for Young Scientists (B) from the Japan Society for the Promotion of Science (no. 25860105) and a grant from the Ministry of Health, Labor, and Welfare of Japan (H24-Shinkou-Ippan-010).

REFERENCES

1. Burnens AP, Stanley J, Schaad UB, Nicolet J. 1993. Novel campylobacter-like organism resembling *Helicobacter fennelliae* isolated from a boy with gastroenteritis and from dogs. *J. Clin. Microbiol.* 31:1916–1917.
2. Hsueh PR, Teng LJ, Hung CC, Chen YC, Yang PC, Ho SW, Luh KT. 1999. Septic shock due to *Helicobacter fennelliae* in a non-human immunodeficiency virus-infected heterosexual patient. *J. Clin. Microbiol.* 37:2084–2086.
3. Smuts HE, Lastovica AJ. 2011. Molecular characterization of the 16S rRNA

- gene of *Helicobacter fennelliae* Isolated from stools and blood cultures from paediatric patients in South Africa. J. Pathog. doi:[10.4061/2011/217376](https://doi.org/10.4061/2011/217376).
4. Rimbara E, Mori S, Kim H, Matsui M, Suzuki S, Takahashi S, Yamamoto S, Mukai M, Shibayama K. 2013. *Helicobacter cinaedi* and *Helicobacter fennelliae* transmission in a hospital from 2008 to 2012. J. Clin. Microbiol. 51:2439–2442.
 5. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: rapid annotations using subsystems technology. BMC Genomics 9:75. doi:[10.1186/1471-2164-9-75](https://doi.org/10.1186/1471-2164-9-75).

Correction for Rimbara et al., Draft Genome Sequence of *Helicobacter fennelliae* Strain MRY12-0050, Isolated from a Bacteremia Patient

Emiko Rimbara,^a Mari Matsui,^a Shigetarou Mori,^a Satowa Suzuki,^a Masato Suzuki,^a Hyun Kim,^a Tsuyoshi Sekizuka,^b Makoto Kuroda,^b Keigo Shibayama^a

Department of Bacteriology II, National Institute of Infectious Diseases, Musashimurayama-shi, Tokyo, Japan^a; Laboratory of Bacterial Genomics, Pathogen Genomics Center, National Institute of Infectious Diseases, Shinjuku-ku, Tokyo, Japan^b

Volume 1, no. 4, [e00512-13](#), 2013. After publication, we detected the presence of a putative cytolethal distending toxin (CDT) gene cluster in *Helicobacter fennelliae* MRY12-0050. In the last paragraph before the accession numbers, the third sentence should read as follows: “*H. cinaedi* and *H. hepaticus* have been reported to express cytolethal distending toxin (CDT), which causes DNA damage to target cells, and a putative CDT gene cluster was also identified in *H. fennelliae* MRY12-0050 (with the gene product sharing 31%, 36%, and 32% amino acid sequence identity with CdtA, CdtB, and CdtC, respectively, of *H. cinaedi* strain CCUG 18818).”

Published 21 July 2016

Copyright © 2016 Rimbara et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](#).