

Genome Sequence of *Paenibacillus polymyxa* ATCC 12321, a Promising Strain for Optically Active (*R,R*)-2,3-Butanediol Production

Ying-Jia Tong, Xiao-Jun Ji, Lu-Gang Liu, Meng-Qiu Shen, He Huang

State Key Laboratory of Materials-Oriented Chemical Engineering, College of Biotechnology and Pharmaceutical Engineering, Nanjing University of Technology, Nanjing, People's Republic of China

***Paenibacillus polymyxa* is a potential strain for (*R,R*)-2,3-butanediol production. Here, we report an annotated draft genome sequence of *P. polymyxa* strain ATCC 12321, which contains 4,429 protein-coding genes and 49 structural RNAs. This genome sequence provides a genetic basis for a better understanding of the mechanism for the accumulation of highly optically active (*R,R*)-2,3-butanediol.**

Received 26 June 2013 Accepted 3 July 2013 Published 1 August 2013

Citation Tong Y-J, Ji X-J, Liu L-G, Shen M-Q, Huang H. 2013. Genome sequence of *Paenibacillus polymyxa* ATCC 12321, a promising strain for optically active (*R,R*)-2,3-butanediol production. *Genome Announc.* 1(4):e00572-13. doi:10.1128/genomeA.00572-13.

Copyright © 2013 Tong et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](http://creativecommons.org/licenses/by/3.0/).

Address correspondence to Xiao-Jun Ji, xiaojunji@njut.edu.cn, or He Huang, biotech@njut.edu.cn.

2,3-Butanediol (2,3-BD) is one of the promising bio-based bulk chemicals due to its extensive industrial applications (1–4). Furthermore, due to its two chiral centers, the molecule of 2,3-BD contains three stereoisomers, namely (*S,S*)-, (*R,R*)-, and *meso*-2,3-BD (5). Among the three stereoisomers, the optically active (*R,R*)-2,3-BD has special applications in the asymmetric synthesis of valuable chiral specialty chemicals (6). Various bacteria were investigated to produce 2,3-BD (1, 2), while only *Paenibacillus polymyxa* (formerly *Bacillus polymyxa*) had the ability to produce (*R,R*)-2,3-BD at over 98% optical purity (7–9). Generally, in addition to (*R,R*)-2,3-BD, the strain produced a small amount of *meso* stereoisomer at the same time (1, 5), and the ratio of these two stereoisomers was usually affected by the environmental conditions, such as aeration and pH (10). However, the mechanism is not clear. Furthermore, the reported mechanisms for the two-stereoisomer formation in *P. polymyxa* were still controversial (5). Studies being performed to elucidate these detailed mechanisms are now focused on the analysis and annotation of a complete genome sequence. To date, three complete genome sequences and two draft genome sequences of *P. polymyxa* strains have been stored in the NCBI database, but these *P. polymyxa* strains all belong to plant-growth-promoting rhizobacteria, not the (*R,R*)-2,3-BD-producing strain. Therefore, sequencing *P. polymyxa* ATCC 12321, which is a promising strain for optically active (*R,R*)-2,3-BD production (8), will not only enrich the genome sequence database of *P. polymyxa*, but it also will help to elucidate the genetic background of this useful strain.

Here, we present the draft genome sequence of strain ATCC 12321, obtained using the Illumina HiSeq 2000 next-generation DNA platform. Sequencing was performed by Shanghai Majorbio Pharm Technology Co., Ltd., with a paired-end library. The reads were trimmed and *de novo* assembled with SOAPdenovo (<http://soap.genomics.org.cn/>) v1.05. Open reading frames (ORFs) were identified by the program Glimmer 3.0 (<http://ccb.jhu.edu/software/glimmer/index.shtml>).

These ORFs were further annotated by comparison with the NCBI nr database and BLASTp (blast 2.2.24). The rRNAs were predicted by RNAmmer (11), and tRNAs were predicted by tRNAscan (12).

The draft genome sequence of strain ATCC 12321 comprises 4,136,795 bp, which is assembled into 92 contigs. The N_{50} quality measurement of the contigs is 172,434 bp and the largest contig assembled is approximately 587,692 bp. It has a G+C content of 46.004%. There are 4,429 predicted protein-coding sequences in the genome sequence. The chromosome has 1 rRNA operon and 48 tRNAs, as predicted by RNAmmer and tRNAscan, respectively.

The genome sequence of *P. polymyxa* ATCC 12321 serves as a basis for further investigation of the molecular basis of its high optically active (*R,R*)-2,3-BD production ability. Relatively detailed annotations will further reveal the mechanism for the (*R,R*)- and *meso*-stereoisomer accumulation ratio variations in response to different environmental conditions.

Nucleotide sequence accession numbers. This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession no. [ARYD000000000](https://www.ncbi.nlm.nih.gov/nuccore/ARYD000000000). The version described in this paper is the first version, accession no. ARYD01000000.

ACKNOWLEDGMENTS

This work was financially supported by the National Science Foundation for Distinguished Young Scholars of China (no. 21225626), the National Basic Research Program of China (no. 2011CBA00807), the National Natural Science Foundation of China (no. 21006049), the National High Technology Research and Development Program of China (no. 2011AA02A207), and the Priority Academic Program Development of Jiangsu Higher Education Institutions.

REFERENCES

1. Ji XJ, Huang H, Ouyang PK. 2011. Microbial 2,3-butanediol production: a state-of-the-art review. *Biotechnol. Adv.* 29:351–364.
2. Ji XJ, Huang H, Nie ZK, Qu L, Xu Q, Tsao GT. 2012. Fuels and chemicals from hemicellulose sugars. *Adv. Biochem. Eng. Biotechnol.* 128:199–224.

3. Zhang WG, Yu DH, Ji XJ, Huang H. 2012. Efficient dehydration of bio-based 2,3-butanediol to butanone over boric acid modified HZSM-5 zeolites. *Green Chem.* 14:3441–3450.
4. Ji XJ, Xia ZF, Fu NH, Nie ZK, Shen MQ, Tian QQ, Huang H. 2013. Cofactor engineering through heterologous expression of an NADH oxidase and its impact on metabolic flux redistribution in *Klebsiella pneumoniae*. *Biotechnol. Biofuels* 6:7.
5. Shen MQ, Ji XJ, Nie ZK, Xia ZF, Yang H, Huang H. 2013. Biotechnological production of 2,3-butanediol stereoisomers: synthetic mechanism and realized methods. *Chin. J. Catal.* 34:351–360.
6. Sadhu KM, Matteson DS, Hurst GD, Kurosky JM. 1984. (*R,R*)-2,3-Butanediol as chiral directing group in the synthesis of (α S)- α -chloro boronic esters. *Organometallics* 3:804–806.
7. De Mas C, Jansen NB, Tsao GT. 1988. Production of optically active 2,3-butanediol by *Bacillus polymyxa*. *Biotechnol. Bioeng.* 31:366–377.
8. Yu B, Sun J, Bommarreddy RR, Song L, Zeng AP. 2011. Novel (*2R,3R*)-2,3-butanediol dehydrogenase from an industrially potential strain *Paenibacillus polymyxa* ATCC 12321. *Appl. Environ. Microbiol.* 77:4230–4233.
9. Gao J, Xu H, Li QJ, Feng XH, Li S. 2010. Optimization of medium for one-step fermentation of inulin extract from jerusalem artichoke tubers using *Paenibacillus polymyxa* ZJ-9 to produce *R,R*-2,3-butanediol. *Bioresour. Technol.* 101:7087–7093.
10. Nakashimada Y, Kanai K, Nishio N. 1998. Optimization of dilution rate, pH and oxygen supply on optical purity of 2,3-butanediol produced by *Paenibacillus polymyxa* in chemostat culture. *Biotechnol. Lett.* 20: 1133–1138.
11. Lagesen K, Hallin P, Rødland EA, Staerfeldt HH, Rognes T, Ussery DW. 2007. RNAmmer: consistent and rapid annotation of ribosomal RNA genes. *Nucleic Acids Res.* 35:3100–3108.
12. Lowe TM, Eddy SR. 1997. tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res.* 25: 955–964.