

Draft Genome Sequence of *Veillonella parvula* HSIVP1, Isolated from the Human Small Intestine

Bartholomeus van den Bogert,^{a,b} Jos Boekhorst,^{c,d} Eddy J. Smid,^{a,e} Erwin G. Zoetendal,^{a,b} Michiel Kleerebezem^{a,b,d,f}

Top Institute Food and Nutrition (TIFN), Wageningen, the Netherlands^a; Laboratory of Microbiology, Wageningen University, Wageningen, the Netherlands^b; Centre for Molecular and Biomolecular Informatics, Radboud University Medical Centre, Nijmegen, the Netherlands^c; NIZO Food Research B.V., Ede, the Netherlands^d; Laboratory of Food Microbiology, Wageningen University, Wageningen, the Netherlands^e; Host-Microbe Interactomics Group, Wageningen University, Wageningen, the Netherlands^f

***Veillonella* species are frequently encountered commensals in the human small intestine. Here, we report the draft genome sequence of the first cultured representative from this ecosystem, *Veillonella parvula* strain HSIVP1. The genome is predicted to encode all the necessary enzymes required for the pathway involved in the conversion of lactate to propanoate.**

Received 28 October 2013 Accepted 1 November 2013 Published 5 December 2013

Citation van den Bogert B, Boekhorst J, Smid EJ, Zoetendal EG, Kleerebezem M. 2013. Draft genome sequence of *Veillonella parvula* HSIVP1, isolated from the human small intestine. *Genome Announc.* 1(6):e00977-13. doi:10.1128/genomeA.00977-13.

Copyright © 2013 van den Bogert et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](http://creativecommons.org/licenses/by/3.0/).

Address correspondence to Michiel Kleerebezem, michiel.kleerebezem@wur.nl.

The genus *Veillonella* encompasses Gram-negative cocci that are commonly encountered in the human oral cavity (1, 2) and parts of the gastrointestinal tract, including the esophagus (3), stomach (4), and small intestine (5–7) (M. M. Leimena and B. van den Bogert, unpublished data). *Veillonella* spp. often co-occur with streptococci in these ecosystems, which is likely related to the capacity of *Veillonella* to effectively utilize lactate, derived from the fermentation of sugars by streptococci, as an energy source (8–10).

Using a selective cultivation approach, small intestine isolates closely related to *Veillonella parvula* were obtained from ileostoma effluent (11). The draft genome sequence of a representative isolate, *V. parvula* HSIVP1, was determined to investigate its metabolic capacity, with a special focus on lactate utilization.

Genomic DNA from *V. parvula* HSIVP1 was sequenced using 454 GS FLX (Roche) technology in combination with titanium chemistry, producing 168,415 reads of ~300 bp, and Illumina HiSeq 2000 technology, producing 10,356,186 paired reads of 50 bp, from 3-kb mate-pair libraries (GATC Biotech, Konstanz, Germany). Pyrosequencing reads were assembled using the Celera Assembler version 6.1 (http://sourceforge.net/apps/mediawiki/wgs-assembler/index.php?title=Main_Page) in 27 contigs. A pseudoassembly was constructed by placing the contigs in their likely order based on paired-read Illumina sequencing data using the SSPACE software version 1.1 (12) and a synteny comparison with the genome of a closely related strain, *V. parvula* DSM 2008 (GenBank accession no. NC_013520). The pseudoassembly was manually screened for inconsistencies using the Artemis comparison tool (13). The HSIVP1 genome was remarkably similar to that of *V. parvula* DSM 2008, displaying only minor chromosomal inversions (<5,000 bp) (B. van den Bogert, unpublished data). The final assembly of the *V. parvula* HSIVP1 genome contains 2,177,885 bp, with an average ~475-fold coverage and a G+C content of 38.51%, and the RAST server (14)-based annotation contains 2,014 predicted protein-coding genes.

The genes in the HSIVP1 genome were assigned to Clusters of

Orthologous Groups (COG) (15) categories, using a BLASTp comparison with the COG database (NCBI, <ftp://ftp.ncbi.nih.gov/pub/COG/COG>) with an alignment *E* value cutoff of 10⁻³. As most *Veillonella* species cannot ferment carbohydrates (16), it is no surprise that the genome of HSIVP1 carries few genes (2.4%) assigned to carbohydrate transport and metabolism. A large fraction of the protein-coding genes was assigned to functions in energy production and conversion, which encompasses most of the genes known to be required for the conversion of lactate to propanoate. This pathway was completely encoded in the HSIVP1 genome and includes the characteristic methylmalonyl-coenzyme A (CoA) decarboxylase that generates a transmembrane electrochemical (Na⁺) gradient (17). This pathway is a critical component of the metabolic relationship in the small intestine ecosystem, where *Veillonella* is proposed to generate energy from the fermentation of lactate produced by (diet-derived) carbohydrate-fermenting lactic acid bacteria (e.g., *Streptococcus*). This proposed metabolic relationship is supported by the high level of expression of the lactate import permease and membrane-associated lactate conversion machinery of *Veillonella* in the small intestine (Leimena and van den Bogert, unpublished).

Nucleotide sequence accession numbers. This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession no. [ASKE00000000](https://www.ncbi.nlm.nih.gov/nuclink/ASKE00000000). The version described in this paper is version ASKE01000000.

ACKNOWLEDGMENTS

This project was partially supported by the Netherlands Bioinformatics Centre (NBIC).

We thank Christopher Bauser and Julia Löcherbach of GATC Biotech (Konstanz, Germany) for assistance in the setup of genome sequencing.

REFERENCES

1. Aas JA, Paster BJ, Stokes LN, Olsen I, Dewhirst FE. 2005. Defining the normal bacterial flora of the oral cavity. *J. Clin. Microbiol.* 43:5721–5732.
2. Keijsers BJ, Zaura E, Huse SM, van der Vossen JM, Schuren FH,

- Montijn RC, ten Cate JM, Crielaard W. 2008. Pyrosequencing analysis of the oral microflora of healthy adults. *J. Dent. Res.* **87**:1016–1020.
3. Pei Z, Bini EJ, Yang L, Zhou M, Francois F, Blaser MJ. 2004. Bacterial biota in the human distal esophagus. *Proc. Natl. Acad. Sci. U. S. A.* **101**: 4250–4255.
 4. Bik EM, Eckburg PB, Gill SR, Nelson KE, Purdom EA, Francois F, Perez-Perez G, Blaser MJ, Relman DA. 2006. Molecular analysis of the bacterial microbiota in the human stomach. *Proc. Natl. Acad. Sci. U. S. A.* **103**:732–737.
 5. van den Bogert B, de Vos WM, Zoetendal EG, Kleerebezem M. 2011. Microarray analysis and barcoded pyrosequencing provide consistent microbial profiles depending on the source of human intestinal samples. *Appl. Environ. Microbiol.* **77**:2071–2080.
 6. Nistal E, Caminero A, Herrán AR, Arias L, Vivas S, de Morales JM, Calleja S, de Miera LE, Arroyo P, Casqueiro J. 2012. Differences of small intestinal bacteria populations in adults and children with/without celiac disease: effect of age, gluten diet, and disease. *Inflamm. Bowel Dis.* **18**: 649–656.
 7. Booiijink CC, El-Aidy S, Rajilić-Stojanović M, Heilig HG, Troost FJ, Smidt H, Kleerebezem M, De Vos WM, Zoetendal EG. 2010. High temporal and inter-individual variation detected in the human ileal microbiota. *Environ. Microbiol.* **12**:3213–3227.
 8. Ng SK, Hamilton IR. 1971. Lactate metabolism by *Veillonella parvula*. *J. Bacteriol.* **105**:999–1005.
 9. Eglund PG, Palmer RJ, Jr, Kolenbrander PE. 2004. Interspecies communication in *Streptococcus gordonii*-*Veillonella atypica* biofilms: signaling in flow conditions requires juxtaposition. *Proc. Natl. Acad. Sci. U. S. A.* **101**: 16917–16922.
 10. Zoetendal EG, Raes J, van den Bogert B, Arumugam M, Booiijink CC, Troost FJ, Bork P, Wels M, de Vos WM, Kleerebezem M. 2012. The human small intestinal microbiota is driven by rapid uptake and conversion of simple carbohydrates. *ISME J.* **6**:1415–1426.
 11. van den Bogert B, Erkus O, Boekhorst J, de Goffau M, Smid EJ, Zoetendal EG, Kleerebezem M. 2013. Diversity of human small intestinal *Streptococcus* and *Veillonella* populations. *FEMS Microbiol. Ecol.* **85**: 376–388.
 12. Boetzer M, Henkel CV, Jansen HJ, Butler D, Pirovano W. 2011. Scaffolding pre-assembled contigs using SSPACE. *Bioinformatics* **27**: 578–579.
 13. Carver TJ, Rutherford KM, Berriman M, Rajandream MA, Barrell BG, Parkhill J. 2005. ACT: the Artemis comparison tool. *Bioinformatics* **21**: 3422–3423.
 14. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: rapid annotations using subsystems technology. *BMC Genomics* **9**:75. doi:10.1186/1471-2164-9-75.
 15. Tatusov RL, Natale DA, Garkavtsev IV, Tatusova TA, Shankavaram UT, Rao BS, Kiryutin B, Galperin MY, Fedorova ND, Koonin EV. 2001. The COG database: new developments in phylogenetic classification of proteins from complete genomes. *Nucleic Acids Res.* **29**:22–28.
 16. De Vos P, Garrity G, Jones D, Krieg NR, Ludwig W, Rainey FA, Schleifer K, Whitman WB. 2009. *Bergey's manual of systematic bacteriology*, 2nd ed, vol 3. Springer Verlag, New York, NY.
 17. Hilpert W, Dimroth P. 1991. On the mechanism of sodium ion translocation by methylmalonyl-CoA decarboxylase from *Veillonella alcalescens*. *Eur. J. Biochem.* **195**:79–86.