



Metagenomic Exploration of *Atelerix albiventris* Gut Microbiome

Makhabbat Amanbayeva,^a Elmira Anarkulova,^{a,b}  Andrey Bogoyavlenskiy,^b Madina Alexyuk,^b Anar Imangazy,^a Vladimir Berezin^b

^aAbai Kazakh National Pedagogical University, Almaty, Kazakhstan

^bResearch and Production Center for Microbiology and Virology, Almaty, Kazakhstan

ABSTRACT Here, we report the metagenomic analysis of the gut of *Atelerix albiventris*, an animal typically kept as a pet in Kazakhstan. In this case, shotgun metagenomic sequencing of the RNA and DNA viral community was performed.

Nowadays, pets are becoming more and more important (1, 2). At the same time, not only familiar cats and dogs but also exotic hedgehogs, lizards, snails, and spiders are being acquired as pets. The introduction of exotic domestic pets in the house creates the need for a more thorough study of the pets' ability to transmit infectious diseases to humans; therefore, the investigation of their microbiomes is an important task in assessing the epidemiological well-being of the population.

Here, the viral and microbial community from the feces of a healthy white-bellied pygmy hedgehog, *Atelerix albiventris* (an ordinary representative of pet shops and minizoo in Almaty, Kazakhstan [43°13'38.6"N, 76°51'52.2"E]), is presented.

Two grams of feces was homogenized in 18 ml of phosphate-buffered saline using the IKA Ultra-Turrax disposable workstation with a disperser tube, filtered using a 0.45- μ m membrane to remove most bacteria, and concentrated by ultracentrifugation using a Beckman Coulter Avanti J30I ultracentrifuge at 29,000 rpm for 2 h at 4°C.

Viral RNA was extracted using a QIAamp viral RNA minikit (Qiagen). The sample extracts were pretreated with RNase-free DNase (Promega). Double-stranded cDNA was obtained with a SuperScript double-stranded cDNA synthesis kit (Invitrogen) according to the manufacturer's instructions. Total DNA was isolated using a PureLink genomic DNA extraction kit (Thermo Fisher Scientific, USA) and stored at -80°C. Genomic DNA and synthesized double-stranded cDNA were pooled. DNA libraries were prepared from 1 ng of the pooled isolated nucleic acids using the Nextera XT DNA sample preparation kit (Illumina, USA). High-throughput sequencing was performed by using an Illumina MiSeq system (paired-end sequencing [2 × 300 bp] with a MiSeq kit v3).

The resulting sequences (852,082 reads) were tested for quality using FastQC v0.11.9 (<https://www.bioinformatics.babraham.ac.uk/projects/fastqc>) and Trimmomatic v0.36 (3) from the Genome Detective tool (4). After the removal of low-quality reads (Q scores of <30) and adapter trimming, 766,874 reads were analyzed by the Kaiju program (5). All tools were run with default parameters unless otherwise specified.

Taxonomic classification of metagenomic data for the *Atelerix albiventris* feces showed that 1% of the sequences corresponded to archaea, 11% to lower eukaryotes, 27% to viruses, 57% to bacteria, and 4% to unclassified organisms (Fig. 1). Although most of the sequences belonged to either normal microflora or microorganisms from food, unfortunately, 1 to 2% of the reads were assigned to microflora potentially capable of causing a number of diseases in humans, including sequences of representatives of *Candida*, *Salmonella*, *Mycobacterium*, *Acinetobacter*, *Klebsiella*, *Chlamydia*, *Yersinia*, *Bartonella*, *Herpesviridae*, *Bunyaviridae*, rabies virus, and the tick-borne encephalitis virus group.

Citation Amanbayeva M, Anarkulova E, Bogoyavlenskiy A, Alexyuk M, Imangazy A, Berezin V. 2021. Metagenomic exploration of *Atelerix albiventris* gut microbiome. *Microbiol Resour Announc* 10:e01342-20. <https://doi.org/10.1128/MRA.01342-20>.

Editor Simon Roux, DOE Joint Genome Institute

Copyright © 2021 Amanbayeva et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Andrey Bogoyavlenskiy, anpav_63@mail.ru.

Received 24 November 2020

Accepted 8 December 2020

Published 7 January 2021




FIG 1 Taxonomic diversity of the microbial and viral communities of *Aterix albiventris*. Taxonomic assignment of the metagenomic reads was based on the NCBI BLAST nonredundant plus eukaryotes database using Kaiju.

Data availability. Raw sequence reads are available under BioProject accession number [PRJNA656141](https://www.ncbi.nlm.nih.gov/bioproject/PRJNA656141) and SRA accession number [SRR12422733](https://www.ncbi.nlm.nih.gov/sra/SRR12422733).

ACKNOWLEDGMENTS

This work was supported by a Grant-in-Aid for Scientific Research from Abai Kazakh National Pedagogical University and by the Ministry of Education and Science of the Republic of Kazakhstan (grant BR05236330).

We declare no conflicts of interest. This article does not contain any studies with human or animal subjects performed by any of the authors.

REFERENCES

- Riley PY, Chomel BB. 2005. Hedgehog zoonoses. *Emerg Infect Dis* 11:1–5. <https://doi.org/10.3201/eid1101.040752>.
- Okada K, Kondo H, Sumi A, Kagawa Y. 2018. A retrospective study of disease incidence in African pygmy hedgehogs (*Aterix albiventris*). *J Vet Med Sci* 80:1504–1510. <https://doi.org/10.1292/jvms.18-0238>.
- Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
- Vilsker M, Moosa Y, Nooij S, Fonseca V, Ghysens Y, Dumon K, Pauwels R, Alcantara LC, Vanden Eynden E, Vandamme A-M, Deforche K, de Oliveira T. 2019. Genome Detective: an automated system for virus identification from high-throughput sequencing data. *Bioinformatics* 35:871–873. <https://doi.org/10.1093/bioinformatics/bty695>.
- Menzel P, Ng KL, Krogh A. 2016. Fast and sensitive taxonomic classification for metagenomics with Kaiju. *Nat Commun* 7:11257. <https://doi.org/10.1038/ncomms11257>.