

# Full-Genome Assembly of Reference Strain *Providencia stuartii* ATCC 33672

K. G. Frey,<sup>a,b</sup> K. A. Bishop-Lilly,<sup>a,b</sup> H. E. Daligault,<sup>c</sup> K. W. Davenport,<sup>c</sup> D. C. Bruce,<sup>c</sup> P. S. Chain,<sup>c</sup> S. R. Coyne,<sup>d</sup> O. Chertkov,<sup>c</sup> T. Freitas,<sup>d\*</sup> J. Jaissle,<sup>d</sup> G. I. Koroleva,<sup>e</sup> J. T. Ladner,<sup>e</sup> T. D. Minogue,<sup>d</sup> G. F. Palacios,<sup>e</sup> C. L. Redden,<sup>a,b</sup> Y. Xu,<sup>c</sup>  S. L. Johnson<sup>c</sup>

Naval Medical Research Center (NMRC), Frederick, Fort Detrick, Maryland, USA<sup>a</sup>; Henry M. Jackson Foundation, Bethesda, Maryland, USA<sup>b</sup>; Los Alamos National Laboratory (LANL), Los Alamos, New Mexico, USA<sup>c</sup>; Diagnostic Systems Division (DSD), United States Army Medical Research Institute of Infectious Diseases (USAMRIID), Fort Detrick, Maryland, USA<sup>d</sup>; Center for Genome Sciences (CGS), USAMRIID, Fort Detrick, Maryland, USA<sup>e</sup>

\* Present address: T. Freitas, Med Fusion, Lewisville, Texas, USA.

**A member of the normal human gut microflora, *Providencia stuartii* is of clinical interest due to its role in nosocomial infections of the urinary tract and because it readily acquires antibiotic resistance. Here, we present the complete genome of *P. stuartii* strain ATCC 33672, consisting of a 4.28-Mbp chromosome and a 48.9-kbp plasmid.**

Received 12 September 2014 Accepted 16 September 2014 Published 23 October 2014

**Citation** Frey KG, Bishop-Lilly KA, Daligault HE, Davenport KW, Bruce DC, Chain PS, Coyne SR, Chertkov O, Freitas T, Jaissle J, Koroleva GI, Ladner JT, Minogue TD, Palacios GF, Redden CL, Xu Y, Johnson SL. 2014. Full-genome assembly of reference strain *Providencia stuartii* ATCC 33672. *Genome Announc*. 2(5):e01082-14. doi:10.1128/genomeA.01082-14.

**Copyright** © 2014 Frey et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](http://creativecommons.org/licenses/by/3.0/).

Address correspondence to S. L. Johnson, shannonj@lanl.gov.

*Providencia stuartii* is a frequent cause of urinary tract infections in hospital patients with long-term indwelling catheters (1). In these patients, bacteriuria may rapidly progress to bacteremia (2, 3). Infection with *P. stuartii* has also been reported to cause peritonitis (4), meningitis (5), pericarditis (6), and infective endocarditis (7). In an unusual case, *P. stuartii* was identified as the cause of a renal abscess mistakenly attributed to an infection with *Pasteurella* (8). Additionally, nosocomial dissemination of multidrug-resistant *P. stuartii* has been reported in burn units (9, 10) and an intensive care unit (10). Members of the genus *Providencia* harbor resistance to aminopenicillins and narrow-spectrum cephalosporins (11) but also readily acquire resistance to other antibiotics through horizontal transfer (12, 13). This is of great concern to public health due to the ability of *P. stuartii* to further disseminate antimicrobial resistance genes (13, 14). We present the complete genome of strain ATCC 33672, consisting of a 4.28-Mbp chromosome and a 48.9-kbp plasmid.

High-quality genomic DNA was extracted from a purified isolate using the Qiagen Genome-tip 500. Specifically, a 100-ml bacterial culture was grown to stationary phase and nucleic acid extracted as per the manufacturer's recommendations. The draft genome of *P. stuartii* ATCC 33672 included a combination of Illumina (15) and 454 technologies (16). For this genome, we constructed and sequenced a 100-bp Illumina library (372-fold genome coverage) and a long-insert paired-end 454 library (6,689 ± 1,140-bp insert; 5-fold genome coverage). Raw data are available in the Short Read Archive (SRA) under accession numbers SRX687103 (Illumina), SRX687104 (Illumina), and SRX687264 (454).

The 454 paired-end data were assembled in Newbler (16), and those consensus sequences were computationally shredded into 2-kbp overlapping fake reads (shreds). The Illumina sequencing

data were assembled with Velvet (17), and the consensus sequences were computationally shredded into 1.5-kbp overlapping shreds. All data were additionally assembled in AllPaths (18), and the consensus sequences were computationally shredded into 5-kbp overlapping shreds. We integrated the 454 Newbler consensus shreds, the Illumina Velvet consensus shreds, AllPaths consensus shreds, and the 454 paired-end library read pairs using parallel Phrap (High Performance Software, LLC). Possible misassemblies were corrected and repeat regions verified using in-house scripts and manual editing in Consed (19–21).

The complete genome assembly of *P. stuartii* ATCC 33672, including a 4,285,951-bp circular chromosome (41.5% G+C content) and a 48,866-bp circular plasmid (42.8% G+C content), was annotated utilizing an Ergatis-based (22) workflow with minor manual curation. The annotated genome contains 4,094 predicted genes, including 3,926 protein-coding genes, 80 tRNA genes, and 22 rRNA genes. A total of 92 virulence genes were noted, including resistance to fluoroquinolones, β-lactams, and aminoglycosides.

**Nucleotide sequence accession numbers.** The nucleotide sequences for *P. stuartii* ATCC 33672 have been deposited in GenBank under accession numbers CP008920 (chromosome) and CP008919 (plasmid).

## ACKNOWLEDGMENTS

Funding for this effort was provided by the Defense Threat Reduction Agency's Joint Science and Technology Office (DTRA J9-CB/JSTO).

This paper is approved by LANL for unlimited release (LA-UR-14-26025).

The views expressed in this article are those of the authors and do not necessarily reflect the official policy or position of the Department of the Navy, Department of Defense, or the U.S. Government.

## REFERENCES

- Warren JW. 1986. *Providencia stuartii*: a common cause of antibiotic-resistant bacteriuria in patients with long-term indwelling catheters. *Rev. Infect. Dis.* 8:61–67. <http://dx.doi.org/10.1093/clinids/8.1.61>.
- Kim BN, Kim NJ, Kim MN, Kim YS, Woo JH, Ryu J. 2003. Bacteraemia due to tribe *Proteaeae*: a review of 132 cases during a decade (1991–2000). *Scand. J. Infect. Dis.* 35:98–103. <http://dx.doi.org/10.1080/0036554021000027015>.
- Woods TD, Watanakunakorn C. 1996. Bacteremia due to *Providencia stuartii*: review of 49 episodes. *South. Med. J.* 89:221–224. <http://dx.doi.org/10.1097/00007611-199602000-00013>.
- Unverdi S, Akay H, Ceri M, Inal S, Altay M, Demiroz AP, Duranay M. 2011. Peritonitis due to *Providencia stuartii*. *Perit. Dial. Int.* 31:216–217. <http://dx.doi.org/10.3747/pdi.2010.00172>.
- Sipahi OR, Bardak-Ozdemir S, Ozgiray E, Aydemir S, Yurtseven T, Yamazhan T, Tasbakan M, Ulusoy S. 2010. Meningitis due to *Providencia stuartii*. *J. Clin. Microbiol.* 48:4667–4668. <http://dx.doi.org/10.1128/JCM.01349-10>.
- Simon C, Dieli M, Brucato A, Pedrotti P, Brambilla P, Curri SF, Senni M, Pericotti S, Suter F, Ferrazzi P. 2010. Images in cardiovascular medicine. Bacterial pericarditis due to *Providencia stuartii*: an atypical case of relapsing pericarditis. *Circulation* 122:e401–e403. <http://dx.doi.org/10.1161/CIRCULATIONAHA.110.943118>.
- Krake PR, Tandon N. 2004. Infective endocarditis due to *Providencia stuartii*. *South. Med. J.* 97:1022–1023.
- Chamberland RR, McElvania TeKippe E, Burnham CA, Kennedy DJ. 2013. Renal abscess caused by a *Providencia stuartii* isolate biochemically misidentified as *Pasteurella*. *J. Clin. Microbiol.* 51:2775–2777. <http://dx.doi.org/10.1128/JCM.00937-13>.
- Saida NB, Thabet L, Messadi A, Bouselmi K, Turki A, Boukadida J. 2008. Clonality of *Providencia stuartii* isolates involved in outbreak that occurred in a burn unit. *Burns J. Int. Soc. Burn Injuries* 34:829–834. <http://dx.doi.org/10.1016/j.burns.2007.09.008>.
- Zavascki AP, Carvalhaes CG, da Silva GL, Tavares Soares SP, de Alcantara LR, Elias LS, Sandri AM, Gales AC. 2012. Outbreak of carbapenem-resistant *Providencia stuartii* in an intensive care unit. *Infect. Control Hosp. Epidemiol.* 33:627–630. <http://dx.doi.org/10.1086/665730>.
- Aubert D, Naas T, Lartigue MF, Nordmann P. 2005. Novel genetic structure associated with an extended-spectrum beta-lactamase *bla*<sub>VEB</sub> gene in a *Providencia stuartii* clinical isolate from Algeria. *Antimicrob. Agents Chemother.* 49:3590–3592. <http://dx.doi.org/10.1128/AAC.49.8.3590-3592.2005>.
- McGann P, Hang J, Clifford RJ, Yang Y, Kwak YI, Kuschner RA, Lesho EP, Waterman PE. 2012. Complete sequence of a novel 178-kilobase plasmid carrying *bla*(NDM-1) in a *Providencia stuartii* strain isolated in Afghanistan. *Antimicrob. Agents Chemother.* 56:1673–1679. <http://dx.doi.org/10.1128/AAC.05604-11>.
- Arpin C, Thabet L, Yassine H, Messadi AA, Boukadida J, Dubois V, Coulange-Mayonnove L, Andre C, Quentin C. 2012. Evolution of an incompatibility group IncA/C plasmid harboring *bla*<sub>CMY-16</sub> and *qnrA6* genes and its transfer through three clones of *Providencia stuartii* during a two-year outbreak in a Tunisian burn unit. *Antimicrob. Agents Chemother.* 56:1342–1349. <http://dx.doi.org/10.1128/AAC.05267-11>.
- Mnif B, Ktari S, Chaari A, Medhioub F, Rhimi F, Bouaziz M, Hammami A. 2013. Nosocomial dissemination of *Providencia stuartii* isolates carrying *bla*<sub>OXA-48</sub>, *bla*<sub>PER-1</sub>, *bla*<sub>CMY-4</sub> and *qnrA6* in a Tunisian hospital. *J. Antimicrob. Chemother.* 68:329–332. <http://dx.doi.org/10.1093/jac/dks386>.
- Bennett S. 2004. Solexa Ltd. *Pharmacogenomics* 5:433–438. <http://dx.doi.org/10.1517/14622416.5.4.433>.
- Margulies M, Egholm M, Altman WE, Attiya S, Bader JS, Bemben LA, Berka J, Braverman MS, Chen YJ, Chen Z, Dewell SB, Du L, Fierro JM, Gomes XV, Godwin BC, He W, Helgesen S, Ho CH, Irzyk GP, Jando SC, Alenquer ML, Jarvie TP, Jirage KB, Kim JB, Knight JR, Lanza JR, Leamon JH, Lefkowitz SM, Lei M, Li J, Lohman KL, Lu H, Makhijani VB, McDade KE, McKenna MP, Myers EW, Nickerson E, Nobile JR, Plant R, Puc BP, Ronan MT, Roth GT, Sarkis GJ, Simons JF, Simpson JW, Srinivasan M, Tartaro KR, Tomasz A, Vogt KA, Volkmer GA, Wang SH, Wang Y, Weiner MP, Yu P, Begley RF, Rothberg JM. 2005. Genome sequencing in microfabricated high-density picolitre reactors. *Nature* 437:376–380. <http://dx.doi.org/10.1038/nature03959>.
- Zerbino DR, Birney E. 2008. Velvet: algorithms for *de novo* short read assembly using de Bruijn graphs. *Genome Res.* 18:821–829. <http://dx.doi.org/10.1101/gr.074492.107>.
- Butler J, MacCallum I, Kleber M, Shlyakhter IA, Belmonte MK, Lander ES, Nusbaum C, Jaffe DB. 2008. ALLPATHS: *de novo* assembly of whole-genome shotgun microreads. *Genome Res.* 18:810–820. <http://dx.doi.org/10.1101/gr.7337908>.
- Ewing B, Green P. 1998. Base-calling of automated sequencer traces using Phred. II. Error Probabilities. *Genome Res.* 8:186–194.
- Ewing B, Hillier L, Wendl MC, Green P. 1998. Base-calling of automated sequencer traces using Phred. I. Accuracy assessment. *Genome Res.* 8:175–185. <http://dx.doi.org/10.1101/gr.8.3.175>.
- Gordon D, Abajian C, Green P. 1998. Consed: a graphical tool for sequence finishing. *Genome Res.* 8:195–202. <http://dx.doi.org/10.1101/gr.8.3.195>.
- Hemmerich C, Buechlein A, Podicheti R, Revanna KV, Dong Q. 2010. An Ergatis-based prokaryotic genome annotation Web server. *Bioinformatics* 26:1122–1124. <http://dx.doi.org/10.1093/bioinformatics/btq090>.