

Genome Sequence Analysis of the Biogenic Amine-Producing Strain *Lactococcus lactis* subsp. *cremoris* CECT 8666 (Formerly GE2-14)

Victor Ladero, Beatriz del Rio, Daniel M. Linares, Maria Fernandez, Baltasar Mayo, M. Cruz Martin, Miguel A. Alvarez

Instituto de Productos Lácteos de Asturias, IPLA-CSIC, Paseo Río Linares s/n, Asturias, Spain

We here report a 2,801,031-bp annotated draft assembly for the *Lactococcus lactis* subsp. *cremoris* GE2-14 genome. This dairy strain produces the biogenic amine putrescine. This sequence may help identify the mechanisms regulating putrescine biosynthesis and throw light on ways to reduce its presence in fermented foods.

Received 12 September 2014 Accepted 16 September 2014 Published 23 October 2014

Citation Ladero V, del Rio B, Linares DM, Fernandez M, Mayo B, Martin MC, Alvarez MA. 2014. Genome sequence analysis of the biogenic amine-producing strain *Lactococcus lactis* subsp. *cremoris* CECT 8666 (formerly GE2-14). *Genome Announc.* 2(5):e01088-14. doi:10.1128/genomeA.01088-14.

Copyright © 2014 Ladero et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/3.0/).

Address correspondence to Victor Ladero, ladero@ipla.csic.es.

Lactococcus lactis is of great economic importance, providing starter cultures for the production of fermented dairy products (1). However, some strains of *L. lactis* can produce putrescine, a toxic biogenic amine (BA) (2, 3).

BA are formed and accumulated in food via microbial metabolism (2). Their ingestion can cause intoxication symptoms (4), and thus, there is a consensus regarding the reduction of BA in foods (5). In dairy products, BA can reach high concentrations (6), with putrescine one of the most abundant (7). Putrescine is mainly produced by the deimination of agmatine (an arginine decarboxylation product) via the agmatine deiminase (AGDI) pathway (7, 8). Environmental and technological factors affect its accumulation (2).

We here report the draft genome of *L. lactis* subsp. *cremoris* CECT 8666 (formerly GE2-14), a strain isolated from artisanal cheese (9) that produces large amounts of putrescine (3). This strain is currently used as a model strain for studying putrescine biosynthesis regulation (10).

A 0.5-kbp genomic library was constructed and subjected to 90-bp paired-end sequencing (providing 30-fold coverage) using a HiSeq 1000 System sequencer (Illumina) (performed at the Beijing Genomics Institute, China). Quality-filtered reads were assembled using Velvet software (<http://www.ebi.ac.uk/~zerbino/velvet/>), resulting in 243 contigs ranging from 202 to 239,606 bp. The total sequence length was 2,801,031 bp, with a G+C content of 35.4%. Annotation was performed by Era7 Bioinformatics (Granada, Spain) using the BG7 pipeline (11), and improved using BLAST analysis results (<http://blast.ncbi.nlm.nih.gov>). The genome contained 2,765 predicted coding sequences. Predicted copies of the 16S, 23S, and 5S rRNA genes were found, as were 53 genes for tRNAs.

Genome analysis confirmed the presence/absence of characteristics identified phenotypically in this strain (9) as carbohydrate utilization genes. Interestingly, this strain is unable to ferment arabinose, yet it possesses the genes required. Detailed analysis detected a frameshift in a permease-encoding (U725_00413) and an α -N-arabinofuranosidase-encoding genes (U725_00414), probably impairing L-arabinose utilization. The genome contained at least four prophages. Indeed, almost one-tenth of the genes (285) were phage related. Several genes with homology to

plasmid replication and mobilization proteins were also found, consistent with the presence of at least four plasmids.

The AGDI cluster involved in putrescine production comprised five genes, *aguR*, *aguB*, *aguD*, *aguA*, and *aguC* (U725_01346 to U725_01350). The AGDI cluster is known to be subject to carbon catabolite repression mediated by the catabolite control protein CcpA (10). Interestingly, in addition to the *ccpA* gene (U725_00364), a second gene involved in carbon catabolite repression (*ccpB*; U725_02733) was identified, although its role in AGDI cluster expression is unknown.

The availability of the *L. lactis* subsp. *cremoris* CECT 8666 genome sequence opens up the possibility of performing transcriptional studies for identifying the genes involved in the regulation of putrescine production. This would improve our knowledge of the factors affecting BA production and accumulation in dairy products, leading to improvements in food safety.

Nucleotide sequence accession numbers. The results of this whole-genome shotgun project were deposited in the DDBJ/EMBL/GenBank database under accession number [AZSI00000000](https://www.ncbi.nlm.nih.gov/nuccore/AZSI00000000) (BioProject PRJNA225671). The version of the genome described here is version AZSI01000000.

ACKNOWLEDGMENTS

This work was funded by the Spanish Ministry of Economy and Competitiveness (AGL2013-45431-R) and the CSIC (201270E144).

We thank Adrian Burton for language assistance.

REFERENCES

1. Limsowitn GKY, Powell IB, Parente E. 1996. Types of starters, p 101–129. *In* Cogan TM, Accolas JP (ed), Dairy starter cultures. VCH, New York, NY.
2. Linares DM, Del Rio B, Ladero V, Martinez N, Fernandez M, Martin MC, Alvarez MA. 2012. Factors influencing biogenic amines accumulation in dairy products. *Front Microbiol.* 3:180. <http://dx.doi.org/10.3389/fmicb.2012.00180>.
3. Ladero V, Rattray FP, Mayo B, Martín MC, Fernández M, Alvarez MA. 2011. Sequencing and transcriptional analysis of the biosynthesis gene cluster of putrescine-producing *Lactococcus lactis*. *Appl. Environ. Microbiol.* 77:6409–6418. <http://dx.doi.org/10.1128/AEM.05507-11>.
4. Ladero V, Calles-Enriquez M, Fernandez M, Alvarez MA. 2010. Toxi-

- ological effects of dietary biogenic amines. *Curr. Nutr. Food Sci.* 6:145–156. <http://dx.doi.org/10.2174/157340110791233256>.
5. EFSA. 2011. Scientific opinion on risk based control of biogenic amine formation in fermented foods. *EFSA J.* 9:2393–2486.
 6. Fernández M, Linares DM, Del Río B, Ladero V, Alvarez MA. 2007. HPLC quantification of biogenic amines in cheeses: correlation with PCR-detection of tyramine-producing microorganisms. *J. Dairy Res.* 74: 276–282. <http://dx.doi.org/10.1017/S0022029907002488>.
 7. Linares DM, Martín MC, Ladero V, Alvarez MA, Fernández M. 2011. Biogenic amines in dairy products. *Crit. Rev. Food Sci. Nutr.* 51:691–703. <http://dx.doi.org/10.1080/10408398.2011.582813>.
 8. Ladero V, Canedo E, Perez M, Martín MC, Fernández M, Alvarez MA. 2012. Multiplex qPCR for the detection and quantification of putrescine-producing lactic acid bacteria in dairy products. *Food Control* 27: 307–313. <http://dx.doi.org/10.1016/j.foodcont.2012.03.024>.
 9. Fernández E, Alegría A, Delgado S, Martín MC, Mayo B. 2011. Comparative phenotypic and molecular genetic profiling of wild *Lactococcus lactis* subsp. *lactis* strains of the *L. lactis* subsp. *lactis* and *L. lactis* subsp. *cremoris* genotypes, isolated from starter-free cheeses made of raw milk. *Appl. Environ. Microbiol.* 77:5324–5335. <http://dx.doi.org/10.1128/AEM.02991-10>.
 10. Linares DM, del Río B, Ladero V, Redruello B, Martín MC, Fernández M, Alvarez MA. 2013. The putrescine biosynthesis pathway in *Lactococcus lactis* is transcriptionally regulated by carbon catabolic repression, mediated by CcpA. *Int. J. Food Microbiol.* 165:43–50. <http://dx.doi.org/10.1016/j.ijfoodmicro.2013.04.021>.
 11. Pareja-Tobes P, Manrique M, Pareja-Tobes E, Pareja E, Tobes R. 2012. BG7: a new approach for bacterial genome annotation designed for next generation sequencing data. *PLoS One* 7:e49239. <http://dx.doi.org/10.1371/journal.pone.0049239>.