

# Draft Genome Sequence of Anammox Bacterium “*Candidatus Scalindua brodae*,” Obtained Using Differential Coverage Binning of Sequencing Data from Two Reactor Enrichments

Daan R. Speth,<sup>a</sup> Lina Russ,<sup>a</sup> Boran Kartal,<sup>a</sup> Huub J. M. op den Camp,<sup>a</sup> Bas E. Dutilh,<sup>b,c,d</sup> Mike S. M. Jetten<sup>a,e</sup>

Ecological Microbiology, IWWR, Radboud University Nijmegen, Nijmegen, The Netherlands<sup>a</sup>; Theoretical Biology and Bioinformatics, Utrecht University, Utrecht, The Netherlands<sup>b</sup>; Centre for Molecular and Biomolecular Informatics, Radboud University Medical Centre, Nijmegen, The Netherlands<sup>c</sup>; Instituto de Biologia, Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brazil<sup>d</sup>; Department of Biotechnology, Delft University of Technology, Delft, The Netherlands<sup>e</sup>

**We present the draft genome of anammox bacterium “*Candidatus Scalindua brodae*,” which at 282 contigs is a major improvement over the highly fragmented genome assembly of related species “*Ca. Scalindua profunda*” (1,580 contigs) which was previously published.**

Received 27 November 2014 Accepted 1 December 2014 Published 8 January 2015

**Citation** Speth DR, Russ L, Kartal B, op den Camp HJM, Dutilh BE, Jetten MSM. 2015. Draft genome sequence of anammox bacterium “*Candidatus Scalindua brodae*,” obtained using differential coverage binning of sequencing data from two reactor enrichments. *Genome Announc.* 3(1):e01415-14. doi:10.1128/genomeA.01415-14.

**Copyright** © 2015 Speth et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Daan R. Speth, d.speth@science.ru.nl.

Anammox bacteria are major players in the global nitrogen cycle, capable of anaerobically oxidizing ammonium to dinitrogen gas, using nitrite as the electron acceptor (1). All currently known anammox bacteria form the monophyletic order *Brocadiales* within the phylum *Planctomycetes* (2). Until now, draft genomes of four anammox species have been reported (3–6). The genome assemblies of “*Candidatus Kuenenia stuttgartiensis*” and “*Ca. Jettenia caeni*” are in 5 and 4 contigs, respectively, whereas the draft genome of “*Ca. Brocadia fulgida*” (411 contigs) is fragmented and the “*Ca. Scalindua profunda*” draft genome (1,580 contigs) is highly fragmented.

Despite advances in culturing techniques, no pure culture of anammox bacteria exists. This restricts genome-sequencing efforts to metagenomic sequencing and binning (7, 8). Here we employed a differential coverage binning approach (9) to increase the confidence of the binning result. We combined sequencing data from Russ et al. (10) with sequencing data from the enrichment culture used as seed for the experimental reactor described in Russ et al. (10). The raw sequence data are available in DDBJ/EMBL/Genbank under accession no. ERX443234 and SRX719339. DNA isolation and sequencing of both enrichments was performed as described previously, using the Powersoil DNA isolation kit (Mo-Bio, Carlsbad, CA, USA) according to the manufacturer’s instructions and the Ion Torrent 200 bp workflow (10).

All data were co-assembled using the CLC genomics workbench (v7.0.4, CLCbio, Aarhus, Denmark) *de novo* assembler, using word size 35 and bubble size 5,000. The obtained contigs were binned with a workflow modified from Albertsen et al. (9) using custom scripts available at [http://www.github.com/dspeth/bioinfo\\_scripts](http://www.github.com/dspeth/bioinfo_scripts). The binned “*Ca. Scalindua brodae*” genome consisted of 282 contigs and was annotated using Prokka 1.10 (11) followed by manual curation. Frameshifts were corrected using the CLC genomics workbench (v7.0.4, CLCbio, Aarhus, Denmark) and a Perl script available at [http://www.github.com/dspeth/bioinfo\\_scripts](http://www.github.com/dspeth/bioinfo_scripts). The

completeness (>92%) of the draft genome was assessed using CheckM (12). The contigs have a total length of 4.1 Mb, average G+C content of 39.6%, and encode 4,016 genes, 39 tRNAs, and 1 rRNA operon.

Hydrazine is a key intermediate in anammox metabolism and the enzymes involved in its turnover are unique to anammox bacteria. The fusion of subunits B and C of the hydrazine synthase operon (*hzsBC*), earlier reported for “*Ca. Scalindua profunda*” (6), was confirmed and additionally the fused genes seemed to have undergone duplication. The presence of two copies of *hzsBC* prohibited resolving their genomic location without mate pair information. As a result, the hydrazine synthase BC gene is present on a separate contig. The hydrazine dehydrogenase is also present in two copies, but their genomic location could be resolved based on sequence difference.

**Nucleotide sequence accession numbers.** This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession no. [JRYO000000000](https://www.ncbi.nlm.nih.gov/nuccore/JRYO000000000). The version described in this paper is the first version, [JRYO010000000](https://www.ncbi.nlm.nih.gov/nuccore/JRYO010000000).

## ACKNOWLEDGMENTS

The work was supported by European Research Council grant 232937 to M.S.M.J. and L.R., NWO VENI grant 863.11.003 to B.K., and BE-Basic grant fp07-2F to D.R.S.

## REFERENCES

- Kartal B, Maalcke WJ, de Almeida NM, Cirpus I, Gloerich J, Geerts W, Op den Camp HJM, Harhangi HR, Janssen-Megens EM, Francoijs K-J, Stunnenberg HG, Keltjens JT, Jetten MSM, Strous M. 2011. Molecular mechanism of anaerobic ammonium oxidation. *Nature* 479:127–130. [http://dx.doi.org/10.1038/nature10453](https://doi.org/10.1038/nature10453).
- Jetten MSM, Op den Camp HJ, Kuenen JG, Strous M. 2010. Description of the order Brocadiales, p 596–603. *In* Krieg NR, Staley JT, Hedlund BP, Paster BJ, Ward N, Ludwig W, Whitman WB (ed), *Bergey’s manual of systematic bacteriology*, vol 4. Springer Verlag, Heidelberg, Germany.
- Strous M, Pelletier E, Manganot S, Rattei T, Lehner A, Taylor MW, Horn M, Daims H, Bartol-Mavel D, Wincker P, Barbe V, Fonknechten

- N, Vallenet D, Segurens B, Schenowitz-Truong C, Médigue C, Collingro A, Snel B, Dutilh BE, Op den Camp HJM, van der Drift C, Cirpus I, van de Pas-Schoonen KT, Harhangi HR, van Niftrik L, Schmid M, Keltjens J, van de Vossenberg J, Kartal B, Meier H, Frishman D, Huynen MA, Mewes H-W, Weissenbach J, Jetten MSM, Wagner M, Le Paslier D. 2006. Deciphering the evolution and metabolism of an anammox bacterium from a community genome. *Nature* 440: 790–794. <http://dx.doi.org/10.1038/nature04647>.
4. Hira D, Toh H, Migita CT, Okubo H, Nishiyama T, Hattori M, Furukawa K, Fujii T. 2012. Anammox organism KSU-1 expresses a NirK-type copper-containing nitrite reductase instead of a NirS-type with cytochrome *cd<sub>1</sub>*. *FEBS Lett* 586:1658–1663. <http://dx.doi.org/10.1016/j.febslet.2012.04.041>.
  5. Ferousi C, Speth DR, Reimann J, Op den Camp HJ, Allen JW, Keltjens JT, Jetten MS. 2013. Identification of the type II cytochrome *c* maturation pathway in anammox bacteria by comparative genomics. *BMC Microbiol* 13:265. <http://dx.doi.org/10.1186/1471-2180-13-265>.
  6. van de Vossenberg J, Woebken D, Maalcke WJ, Wessels HJCT, Dutilh BE, Kartal B, Janssen-Megens EM, Roeselers G, Yan J, Speth D, Glerich J, Geerts W, van der Biezen E, Pluk W, Francoijs K-J, Russ L, Lam P, Malfatti SA, Tringe SG, Haaijer SCM, Op den Camp HJM, Stunnenberg HG, Amann R, Kuypers MMM, Jetten MSM. 2013. The metagenome of the marine anammox bacterium “*Candidatus Scalindua profunda*” illustrates the versatility of this globally important nitrogen cycle bacterium. *Environ Microbiol* 15:1275–1289. <http://dx.doi.org/10.1111/j.1462-2920.2012.02774.x>.
  7. van der Star WRL, Miclea AI, van Dongen UGJM, Muyzer G, Picioranu C, van Loosdrecht MCM. 2008. The membrane bioreactor: A novel tool to grow anammox bacteria as free cells. *Biotechnol Bioeng* 101:286–294. <http://dx.doi.org/10.1002/bit.21891>.
  8. Sharon I, Banfield JF. 2013. Genomes from metagenomics. *Science* 342: 1057–1058. <http://dx.doi.org/10.1126/science.1247023>.
  9. Albertsen M, Hugenholtz P, Skarshewski A, Nielsen KL, Tyson GW, Nielsen PH. 2013. Genome sequences of rare, uncultured bacteria obtained by differential coverage binning of multiple metagenomes. *Nat Biotechnol* 31:533–538. <http://dx.doi.org/10.1038/nbt.2579>.
  10. Russ L, Speth DR, Jetten MSM, Op den Camp HJM, Kartal B. 2014. Interactions between anaerobic ammonium and sulfur-oxidizing bacteria in a laboratory scale model system. *Environ Microbiol* 16:3487–3498. <http://dx.doi.org/10.1111/1462-2920.12487>.
  11. Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068–2069. <http://dx.doi.org/10.1093/bioinformatics/btu153>.
  12. Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. 2014. CheckM: assessing the quality of microbial genomes recovered from isolates, single cells and metagenomes. *PeerJ PrePrints* 2:e554v1. doi: <http://dx.doi.org/10.7287/peerj.preprints.554v1>.