

Complete Genome Sequence of the *Opitutaceae* Bacterium Strain TAV5, a Potential Facultative Methyloph of the Wood-Feeding Termite *Reticulitermes flavipes*

Malini Kotak,^a Jantiya Isanapong,^b Lynne Goodwin,^c David Bruce,^c Amy Chen,^d Cliff S. Han,^c Marcel Huntemann,^d Natalia Ivanova,^d Miriam L. Land,^e Matt Nolan,^d Amrita Pati,^d Tanja Woyke,^d Jorge L. M. Rodrigues^f

Department of Biology, University of Texas, Arlington, Texas, USA^a; King Mongkut's University of Technology North Bangkok, Bangkok, Thailand^b; Los Alamos National Laboratory, Los Alamos, New Mexico, USA^c; DOE Joint Genome Institute, Walnut Creek, California, USA^d; Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA^e; Department of Land, Water and Air Resources, University of California-Davis, Davis, California, USA^f

The *Opitutaceae* bacterium strain TAV5, a member of the phylum *Verrucomicrobia*, was isolated from the wood-feeding termite hindgut. We report here its complete genome sequence, which contains a chromosome and a plasmid of 7,317,842 bp and 99,831 bp, respectively. The genomic analysis reveals genes for methyloph, lignocellulose degradation, and ammonia and sulfate assimilation.

Received 17 January 2015 Accepted 28 January 2015 Published 5 March 2015

Citation Kotak M, Isanapong J, Goodwin L, Bruce D, Chen A, Han CS, Huntemann M, Ivanova N, Land ML, Nolan M, Pati A, Woyke T, Rodrigues JLM. 2015. Complete genome sequence of the *Opitutaceae* bacterium strain TAV5, a potential facultative methyloph of the wood-feeding termite *Reticulitermes flavipes*. *Genome Announc* 3(2):e00060-15. doi:10.1128/genomeA.00060-15.

Copyright © 2015 Kotak et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/3.0/).

Address correspondence to Jorge L. M. Rodrigues, jmrodrigues@ucdavis.edu.

The ability to grow on single-carbon compounds other than CO₂ is a distinctive feature of methylophs. Among this group, members of the *Verrucomicrobia* have been identified as the first phylum outside of the *Proteobacteria* to be characterized as methanotrophs (1). The genomes of three species isolated from geographically distant but geothermally similar environments were previously sequenced and studied (2–4). Here, we report the complete genome of the *Opitutaceae* bacterium strain TAV5, a mesophilic verrucomicrobium isolated from the hindgut of the wood-feeding termite *Reticulitermes flavipes* (5), containing genes associated with methylophic competency.

The genomic DNA of TAV5 was isolated using a cetyltrimethylammonium bromide method, and the genome was sequenced using a combination of Illumina HiSeq 2000 and 454 FLX Titanium systems. The individual reads were assembled with the Newbler assembler (Roche). Prodigal (6) was used to identify the genes, and manual curation was done using the Joint Genome Institute (JGI) GenePRIMP pipeline (7). The translated protein-coding genes (coding sequences [CDSs]) were used to search the National Center for Biotechnology Information (NCBI) nonredundant, UniProt, TIGRFam, Pfam, PRIAM, KEGG, COG, and InterPro databases, and a product description for each protein was predicted. Noncoding genes and miscellaneous features were predicted using tRNAscan-SE (8), RNAmmer (9), Rfam (10), TM-HMM (6), and signalP (11).

The TAV5 genome is composed of a chromosome that is 7,317,842 bp long and contains 6,000 genes, with a G+C content of 63.45%, and a plasmid that is 99,831 bp long and contains 96 genes, with a G+C content of 54.44%. Together, they contain 5,950 CDSs, 48 tRNA genes, and 1 complete rRNA operon. The number of CDSs with a predicted function is 4,086 (67.03% of the total), the number of KEGG orthologs is 1,912 (31.36%), and the number of Cluster of Orthologous Group classifications is

2,984 (48.95%). An analysis of the genome revealed the presence of enzymes for formate metabolism, such as formate dehydrogenase, a formate transporter, and enzymes of the serine pathway, like serine hydroxymethyltransferase, glycerate kinase, and phosphoenolpyruvate carboxykinase.

The TAV5 genome contains a number of glycoside hydrolases (GH) involved in the degradation of cellulose (GH 5 and 9) and hemicellulose (GH 8, 10, 26, 28, and 53), as observed for the TAV1 and TAV2 genomes (12, 13). The genome has genes for the enzymes 3-carboxymuconate cyclase (EC 5.5.1.5) and 4-carboxymuconolactone decarboxylase (EC 4.1.1.44), which are involved in the degradation of protocatechuate that is derived from lignin (14), as well as genes coding for dioxygenases and dienelactone hydrolase (EC 3.1.1.45), known for ring cleavage of aromatic compounds. These enzymes structurally modify lignin, improving the accessibility of polysaccharides to glycoside hydrolases and increasing the efficiency of degradation (15). The TAV5 genome contains the *ccb₃*-type cytochrome oxidase gene, implying the role of TAV5 in oxygen removal. Furthermore, genes for ammonia and sulfate assimilation, the urea cycle, and nitrogen fixation were observed. The availability of this genome will allow the understanding of metabolic and physiological properties carried out by members of the phylum *Verrucomicrobia* present in the termite hindgut.

Nucleotide sequence accession numbers. The complete genome sequence of the *Opitutaceae* bacterium TAV5 was deposited in GenBank under the accession numbers CP007053.1 (chromosome) and CP007054.1 (plasmid).

ACKNOWLEDGMENT

The work conducted by the U.S. Department of Energy Joint Genome Institute is supported by the Office of Science of the U.S. Department of Energy under Contract DE-AC02-05CH11231.

REFERENCES

- Op den Camp HJ, Islam T, Stott MB, Harhangi HR, Hynes A, Schouten S, Jetten MS, Birkeland NK, Pol A, Dunfield PF. 2009. Environmental, genomic and taxonomic perspectives on methanotrophic *Verrucomicrobia*. *Environ Microbiol Rep* 1:293–306. <http://dx.doi.org/10.1111/j.1758-2229.2009.00022.x>.
- Dunfield PF, Yuryev A, Senin P, Smirnova AV, Stott MB, Hou S, Ly B, Saw JH, Zhou Z, Ren Y, Wang J, Mountain BW, Crowe MA, Weatherby TM, Bodelier PL, Liesack W, Feng L, Wang L, Alam M. 2007. Methane oxidation by an extremely acidophilic bacterium of the phylum *Verrucomicrobia*. *Nature* 450:879–882. <http://dx.doi.org/10.1038/nature06411>.
- Pol A, Heijmans K, Harhangi HR, Tedesco D, Jetten MS, Op den Camp HJ. 2007. Methanotrophy below pH 1 by a new *Verrucomicrobia* species. *Nature* 450:874–878. <http://dx.doi.org/10.1038/nature06222>.
- Islam T, Jensen S, Reigstad LJ, Larsen O, Birkeland NK. 2008. Methane oxidation at 55°C and pH 2 by a thermoacidophilic bacterium belonging to the *Verrucomicrobia* phylum. *Proc Natl Acad Sci U S A* 105:300–304. <http://dx.doi.org/10.1073/pnas.0704162105>.
- Stevenson BS, Eichorst SA, Wertz JT, Schmidt TM, Breznak JA. 2004. New strategies for cultivation and detection of previously uncultured microbes. *Appl Environ Microbiol* 70:4748–4755. <http://dx.doi.org/10.1128/AEM.70.8.4748-4755.2004>.
- Hyatt D, Chen GL, Locascio PF, Land ML, Larimer FW, Hauser LJ. 2010. Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics* 11:119. <http://dx.doi.org/10.1186/1471-2105-11-119>.
- Pati A, Ivanova NN, Mikhailova N, Ovchinnikova G, Hooper SD, Lykidis A, Kyrpides NC. 2010. GenePRIMP: a gene prediction improvement pipeline for prokaryotic genomes. *Nat Methods* 7:455–457. <http://dx.doi.org/10.1038/nmeth.1457>.
- Lowe TM, Eddy SR. 1997. tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res* 25:955–964. <http://dx.doi.org/10.1093/nar/25.5.0955>.
- Lagesen K, Hallin P, Rødland EA, Staerfeldt HH, Rognes T, Ussery DW. 2007. RNAMmer: consistent and rapid annotation of ribosomal RNA genes. *Nucleic Acids Res* 35:3100–3108. <http://dx.doi.org/10.1093/nar/gkm160>.
- Griffiths-Jones S, Bateman A, Marshall M, Khanna A, Eddy SR. 2003. Rfam: an RNA family database. *Nucleic Acids Res* 31:439–441. <http://dx.doi.org/10.1093/nar/gkg006>.
- Bendtsen JD, Nielsen H, von Heijne G, Brunak S. 2004. Improved prediction of signal peptides: SignalP 3.0. *J Mol Biol* 340:783–795. <http://dx.doi.org/10.1016/j.jmb.2004.05.028>.
- Isanapong J, Goodwin L, Bruce D, Chen A, Detter C, Han J, Han CS, Held B, Huntmann M, Ivanova N, Land ML, Mavromatis K, Nolan M, Pati A, Pennacchio L, Pitluck S, Szeto E, Tapia R, Woyke T, Rodrigues JL. 2012. High-quality draft genome sequence of the *Opi-tutaceae* bacterium strain TAV1, a symbiont of the wood-feeding termite *Reticulitermes flavipes*. *J Bacteriol* 194:2744–2745. <http://dx.doi.org/10.1128/JB.00264-12>.
- Isanapong J, Sealy Hambricht W, Willis AG, Boonmee A, Callister SJ, Burnum KE, Paša-Tolić L, Nicora CD, Wertz JT, Schmidt TM, Rodrigues JL. 2013. Development of an ecophysiological model for *Diplosphaera colotermitum* TAV2, a termite hindgut *Verrucomicrobium*. *ISME J* 7:1803–1813. <http://dx.doi.org/10.1038/ismej.2013.74>.
- Harwood CS, Parales RE. 1996. The β -ketoadipate pathway and the biology of self-identity. *Annu Rev Microbiol* 50:553–590. <http://dx.doi.org/10.1146/annurev.micro.50.1.553>.
- Brune A. 2014. Symbiotic digestion of lignocellulose in termite guts. *Nat Rev Microbiol* 12:168–180. <http://dx.doi.org/10.1038/nrmicro3182>.