

Complete Genome Sequence of *Terribacillus aidingensis* Strain MP602, a Moderately Halophilic Bacterium Isolated from *Cryptomeria fortunei* in Tianmu Mountain in China

Peng Lu,^a Mengying Lei,^a Fenghu Xiao,^b Liqin Zhang,^b  Yongjun Wang^{a,b}

Key Laboratory of Forest Protection, College of Forestry and Biotechnology, Zhejiang Agriculture and Forestry University, Lin'An, Zhejiang, China^a; National Joint Engineering Laboratory of Biopesticide Preparation, Lin'An, Zhejiang, China^b

P.L. and M.L. contributed equally to this work.

***Terribacillus aidingensis* strain MP602, which was isolated from an ancient tree (*Cryptomeria fortunei*) in Tianmu Mountain in China, has antagonistic activity against several certain phytopathogenic fungi. Here, we report the genome sequence of this strain. This is the first complete genome report of the *Terribacillus* genus.**

Received 1 February 2015 Accepted 4 February 2015 Published 19 March 2015

Citation Lu P, Lei M, Xiao F, Zhang L, Wang Y. 2015. Complete genome sequence of *Terribacillus aidingensis* strain MP602, a moderately halophilic bacterium isolated from *Cryptomeria fortunei* in Tianmu Mountain in China. *Genome Announc* 3(2):e00126-15. doi:10.1128/genomeA.00126-15.

Copyright © 2015 Lu et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](http://creativecommons.org/licenses/by/3.0/).

Address correspondence to Yongjun Wang, wangyj@zafu.edu.cn.

The genus *Terribacillus* belongs to the family *Bacillaceae* and was established in 2007 (1). Currently this genus comprises four validly published species: *Terribacillus saccharophilus* and *T. halophilus* were isolated from field soil in Japan (1); *T. goriensis* DSM 18252^T was isolated from coastal water of the east coast of Korea (2); and *T. aidingensis* Y17-61^T was isolated from sediments of Aiding Salt Lake in the Xinjiang region of China (3). It has been reported that *T. aidingensis* was efficient in biological control against gray mold in strawberry (4). *Terribacillus aidingensis* strain MP602 was isolated from an ~200-year-old tree (*Cryptomeria fortunei*) in Tianmu Mountain in China. This bacterium displays a moderate tolerance to salts (up to 14%). Interestingly, it showed high antagonistic activity against several plant pathogenetic fungi, such as *Botrytis cinerea*, *Magnaporthe grisea*, *Rhizoctonia solani*, and so on. Therefore, this bacterium has been evaluated as an inoculant for plant disease control. Genome sequencing of *T. aidingensis* MP602 was conducted to obtain additional insights into physiological characteristics involved in tolerance to salts and plant disease control and for future studies examining the molecular basis of these traits.

The completed genome sequence of *T. aidingensis* MP602 was developed at the Beijing Genomics Institute (BGI, Beijing, China) using Solexa paired-end sequencing technology (5). Using total genomic DNA, two gene libraries were prepared from sheared DNA fractions (~500 bp and ~2 kb) using Illumina paired-end sample preparation kits (Illumina, Inc.) according to the manufacturer's instructions. The DNA was sequenced using an Illumina Solexa GA IIx instrument. The sequencing run yielded 2,189,000 filtered paired-end reads (500-bp insert; 354 Mb in total) and a total of 3,925,000 reads for the 2-kb inserts (321 Mb in total) that were used for deep sequencing. The total sequence data provided 131-fold coverage of the genome. About 98.2% of the reads were assembled into 20 scaffolds with 251 contigs using the SOAPdenovo alignment tool (<http://soap.genomics.org.cn/index.html#intro2>). Gaps were

filled by Sanger sequencing of PCR products by custom primer walks and long-distance PCR amplification of the regions between each pair of scaffolds and contigs. The open reading frames (ORFs) were predicted using the NCBI Prokaryotic Genomes Automatic Annotation Pipeline (PGAAP) (6), and tRNA and rRNA genes were identified by tRNAscan-SE version 1.3 (7) and RNAmmer version 1.2 (8). The metabolic pathways were examined through the KEGG Automatic Annotation Server (<http://www.genome.jp/kegg>).

The completed genome of *T. aidingensis* MP602 contained two replicons, which comprised a circular chromosome (3,488,643 bp) containing 3,587 ORFs and a circular plasmid, which was named pT1 (88,439 bp), carrying 23 ORFs. The GC content of the chromosome is 42.76%, while the content of plasmid pT1 is 37.32%. This genome carried 56 tRNA genes and 24 rRNA operons, which are all located in the chromosome.

Nucleotide sequence accession numbers. The complete genome sequence of the *T. aidingensis* MP602 chromosome is available in GenBank under the accession number [CP008876](https://www.ncbi.nlm.nih.gov/nuccore/CP008876). The GenBank accession number for plasmid pT1 is [CP008877](https://www.ncbi.nlm.nih.gov/nuccore/CP008877).

ACKNOWLEDGMENTS

This work was supported by grant LY12C14006 from the Zhejiang Provincial Natural Science Foundation of China and grant KF201327 from the Open Fund for the Top-Priority-Discipline Forestry of Zhejiang province, Agriculture and Forestry University, China.

REFERENCES

1. An S-Y, Asahara M, Goto K, Kasai H, Yokota A. 2007. *Terribacillus saccharophilus* gen. nov., sp. nov. and *Terribacillus halophilus* sp. nov., spore-forming bacteria isolated from field soil in Japan. *Int J Syst Evol Microbiol* 57:51–55. <http://dx.doi.org/10.1099/ijs.0.64340-0>.
2. Kim Y-G, Hwang CY, Yoo KW, Moon HT, Yoon J-H, Cho BC. 2007. *Pelagibacillus goriensis* gen. nov., sp. nov., a moderately halotolerant bacterium isolated from coastal water off the east coast of Korea. *Int J Syst Evol Microbiol* 57:1554–1560. <http://dx.doi.org/10.1099/ijs.0.64820-0>.

3. Liu W, Jiang L, Guo C, Yang SS. 2010. *Terribacillus aidingensis* sp. nov., a moderately halophilic bacterium. *Int J Syst Evol Microbiol* 60:2940–2945. <http://dx.doi.org/10.1099/ijss.0.017228-0>.
4. Essghaier B, Fardeau ML, Cayol JL, Hajlaoui MR, Boudabous A, Jijakli H, Sadfi-Zouaoui N. 2009. Biological control of grey mould in strawberry fruits by halophilic bacteria. *J Appl Microbiol* 106:833–846. <http://dx.doi.org/10.1111/j.1365-2672.2008.04053.x>.
5. Bennett S. 2004. Solexa Ltd. *Pharmacogenomics* 5:433–438. <http://dx.doi.org/10.1517/14622416.5.4.433>.
6. Klimke W, Agarwala R, Badretdin A, Chetvernin S, Ciufu S, Fedorov B, Kiryutin B, O'Neill K, Resch W, Resenchuk S, Schafer S, Tolstoy I, Tatusova T. 2009. The National Center for Biotechnology Information's protein clusters database. *Nucleic Acids Res* 37:D216–D223. <http://dx.doi.org/10.1093/nar/gkn734>.
7. Lowe TM, Eddy SR. 1997. tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res* 25:955–964. <http://dx.doi.org/10.1093/nar/25.5.0955>.
8. Lagesen K, Hallin P, Rødland EA, Staerfeldt HH, Rognes T, Ussery DW. 2007. RNAmmer: consistent and rapid annotation of ribosomal RNA genes. *Nucleic Acids Res* 35:3100–3108. <http://dx.doi.org/10.1093/nar/gkm160>.