

Draft Genome Sequence of *Exophiala mesophila*, a Black Yeast with High Bioremediation Potential

Hakim Tafer, Ksenija Lopandic, Barbara Blasi, Caroline Poyntner, Katja Sterflinger

VIBT-Extremophile Center, University of Natural Resources and Life Sciences, Vienna, Austria

The fungal genus *Exophiala* comprises both pathogen species, which cause severe infections in humans, and environmental species, which are able to degrade alkylbenzene compounds. The draft genome sequence of *Exophiala mesophila* presented here is the first genome assembly of an alkylbenzene-degrading organism belonging to the genus *Exophiala*.

Received 11 February 2015 Accepted 24 February 2015 Published 9 April 2015

Citation Tafer H, Lopandic K, Blasi B, Poyntner C, Sterflinger K. 2015. Draft genome sequence of *Exophiala mesophila*, a black yeast with high bioremediation potential. *Genome Announc* 3(2):e00203-15. doi:10.1128/genomeA.00203-15.

Copyright © 2015 Tafer et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](http://creativecommons.org/licenses/by/3.0/).

Address correspondence to Katja Sterflinger, katja.sterflinger@boku.ac.at.

So-called black yeasts are ascomycetes exhibiting hyphal and yeast-like morphology with budding cells. Within the black yeasts, the genus *Exophiala* seems to be an evolutionary hot spot, with a high diversification and emerging adaptation toward human environments and human hosts. *Exophiala dermatitidis*, in particular, is both an opportunistic human pathogen and a fungus found in human environments, like bathrooms, sauna facilities, or dishwashers. Some species show a high affinity for contaminated sites and are able to degrade a variety of xenobiotics, including hydrocarbons, under desiccation and under extreme pH and temperature conditions. Thus, they are regarded as promising candidates for bioremediation (1). *Exophiala mesophila* was originally isolated from a silicone seal in a hospital but is also frequently found in tap water, saunas, and steam bath facilities (2).

Genomic DNA was isolated from *E. mesophila* (strain CBS 120910) grown on 2% malt extract agar using a cetyltrimethylammonium bromide (CTAB)-based protocol. The elimination of melanin from DNA was performed by two phenol-chloroform purification steps. Genome sequencing was carried out using the Ion Proton Technology (Ion AmpliSeq library preparation kit, Template OT2 200 kit, Ion PI sequencing 200 kit, and the Ion PI chip kit version 2; Life Technologies, Carlsbad, CA), according to the instructions of the producers. A total of 12.2 Gb, with a median read length of 169 bp, were assembled into a 30-Mb genome containing 507 contigs (N_{50} , 121,790). The genome assembly was performed with Newbler 2.9.

The presence of genes involved in toluene degradation was investigated in *E. mesophila* and, for the purposes of comparison, was also investigated in the yeast *Saccharomyces cerevisiae* S288C. A list of 12 genes related to toluene degradation was compiled from Parales et al. (3) and mapped to the three genomes using Scipio (4), with a score threshold of 0.3. Only 2 genes belonging to

the toluene degradation pathway (benzaldehyde dehydrogenase and benzylalcohol dehydrogenase) were found in yeast S288C, while 9 genes (ketoacyl-CoA thiolase, catechol 1,2-dioxygenase, protocatechuate dioxygenase, cytochrome P450, benzaldehyde dehydrogenase, benzylalcohol dehydrogenase, *cis,cis*-muconate-lactonizing enzyme, β -carboxy muconate enzyme, and β -ketoacyl-CoA transferase) were found in *E. mesophila*, in line with its ability to digest alkylbenzene compounds.

Nucleotide sequence accession numbers. This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession no. [JSEI000000000](https://www.ncbi.nlm.nih.gov/nuclink/JSEI000000000). The version described in this paper is [JSEI010000000](https://www.ncbi.nlm.nih.gov/nuclink/JSEI010000000).

ACKNOWLEDGMENTS

This project was financed by the Austrian Science Fund (FWF) (project P 25119-B16). The equipment of the VIBT-Extremophile Center used in this study was financed by BOKU-Equipment GesmbH. The computational results presented have been achieved in part using the Vienna Scientific Cluster (VSC).

REFERENCES

1. Prenafeta-Boldú FX, Guvernau M, Gallastegui G, Viñas M, de Hoog GS, Elías A. 2012. Fungal/bacterial interactions during the biodegradation of TEX hydrocarbons (toluene, ethylbenzene and *p*-xylene) in gas biofilters operated under xerophilic conditions. *FEMS Microbiol Ecol* 80:722–734. <http://dx.doi.org/10.1111/j.1574-6941.2012.01344.x>.
2. Listemann H, Freiesleben H. 1996. *Exophiala mesophila* spec. nov. *Mycoses* 39:1–3. <http://dx.doi.org/10.1111/j.1439-0507.1996.tb00076.x>.
3. Parales RE, Parales JV, Pelletier DA, Ditty JL. 2008. Diversity of microbial toluene degradation pathways. *Adv Appl Microbiol* 64:1–73. [http://dx.doi.org/10.1016/S0065-2164\(08\)00401-2](http://dx.doi.org/10.1016/S0065-2164(08)00401-2).
4. Keller O, Odronitz F, Stanke M, Kollmar M, Waack S. 2008. Scipio: using protein sequences to determine the precise exon/intron structures of genes and their orthologs in closely related species. *BMC Bioinformatics* 9:278. <http://dx.doi.org/10.1186/1471-2105-9-278>.