

Whole-Genome Shotgun Sequence of the Keratinolytic Bacterium *Lysobacter* sp. A03, Isolated from the Antarctic Environment

Jamile Queiroz Pereira,^{a,b} Adriana Ambrosini,^a Fernando Hayashi Sant'Anna,^a Michele Tadra-Sfeir,^c Helisson Faoro,^c Fábio Oliveira Pedrosa,^c Emanuel Maltempo Souza,^c Adriano Brandelli,^b Luciane M. P. Passaglia^a

Departamento de Genética, Instituto de Biociências, Universidade Federal do Rio Grande do Sul (UFRGS), Porto Alegre, RS, Brazil^a; Instituto de Ciências e Tecnologia de Alimentos, Universidade Federal do Rio Grande do Sul (ICTA-UFRGS), Porto Alegre, RS, Brazil^b; Departamento de Bioquímica e Biologia Molecular, Universidade Federal do Paraná (UFPR), Curitiba, PR, Brazil^c

***Lysobacter* sp. strain A03 is a protease-producing bacterium isolated from decomposing-penguin feathers collected in the Antarctic environment. This strain has the ability to degrade keratin at low temperatures. The A03 genome sequence provides the possibility of finding new genes with biotechnological potential to better understand its cold-adaptation mechanism and survival in cold environments.**

Received 19 February 2015 Accepted 23 February 2015 Published 2 April 2015

Citation Pereira JQ, Ambrosini A, Sant'Anna FH, Tadra-Sfeir M, Faoro H, Pedrosa FO, Souza EM, Brandelli A, Passaglia LMP. 2015. Whole-genome shotgun sequence of the keratinolytic bacterium *Lysobacter* sp. A03, isolated from the Antarctic environment. *Genome Announc* 3(2):e00246-15. doi:10.1128/genomeA.00246-15.

Copyright © 2015 Pereira et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/3.0/).

Address correspondence to Luciane M. P. Passaglia, luciane.passaglia@ufrgs.br.

Lysobacter is a genus of Gram-negative bacteria first described in 1978 (1) that belongs to the family *Xanthomonadaceae*, within the *Gammaproteobacteria*. They are characterized by gliding motility, a high G+C content, and the production of a broad range of proteases and antibiotics, thus representing a source of biocontrol agents (2).

The Antarctic strain A03 was isolated from decomposing-penguin feathers collected on King George Island, Antarctica, and was identified as a *Lysobacter* sp. by both 16S rRNA and 16S-23S rRNA intergenic transcribed spacer gene sequencing. The isolate was able to grow preferentially in feather meal broth (FMB) substrate and showed high proteolytic activity at temperatures of approximately 20°C, within the range of psychrophilic microorganisms (3). Considering its biotechnological potential, mainly due to its production of various extracellular proteases, the genome sequence of the *Lysobacter* sp. A03 strain was obtained and the preliminary analysis is presented here.

A03 whole-genome shotgun sequencing was performed on the MiSeq Illumina platform using the MiSeq reagent kit, version 2. A total of 109,889 paired-end reads were obtained, with an average length of 240 bp and approximately 18-fold coverage. The assembly was performed using CLC Genomics Workbench (<http://www.clcbio.com/products/clc-genomics-workbench/>), A5-miseq (4), CISA (5), and SPADES (6), and considering the lower N_{50} value (51,277) and the smaller number of contigs (101), the assembly constructed by SPADES was chosen. The CheckM (7) program was used to assess the quality of the microbial genome, and the automatic annotation of the genome sequence was performed in the RAST server (8).

The draft genome sequence of strain A03 comprised 2,873,548 bp representing approximately 99.1% of the genome size, with a G+C content of 65.79%. A total of 2,615 coding sequences (CDSs), 46 tRNA genes, and 2 rRNA genes were predicted. As expected, many peptidase-coding genes were found,

including a sequence coding for an extracellular keratinase; furthermore, three genes involved in cold shock response (2 *cspA* and 1 *cspG*) and six beta-lactamase resistance genes were predicted.

The availability of the genome sequence from *Lysobacter* sp. A03 may provide methods for searching for new biotechnologically relevant enzymes and might increase the understanding of the possible mechanisms that lead to the cold adaptation of life.

Nucleotide sequence accession numbers. This whole-genome shotgun project has been deposited at GenBank under the accession no. [JXSS000000000](https://www.ncbi.nlm.nih.gov/nuccore/JXSS000000000). The version described in this paper is version [JXSS010000000](https://www.ncbi.nlm.nih.gov/nuccore/JXSS010000000).

ACKNOWLEDGMENTS

This work was supported by the “Coordenação de Aperfeiçoamento de Pessoal de Nível Superior” (grant 42001013068D3) and “Conselho Nacional de Desenvolvimento Científico e Tecnológico—Instituto Nacional de Ciência e Tecnologia de Fixação Biológica do Nitrogênio” (project 573828/2008-3), Brazil.

We thank the “Projeto Antártico Brasileiro (PROANTAR)” and the “Laboratório de Ornitologia e Animais Marinhos” of the Universidade do Vale do Rio dos Sinos (UNISINOS) for providing the samples from which the strain was isolated.

REFERENCES

- Christensen P, Cook FD. 1978. *Lysobacter*, a new genus of nonfruiting, gliding bacteria with a high base ratio. *Int J Syst Evol Microbiol* 28:367–393. [http://dx.doi.org/10.1099/00207713-28-3-367](https://doi.org/10.1099/00207713-28-3-367).
- Zhou L, Li M, Yang J, Wei L, Ji G. 2014. Draft genome sequence of antagonistic agent *Lysobacter antibioticus* 13-6. *Genome Announc* 2(5):e00566-14. [http://dx.doi.org/10.1128/genomeA.00566-14](https://doi.org/10.1128/genomeA.00566-14).
- Pereira JQ, Lopes FC, Medina LFC, Petry MV, Brandelli A. 2014. Isolation of three novel Antarctic psychrotolerant feather-degrading bacteria and partial purification of keratinolytic enzyme from *Lysobacter* sp. A03. *Int Biodeterior Biodegr* 88:1–7. [http://dx.doi.org/10.1016/j.ibiod.2013.11.012](https://doi.org/10.1016/j.ibiod.2013.11.012).
- Coil D, Jospin G, Darling AE. 2014. A5-miseq: an updated pipeline to assemble microbial genomes from Illumina MiSeq data. *Bioinformatics* 31:587–589. [http://dx.doi.org/10.1093/bioinformatics/btu661](https://doi.org/10.1093/bioinformatics/btu661).

5. Lin SH, Liao YC. 2013. CISA: contig integrator for sequence assembly of bacterial genomes. *PLoS One* 8:e60843. <http://dx.doi.org/10.1371/journal.pone.0060843>.
6. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <http://dx.doi.org/10.1089/cmb.2012.0021>.
7. Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. 2014. CheckM: assessing the quality of microbial genomes recovered from isolates, single cells and metagenomes. *PeerJ PrePrints* 2:e554v1. <http://dx.doi.org/10.7287/peerj.preprints.554v1>.
8. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75. <http://dx.doi.org/10.1186/1471-2164-9-75>.