

Complete Genome Sequence of *Bifidobacterium dentium* Strain JCM 1195^T, Isolated from Human Dental Caries

Hidehiro Toh,^a Jun-ichiro Hayashi,^b Kenshiro Oshima,^c Akiko Nakano,^d Yukiko Takayama,^c Kageyasu Takanashi,^c Hidetoshi Morita,^e Masahira Hattori^c

Medical Institute of Bioregulation, Kyushu University, Fukuoka, Japan^a; School of Dentistry, Aichi Gakuin University, Nagoya, Japan^b; Graduate School of Frontier Sciences, University of Tokyo, Kashiwa, Japan^c; Department of Microbiology and Immunology, Teikyo University School of Medicine, Tokyo, Japan^d; School of Veterinary Medicine, Azabu University, Sagami-hara, Japan^e

H.T. and J.H. contributed equally to this work.

***Bifidobacterium dentium* strain JCM 1195^T was isolated from human dental caries. Here, we report the complete genome sequence of this organism.**

Received 25 February 2015 Accepted 26 February 2015 Published 9 April 2015

Citation Toh H, Hayashi J, Oshima K, Nakano A, Takayama Y, Takanashi K, Morita H, Hattori M. 2015. Complete genome sequence of *Bifidobacterium dentium* strain JCM 1195^T, isolated from human dental caries. *Genome Announc* 3(2):e00284-15. doi:10.1128/genomeA.00284-15.

Copyright © 2015 Toh et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Hidetoshi Morita, morita@azabu-u.ac.jp.

Bifidobacterium is frequently isolated from the human intestine, but *Bifidobacterium dentium* is known to be present in the human oral cavity, as well as in the intestine. *B. dentium* is frequently isolated from active carious lesions and thus may contribute to the pathogenesis of dental caries. Possession of an enzyme activity that degrades artificial trypsin substrates, such as benzoylarginine- β -naphthylamide, has been proposed as a possible virulence factor in the suspected periodontal pathogens (1, 2). *B. dentium*, which is one of the periodontal bacterial isolates, also has this activity (3). *B. dentium* belongs to the *Bifidobacterium adolescentis* group (4).

B. dentium strain JCM 1195^T (DSM 20436^T) was isolated from human dental caries (5). We determined the complete genome sequence of *B. dentium* JCM 1195^T using a whole-genome shotgun strategy with Sanger sequencing (ABI 3730xl sequencers). We constructed small-insert (2-kb) and large-insert (10-kb) genomic DNA libraries and generated 33,024 sequence reads (9.3-fold coverage) for *B. dentium* JCM 1195^T from both ends of the genomic clones. Data were assembled with the Phred-Phrap-Consed program. Gap closing and resequencing of low-quality regions were conducted by Sanger sequencing to obtain the high-quality finished sequence. The overall accuracy of the finished sequence was estimated to have an error rate of <1 per 10,000 bases (Phrap score of ≥ 40). An initial set of predicted protein-coding genes was identified using Glimmer version 3.0 (6). Genes consisting of <120 bp and those containing overlaps were eliminated. The tRNA genes were predicted by tRNAscan-SE (7), and the rRNA genes were detected by a BLASTn search using known *Bifidobacterium* rRNA sequences as queries.

The genome sequence of *B. dentium* JCM 1195^T consists of a circular chromosome of 2,635,669 bp with no plasmid. The genome size is larger than those of the other species in the *B. adolescentis* group, such as *B. adolescentis*, *B. angulatum*, *B. catenulatum*, and *B. pseudocatenulatum*. JCM 1195^T contained a clustered regularly interspaced short palindromic repeats (CRISPR) (8) region

(1,831,394 to 1,836,771), and five CRISPR-associated genes (BBDE_1555 to BBDE_1559) were encoded upstream of the CRISPR region. The chromosome contained 2,141 predicted protein-coding genes, 2,066 (97%) of which were conserved in the genome of *B. dentium* Bd1 (9). JCM 1195^T contained seven pilus gene clusters, all of which also were found in the genome of *B. dentium* Bd1 (10). Of the 2,141 protein-coding genes, 1,307 (61%) were conserved in the genome of *B. adolescentis* ATCC 15703^T (accession no. AP009256). The remaining 834 genes contained nine carbohydrate utilization gene clusters, which consist of a carbohydrate transporter, glycosyl hydrolase, and transcriptional regulator (BBDE_0114–BBDE_0129, BBDE_0465–BBDE_0469, BBDE_0627–BBDE_0633, BBDE_1007–BBDE_1010, BBDE_1208–BBDE_1212, BBDE_1611–BBDE_1616, BBDE_1980–BBDE_1986, BBDE_1996–BBDE_2043, and BBDE_2051–BBDE_2055). The genome information of this species will be useful for further studies of its physiology, taxonomy, and ecology.

Nucleotide sequence accession number. The sequence data for the genome have been deposited in DDBJ/GenBank/EMBL under the accession number [AP012326](https://doi.org/10.1101/012326).

ACKNOWLEDGMENTS

This research was supported by the Research Project Grant (2012-K-5) awarded by Azabu University and Grants-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science, and Technology of Japan (MEXT) (grant 24593135 to J.H.).

REFERENCES

1. Bedi GS, Williams T. 1994. Purification and characterization of a collagen-degrading protease from *Porphyromonas gingivalis*. *J Biol Chem* 269:599–606.
2. Pederson ED, Miller JW, Matheson S, Simonson LG, Chadwick DE, Covill PJ, Turner DW, Lamberts BL, Morton HE. 1994. Trypsin-like activity levels of *Treponema denticola* and *Porphyromonas gingivalis* in adults with periodontitis. *J Clin Periodontol* 21:519–525. <http://dx.doi.org/10.1111/j.1600-051X.1994.tb01167.x>.
3. Maiden MF, Tanner A, Macuch PJ. 1996. Rapid characterization of

- periodontal bacterial isolates by using fluorogenic substrate tests. *J Clin Microbiol* 34:376–384.
4. Felis GE, Dellaglio F. 2007. Taxonomy of lactobacilli and bifidobacteria. *Curr Issues Intest Microbiol* 8:44–61.
 5. Scardovi V, Crociani F. 1974. *Bifidobacterium catenulatum*, *Bifidobacterium dentium*, and *Bifidobacterium angulatum*: three new species and their deoxyribonucleic acid homology relationships. *Int J Syst Bacteriol* 24: 6–20. <http://dx.doi.org/10.1099/00207713-24-1-6>.
 6. Delcher AL, Bratke KA, Powers EC, Salzberg SL. 2007. Identifying bacterial genes and endosymbiont DNA with Glimmer. *Bioinformatics* 23:673–679. <http://dx.doi.org/10.1093/bioinformatics/btm009>.
 7. Schattner P, Brooks AN, Lowe TM. 2005. The tRNAscan-SE, snoscan and snoGPS Web servers for the detection of tRNAs and snoRNAs. *Nucleic Acids Res* 33:W686–W689. <http://dx.doi.org/10.1093/nar/gki366>.
 8. Horvath P, Coûté-Monvoisin AC, Romero DA, Boyaval P, Fremaux C, Barrangou R. 2009. Comparative analysis of CRISPR loci in lactic acid bacteria genomes. *Int J Food Microbiol* 131:62–70. <http://dx.doi.org/10.1016/j.ijfoodmicro.2008.05.030>.
 9. Ventura M, Turrioni F, Zomer A, Foroni E, Giubellini V, Bottacini F, Canchaya C, Claesson MJ, He F, Mantzourani M, Mulas L, Ferrarini A, Gao B, Delledonne M, Henrissat B, Coutinho P, Oggioni M, Gupta RS, Zhang Z, Beighton D, Fitzgerald GF, O'Toole PW, van Sinderen D. 2009. The *Bifidobacterium dentium* Bd1 genome sequence reflects its genetic adaptation to the human oral cavity. *PLoS Genet* 5:e1000785. <http://dx.doi.org/10.1371/journal.pgen.1000785>.
 10. Foroni E, Serafini F, Amidani D, Turrioni F, He F, Bottacini F, O'Connell Motherway M, Viappiani A, Zhang Z, Rivetti C, van Sinderen D, Ventura M. 2011. Genetic analysis and morphological identification of pilus-like structures in members of the genus *Bifidobacterium*. *Microb Cell Fact* 10(Suppl 1):S16. <http://dx.doi.org/10.1186/1475-2859-10-S1-S16>.