

Draft Genome Sequence of Antarctic *Pseudomonas* sp. Strain KG01 with Full Potential for Biotechnological Applications

María S. Pavlov,^a Felipe Lira,^b José L. Martínez,^b Jorge Olivares,^a Sergio H. Marshall^{a,c}

Laboratorio de Genética e Inmunología Molecular, Instituto de Biología, Facultad de Ciencias, Pontificia Universidad Católica de Valparaíso, Valparaíso, Chile^a; Departamento de Biotecnología Microbiana, Centro Nacional de Biotecnología CSIC, Madrid, Spain^b; Fraunhofer Chile Research Foundation, Center For Systems Biotechnology, Santiago, Chile^c

M.S.P. and F.L. contributed equally to this work.

We report here the draft genome sequence of a free-living psychrotolerant, *Pseudomonas* sp. strain KG01, isolated from an Antarctic soil sample and displaying interesting antimicrobial and surfactant activities. The sequence is 6.3 Mb long and includes 5,648 predicted-coding sequences.

Received 6 July 2015 Accepted 10 July 2015 Published 20 August 2015

Citation Pavlov MS, Lira F, Martínez JL, Olivares J, Marshall SH. 2015. Draft genome sequence of Antarctic *Pseudomonas* sp. strain KG01 with full potential for biotechnological applications. *Genome Announc* 3(4):e00906-15. doi:10.1128/genomeA.00906-15.

Copyright © 2015 Pavlov et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/3.0/).

Address correspondence to Sergio H. Marshall, smarshal@ucv.cl.

The genus *Pseudomonas*, described as early as 1894, is at present considered one of the most diverse and ubiquitous bacterial genera whose species have been isolated worldwide in all kinds of environments, from the poles to the tropics (1). The genus comprises a large number of strains able to produce a complex array of secondary metabolites with biotechnological relevance (2, 3). The recent technical advances in the genomic and metabolomic fields combined with sequencing of Antarctic microorganisms constitute a pivotal tool to augment traditional methods of studying and discovering valuable natural products (4). This *in silico* approach has successfully been demonstrated in the discovery of new antibiotics and plant growth-promoting substances produced by several *Pseudomonas* spp. isolated from temperate-latitude environments (5).

In this framework, we present here the draft sequence of the genome of *Pseudomonas* sp. strain KG01, a strain isolated from Antarctic soil with a full diversity of relevant bioactivities, such as surfactant production and antimicrobial compound secretion (M. S. Pavlov, unpublished data).

To access the genomic information, DNA was extracted using the Gnome DNA isolation kit (MPI Biomedicals, CA) from an overnight bacterial culture and subjected to whole-genome shotgun sequencing. Paired-end libraries, with an average insert size of 300 to 400 bp, were generated using a read length of 2×300 bp. Sequencing was performed using Illumina technology, according to the manufacturer's protocols for Illumina MiSeq at the facilities of Parque Científico de Madrid (Madrid, Spain). Shotgun sequencing produced a total of $2,054,712 \times 2$ reads. Raw sequences were filtered, and high-quality reads were assembled *de novo* using the MIRA 4.0 software (6) with the parameters established for the correct orientation and insert size determination. The resulting contigs were manually reassembled using Gap5 (Staden package version 2.0.0 b10) (7) to increase their length. The final KG01 assembly included 48 contigs (88.8-fold coverage), with the largest contig measuring 0.70 Mb. The draft genome sequence of *Pseu-*

domonas sp. strain KG01 comprises a total consensus sequence with an estimated size of 6.3 Mb and 60.1% G+C content.

Gene prediction and annotation were performed using the RAST server (<http://rast.nmpdr.org/>) (8). Using this approach, 5,648 coding sequences and 59 tRNA genes distributed in 16 scaffolds in the genome of *Pseudomonas* sp. strain KG01 were predicted.

A pyocin-like prophage sequence integrated between *mutS* and *recA-recX* (contig 1), two previously described siderophore biosynthetic gene clusters for *Pseudomonas* (pyoverdine, contig 2; pseudomonine, contig 161), an aryl-polyene gene cluster (contig 7), and an ectoine production gene cluster (contig 9) were identified using the antiSMASH server. In addition to previously *in vitro*-acquired evidences, the actual genomic information suggests a putative high potential of *Pseudomonas* sp. strain KG01 to produce biotechnological valuable substances with applications in different fields of biotechnology.

Nucleotide sequence accession numbers. This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession no. LFMW00000000. The version described in this paper is version LFMW01000000.

ACKNOWLEDGMENTS

This research was partially funded by CORFO-INACH grant 11IDL2-10651 and INACH grant DT_02-13. M.S.P. and F.L. were recipients of a CONICYT and La Caixa fellowship, respectively.

REFERENCES

1. Peix A, Ramírez-Bahena MH, Velázquez E. 2009. Historical evolution and current status of the taxonomy of genus *Pseudomonas*. *Infect Genet Evol* 9:1132–1147. <http://dx.doi.org/10.1016/j.meegid.2009.08.001>.
2. Gross H, Loper JE. 2009. Genomics of secondary metabolite production by *Pseudomonas* spp. *Nat Prod Rep* 26:1408–1446. <http://dx.doi.org/10.1039/b817075b>.

3. Raaijmakers JM, Mazzola M. 2012. Diversity and natural functions of antibiotics produced by beneficial and plant pathogenic bacteria. *Annu Rev Phytopathol* 50:403–424. <http://dx.doi.org/10.1146/annurev-phyto-081211-172908>.
4. Harvey AL, Edrada-Ebel R, Quinn RJ. 2015. The re-emergence of natural products for drug discovery in the genomics era. *Nat Rev Drug Discov* 14:111–129. <http://dx.doi.org/10.1038/nrd4510>.
5. Loper JE, Hassan KA, Mavrodi DV, Davis EW, Lim CK, Shaffer BT, Elbourne LD, Stockwell VO, Hartney SL, Breakwell K, Henkels MD, Tetu SG, Rangel LI, Kidarsa TA, Wilson NL, van de Mortel JE, Song C, Blumhagen R, Radune D, Hostetler JB, Brinkac LM, Durkin AS, Kluepfel DA, Wechter WP, Anderson AJ, Kim YC, Pierson LS, Pierson EA, Lindow SE, Kobayashi DY, Raaijmakers JM, Weller DM, Thomashow LS, Allen AE, Paulsen IT. 2012. Comparative genomics of plant-associated *Pseudomonas* spp.: insights into diversity and inheritance of traits involved in multitrophic interactions. *PLoS Genet* 8:e1002784. <http://dx.doi.org/10.1371/journal.pgen.1002784>.
6. Chevreux B, Wetter T, Suhai S. 1999. Genome sequence assembly using trace signals and additional sequence information, p 45–56. *In* Computer science and biology. Proceedings of the German Conference on Bioinformatics, GCB '99 GCB, Hannover, Germany.
7. Bonfield JK, Whitwham A. 2010. Gap5—editing the billion fragment sequence assembly. *Bioinformatics* 26:1699–1703. <http://dx.doi.org/10.1093/bioinformatics/btq268>.
8. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75. <http://dx.doi.org/10.1186/1471-2164-9-75>.