

Complete Genome Sequence of the Fish Pathogen *Flavobacterium columnare* Strain C#2

Ryan P. Bartelme,^a Ryan J. Newton,^a Yongtao Zhu,^b Nan Li,^c Benjamin R. LaFrentz,^d Mark J. McBride^b

School of Freshwater Sciences, University of Wisconsin—Milwaukee, Milwaukee, Wisconsin, USA^a; Department of Biological Sciences, University of Wisconsin—Milwaukee, Milwaukee, Wisconsin, USA^b; Institute of Hydrobiology, Chinese Academy of Sciences, Wuhan, Hubei Province, China^c; United States Department of Agriculture-Agricultural Research Service (USDA-ARS), Aquatic Animal Health Research Unit, Auburn, Alabama, USA^d

***Flavobacterium columnare* is a Gram-negative bacterial pathogen that causes columnaris disease of freshwater fish. *Flavobacterium columnare* strain C#2 was isolated from a diseased warm-water fish and is typed as genomovar II. The genome consists of a single 3.33-Mb circular chromosome with 2,689 predicted coding genes.**

Received 10 May 2016 Accepted 13 May 2016 Published 23 June 2016

Citation Bartelme RP, Newton RJ, Zhu Y, Li N, LaFrentz BR, McBride MJ. 2016. Complete genome sequence of the fish pathogen *Flavobacterium columnare* strain C#2. Genome Announc 4(3):e00624-16. doi:10.1128/genomeA.00624-16.

Copyright © 2016 Bartelme et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Ryan P. Bartelme, bartelm7@uwm.edu.

Flavobacterium columnare is an important pathogen of freshwater fish (1–3). Although large die-offs among wild and farmed fish are caused by *F. columnare*, little is known regarding its virulence mechanisms or its ecology (4). Strains of *F. columnare* are classified into genomovars based on restriction digestion of the 16S rRNA gene and the 16S–23S rRNA internal transcribed spacer (2, 3), and *F. columnare* strain C#2 (originally referred to as *F. columnare* strain #2 [1, 5]) is a genomovar II strain. The genome sequence of a genomovar I strain was previously reported (6). *F. columnare* strain C#2 was selected for genome sequencing because it is a virulent genomovar II strain and is amenable to genetic manipulations (5).

A single colony of strain C#2 was grown overnight in 10 ml of modified Shieh medium (7) at 25°C with shaking. Genomic DNA (gDNA) was extracted from 10 ml of culture using the Qiagen DNeasy blood and tissue kit (Qiagen, Hilden, Germany). AMPure bead size-selected 20-kb libraries were constructed according to the Pacific Biosciences RSII protocol. Two PacBio single-molecule real-time (SMRT) cells were loaded with 0.01 and 0.015 nM concentrations of library with Pacific Biosciences sequencing reagent 4.0, C4 chemistry, and P6 version 2 polymerase. Sequencing produced 66,158 reads, with an N_{50} read length of 44,216 bp and mean read length of 20,496 bp, representing 296.7× coverage of the genome. Genome assembly was done using the PacBio PBcR HGAP 2.3.0 pipeline, with default settings (8). The assembly produced two contigs of 3,330,796 and 8,000 bases. The shorter 8-kb contig was eliminated, as the confidence scoring was too low for inclusion in the assembly (Quality score, mapQV < 10). Gepard comparison (9) of the *F. columnare* strain C#2 genome to that of *F. columnare* 94-081 (also genomovar II; accession no. CP013992), revealed three large syntenic regions split by a single chromosomal inversion and rearrangement. Similar chromosomal changes have been observed in other members of the genus *Flavobacterium* (10). The polished assembly was trimmed using the check_circularity.pl script from SPRAI (11), and the resulting

single circular contig was annotated using the Prokaryotic Genome Annotation Pipeline through NCBI (12).

The C#2 genome has a G+C content of 30.97%, 13 rRNA operons, 93 tRNAs, and 2 clustered regularly interspaced short palindromic repeat (CRISPR) arrays, and 93.4% of the open reading frames (ORFs) correspond to predicted coding genes. The genome was analyzed for secretion systems and potential secreted virulence factors (13, 14). *F. columnare* strain C#2 contains the core genes of the *Bacteroidetes*-specific type IX secretion system (15). Genes encoding potential secreted virulence factors, such as chondroitinases, proteases, and adhesins, were also identified. *F. columnare* strain C#2 exhibits gliding motility, which is potentially important in *columnaris* pathology, and the annotated genome contains all of the genes known to be required for this process (16, 17). The availability of a complete genome and the genetic manipulability of *F. columnare* strain C#2 enable further work toward understanding *F. columnare* virulence factors and the development of avirulent vaccine strains to prevent outbreaks of columnaris disease in aquaculture settings.

Nucleotide sequence accession number. The genome sequence of *Flavobacterium columnare* strain C#2 has been deposited in GenBank with the accession no. CP015107. This paper describes the first version of the genome deposited.

ACKNOWLEDGMENTS

We thank the Great Lakes Genomics Center (School of Freshwater Sciences, University of Wisconsin Milwaukee) for their assistance in library preparation, sequencing, and use of the PacBio SMRT Portal. We thank Andrew Goodwin, Swapna Thomas-Jinu, Andrew Staroscik, and David Hunnicutt for advice regarding nomenclature for strain C#2. We thank Sandra L. McLellan for her contributions optimizing gDNA extraction and acquiring some of the funding provided for this work.

FUNDING INFORMATION

This work, including the efforts of Ryan J. Newton, was funded by a University of Wisconsin System Incentive grant to the School of Freshwater Sciences. This work, including the efforts of Mark J. McBride, was

funded by the Wisconsin Sea Grant Institute, University of Wisconsin (519K805), and by USDA CRIS project number 5090-31320-002-00D.

REFERENCES

1. Thomas-Jinu S, Goodwin AE. 2004. Morphological and genetic characteristics of *Flavobacterium columnare* isolates: correlations with virulence in fish. *J Fish Dis* 27:29–35. <http://dx.doi.org/10.1046/j.1365-2761.2003.00507.x>.
2. Darwish AM, Ismaiel AA. 2005. Genetic diversity of *Flavobacterium columnare* examined by restriction fragment length polymorphism and sequencing of the 16S ribosomal RNA gene and the 16S-23S rDNA spacer. *Mol Cell Probes* 19:267–274. <http://dx.doi.org/10.1016/j.mcp.2005.04.003>.
3. LaFrentz BR, Waldbieser GC, Welch TJ, Shoemaker CA. 2014. Intra-genomic heterogeneity in the 16S rRNA genes of *Flavobacterium columnare* and standard protocol for genomovar assignment. *J Fish Dis* 37:657–669. <http://dx.doi.org/10.1111/jfd.12166>.
4. McBride MJ. 2014. The family *Flavobacteriaceae*, p 643–676. In Rosenberg E, DeLong E, Lory S, Stackebrandt E, Thompson F (ed), *The prokaryotes*, vol 11, 4th ed. Springer Berlin Heidelberg, Berlin, Germany.
5. Staroscik AM, Hunnicutt DW, Archibald KE, Nelson DR. 2008. Development of methods for the genetic manipulation of *Flavobacterium columnare*. *BMC Microbiol* 8:115. <http://dx.doi.org/10.1186/1471-2180-8-115>.
6. Tekedar HC, Karsi A, Gillaspay AF, Dyer DW, Benton NR, Zaitshik J, Vamenta S, Banes MM, Gülsoy N, Aboko-Cole M, Waldbieser GC, Lawrence ML. 2012. Genome sequence of the fish pathogen *Flavobacterium columnare* ATCC 49512. *J Bacteriol* 194:2763–2764. <http://dx.doi.org/10.1128/JB.00281-12>.
7. Decostere A, Haesebrouck F, Devriese LA. 1997. Shieh medium supplemented with tobramycin for selective isolation of *Flavobacterium columnare* (*Flexibacter columnaris*) from diseased fish. *J Clin Microbiol* 35:322–324.
8. Chin C-S, Alexander DH, Marks P, Klammer AA, Drake J, Heiner C, Clum A, Copeland A, Huddleston J, Eichler EE, Turner SW, Korlach J. 2013. Nonhybrid, finished microbial genome assemblies from long-read SMRT sequencing data. *Nat Methods* 10:563–569. <http://dx.doi.org/10.1038/nmeth.2474>.
9. Krumsiek J, Arnold R, Rattei T. 2007. Gepard: a rapid and sensitive tool for creating dotplots on genome scale. *Bioinformatics* 23:1026–1028. <http://dx.doi.org/10.1093/bioinformatics/btm039>.
10. Touchon M, Barbier P, Bernardet JF, Loux V, Vacherie B, Barbe V, Rocha EPC, Duchaud E. 2011. Complete genome sequence of the fish pathogen *Flavobacterium branchiophilum*. *Appl Environ Microbiol* 77:7656–7662. <http://dx.doi.org/10.1128/AEM.05625-11>.
11. Imai T. 2015. SPRAI: single pass read accuracy improver, version 0.9.9.10. Sprai, Tokyo, Japan.
12. Tatusova T, Ciufu S, Federhen S, Fedorov B, McVeigh R, O'Neill K, Tolstoy I, Zaslavsky L. 2015. Update on RefSeq microbial genomes resources. *Nucleic Acids Res* 43:D599–D605. <http://dx.doi.org/10.1093/nar/gku1062>.
13. Chagnot C, Zorgani MA, Astruc T, Desvaux M. 2013. Proteinaceous determinants of surface colonization in bacteria: bacterial adhesion and biofilm formation from a protein secretion perspective. *Front Microbiol* 4:303. <http://dx.doi.org/10.3389/fmicb.2013.00303>.
14. Sandkvist M. 2001. Type II secretion and pathogenesis. *Infect Immun* 69:3523–3535. <http://dx.doi.org/10.1128/IAI.69.6.3523-3535.2001>.
15. Abby SS, Cury J, Guglielmini J, Néron B, Touchon M, Rocha EPC. 2016. Identification of protein secretion systems in bacterial genomes. *Sci Rep* 6:23080. <http://dx.doi.org/10.1038/srep23080>.
16. Klesius PH, Shoemaker CA, Evans JJ. 2008. *Flavobacterium columnare* chemotaxis to channel catfish mucus. *FEMS Microbiol Lett* 288:216–220. <http://dx.doi.org/10.1111/j.1574-6968.2008.01348.x>.
17. McBride MJ, Zhu Y. 2013. Gliding motility and Por secretion system genes are widespread among members of the phylum *Bacteroidetes*. *J Bacteriol* 195:270–278. <http://dx.doi.org/10.1128/JB.01962-12>.