

# Complete Genome Sequence of *Bacillus subtilis* subsp. *subtilis* Strain $\Delta 6$

Daniel R. Reuß,<sup>a</sup> Andrea Thürmer,<sup>b</sup> Rolf Daniel,<sup>b</sup> Wim J. Quax,<sup>c</sup> Jörg Stülke<sup>a</sup>

Department of General Microbiology, Georg-August-University Göttingen, Göttingen, Germany<sup>a</sup>; Department of Genomics and Applied Microbiology and Göttingen Genomics Laboratory, Georg-August-University Göttingen, Göttingen, Germany<sup>b</sup>; Groningen Research Institute of Pharmacy, University of Groningen, Groningen, The Netherlands<sup>c</sup>

***Bacillus subtilis*  $\Delta 6$  is a genome-reduced strain that was cured from six prophages and AT-rich islands. This strain is of great interest for biotechnological applications. Here, we announce the full-genome sequence of this strain. Interestingly, the conjugative element ICEBs1 has most likely undergone self-excision in *B. subtilis*  $\Delta 6$ .**

Received 6 June 2016 Accepted 10 June 2016 Published 28 July 2016

Citation Reuß DR, Thürmer A, Daniel R, Quax WJ, Stülke J. 2016. Complete genome sequence of *Bacillus subtilis* subsp. *subtilis* strain  $\Delta 6$ . *Genome Announc* 4(4):e00759-16. doi:10.1128/genomeA.00759-16.

Copyright © 2016 Reuß et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Jörg Stülke, [jstuelk@gwdg.de](mailto:jstuelk@gwdg.de).

*Bacillus subtilis*  $\Delta 6$  is a derivative of the laboratory wild-type strain *B. subtilis* 168, which was cured from six prophages and AT-rich islands. For this purpose, the prophages SP $\beta$  and PBSX, the prophage-like elements prophage 1, prophage 3, and *skin*, as well as the polyketide synthesis operon *pks* were deleted. Interestingly, this genome reduction by 7.7% did not have a major impact on physiology, metabolic flux patterns, or genetic competence (1).

*B. subtilis*  $\Delta 6$  is a promising starting point for further genome reduction. Moreover, it can serve as a chassis strain in the context of biotechnological applications, that is, highly efficient protein secretion and vitamin production (2–4). Indeed, *B. subtilis*  $\Delta 6$  has recently been used to obtain a total genome reduction of 13.6% (5). For a better understanding with respect to future projects, we have sequenced the genome of *B. subtilis*  $\Delta 6$ . The chromosomal DNA was isolated from a stationary phase culture using a commercially available kit (peqGOLD Bacterial DNA Kit, VWR International GmbH). We obtained 6.63 million reads from an Illumina 75-bp single-read run and mapped them to the *B. subtilis* 168 genome (GenBank accession number NC\_000964) (6) using the Geneious Read Mapper (Geneious version 9.0.5 software, Biomatters, Ltd.) (7). The alignment showed a 118-fold average coverage and a 99.5% pairwise identity to the reference genome of *B. subtilis* 168. The insertion and the correct sequence of the chloramphenicol resistance gene at the *pks* operon locus were verified by a standard PCR. The final genome sequence of *B. subtilis*  $\Delta 6$  has a length of 3,876,919 bp.

We identified 28 variations (single-nucleotide polymorphism, deletion, insertion, and substitution) with a minimal coverage of 25 $\times$  and a minimum variant frequency of 0.8. Four of these mutations have an effect on the amino acids sequence of the encoded protein (*carA*, *yobM*, *ywbD*, and *walH*), whereas four mutations are silent (*yczC*, *yjnA*, *glcF*, and *amyX*). The remaining 20 variants are located in intergenic and RNA-encoding regions. All variations can be requested from the corresponding author. In addition, we could confirm the presence of all six deletions performed by Westers et al. (1). Interestingly, *B. subtilis*  $\Delta 6$  contains a seventh

large deletion of 20.5 kb (25 genes; genome position: 529,422 to 549,925 bp). This deletion corresponds to the mobile genetic element ICEBs1 (8), which likely has undergone self-excision, as it has been reported for other *B. subtilis* strains (9). Taken together, *B. subtilis*  $\Delta 6$  is lacking 376 genes at seven different locations covering 8.03% of the reference genome of *B. subtilis* 168. These deletions increased the GC content from 43.5% to 43.9%.

**Nucleotide sequence accession number.** The genome sequence of *B. subtilis* subsp. *subtilis* strain  $\Delta 6$  is deposited in GenBank under the accession number CP015975.

## ACKNOWLEDGMENTS

D.R.R. was supported by the Göttingen Graduate School for Neuroscience and Molecular Biosciences (DFG grant GSC 226/2).

## FUNDING INFORMATION

This work, including the efforts of Daniel Reuß, was funded by Deutsche Forschungsgemeinschaft (DFG) (GSC 226/2).

D.R.R. was supported by the Göttingen Graduate School for Neuroscience and Molecular Biosciences (DFG grant GSC 226/2).

## REFERENCES

- Westers H, Dorenbos R, van Dijk JM, Kabel J, Flanagan T, Devine KM, Jude F, Seror SJ, Beekman AC, Darmon E, Eschevins C, de Jong A, Bron S, Kuipers OP, Albertini AM, Antelmann H, Hecker M, Zamboni N, Sauer U, Bruand C, Ehrlich DS, Alonso JC, Salas M, Quax WJ. 2003. Genome engineering reveals large dispensable regions in *Bacillus subtilis*. *Mol Biol Evol* 20:2076–2090. <http://dx.doi.org/10.1093/molbev/msg219>.
- Van Dijk JM, Hecker M. 2013. *Bacillus subtilis*: from soil bacterium to super-secreting cell factory. *Microb Cell Fact* 12:3. <http://dx.doi.org/10.1186/1475-2859-12-3>.
- Juhas M, Reuß DR, Zhu B, Commichau FM. 2014. *Bacillus subtilis* and *Escherichia coli* essential genes and minimal cell factories after one decade of genome engineering. *Microbiology* 160:2341–2351. <http://dx.doi.org/10.1099/mic.0.079376-0>.
- Commichau FM, Alzinger A, Sande R, Bretzel W, Meyer FM, Chevreux B, Wyss M, Hohmann HP, Prágai Z. 2014. Overexpression of a non-native deoxyxylulose-dependent vitamin B<sub>6</sub> pathway in *Bacillus subtilis* for the

- production of pyridoxine. *Metag Eng* 25:38–49. <http://dx.doi.org/10.1016/j.jymben.2014.06.007>.
5. Wenzel M, Altenbuchner J. 2015. Development of a markerless gene deletion system for *Bacillus subtilis* based on the mannose phosphoenolpyruvate-dependent phosphotransferase system. *Microbiology* 161:1942–1949. <http://dx.doi.org/10.1099/mic.0.000150>.
  6. Barbe V, Cruveiller S, Kunst F, Lenoble P, Meurice G, Sekowska A, Vallenet D, Wang T, Moszer I, Médigue C, Danchin A. 2009. From a consortium sequence to a unified sequence: the *Bacillus subtilis* 168 reference genome a decade later. *Microbiology* 155:1758–1775. <http://dx.doi.org/10.1099/mic.0.027839-0>.
  7. Kears M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Duran C, Thierer T, Ashton B, Meintjes P, Drummond A. 2012. Geneious basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 28:1647–1649. <http://dx.doi.org/10.1093/bioinformatics/bts199>.
  8. Auchtung JM, Lee CA, Monson RE, Lehman AP, Grossman AD. 2005. Regulation of a *Bacillus subtilis* mobile genetic element by intercellular signaling and the global DNA damage response. *Proc Natl Acad Sci USA* 102:12554–12559. <http://dx.doi.org/10.1073/pnas.0505835102>.
  9. Reuß DR, Schuldes J, Daniel R, Altenbuchner J. 2015. Complete genome sequence of *Bacillus subtilis* subsp. *subtilis* strain 3NA. *Genome Announc* 3(2):e00084-15. <http://dx.doi.org/10.1128/genomeA.00084-15>.