

Draft Genome Sequence of *Curtobacterium* sp. Strain UCD-KPL2560 (Phylum *Actinobacteria*)

Brian A. Klein,^a Katherine P. Lemon,^{a,b} Lina L. Faller,^a Guillaume Jospin,^c Jonathan A. Eisen,^{c,d} David A. Coil^c

Department of Microbiology, The Forsyth Institute, Cambridge, Massachusetts, USA^a; Division of Infectious Diseases, Boston Children's Hospital, Harvard Medical School, Boston, Massachusetts, USA^b; Genome Center, University of California Davis, Davis, California, USA^c; Department of Evolution and Ecology and Department of Medical Microbiology and Immunology, University of California Davis, Davis, California, USA^d

Here, we present the draft genome sequence of the actinobacterium *Curtobacterium* sp. strain UCD-KPL2560, which was isolated from the running surface of an indoor track field house in Medford, MA, USA (42.409716°N, -71.115169°W). The genome assembly contains 3,480,487 bp in 156 contigs.

Received 9 August 2016 Accepted 17 August 2016 Published 6 October 2016

Citation Klein BA, Lemon KP, Faller LL, Jospin G, Eisen JA, Coil DA. 2016. Draft genome sequence of *Curtobacterium* sp. strain UCD-KPL2560 (phylum *Actinobacteria*). *Genome Announc* 4(5):e01040-16. doi:10.1128/genomeA.01040-16.

Copyright © 2016 Klein et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Katherine P. Lemon, klemon@forsyth.org.

Members of the genus *Curtobacterium*, which was accepted as a genus in 1972 and refined in 1986 (1, 2), have been previously isolated from soil (3, 4), plants (5), cheese-processing equipment (6), and residential carpet (7). *Curtobacterium* spp. are characterized as Gram-positive aerobic bacilli that frequently demonstrate yellow-orange pigment when grown *in vitro* (2). *Curtobacterium* spp. are not common pathogens of humans; however, *Curtobacterium* spp. are recognized plant pathogens (8), and rare plant-to-human infections, as well as occasional human host isolation, are reported (9, 10).

We isolated *Curtobacterium* sp. strain UCD-KPL2560 from the flooring of an indoor track facility in Medford, MA, on 15 October 2015 as part of a project to produce reference genomes for microorganisms residing in the built environment. A nylon-flocked swab (Copan) dipped in sterile buffer (0.1 M NaCl and 0.1% Tween) was rubbed over the track surface starting block area (5 cm²) for 1 min, inoculated onto brain heart infusion agar containing fosfomycin (20 µg/ml), and incubated aerobically at 37°C for five days. A small round orange colony was selected from the original isolation plate and subcultured for purity. For 16S rRNA identification, an initial DNA extraction was performed with the DNeasy purification kit (Qiagen). A putative genus name was assigned to the isolate following PCR amplification (primers 27F and 1492R) and subsequent Sanger sequencing of the 16S rRNA gene. Genomic DNA for whole-genome sequencing was extracted using the MasterPure complete DNA/RNA purification kit (Epicentre).

Illumina paired-end libraries were generated using a Nextera DNA sample prep kit (Illumina). We selected 600- to 900-bp fragments using a Pippin Prep (Sage Science). Resultant libraries were sequenced on an Illumina MiSeq, with a read length of 300 bp. This produced a total of 3,920,338 paired-end reads. Quality trimming and error correction of the reads resulted in 3,506,299 high-quality reads using the A5-MiSeq assembly pipeline (version 05/22/2015) (11, 12). The resulting assembly contained 156 scaffolds (minimum, 598 bp; maximum, 215,854 bp; N_{50} , 36,322 bp). The

final assembly contained 3,480,487 bp, with a G+C content of 72% and a coverage estimate of 114% (EC value). Genome completeness was assessed using PhyloSift and CheckM (13, 14); all 37 PhyloSift marker genes were present, with 36 in single copy, and the ribosomal subunit L14 present in four copies. CheckM gave a 99% completeness estimation.

Annotation was performed using the RAST server (default settings on 14 April 2016) (15). *Curtobacterium* sp. strain UCD-KPL2560 contains 3,201 predicted coding sequences (CDSs), four predicted rRNAs, and 53 predicted tRNAs. It has 12 type IV pilus genes, 34 motility genes, and four auxin biosynthesis genes. At least two partial phages are predicted, and no clustered regularly interspaced short palindromic repeat (CRISPR) systems were identified (15–17).

We attempted to assign a putative species designation to *Curtobacterium* sp. UCD-KPL2560 using PhyloPhlAn and by generating a 16S-rRNA-gene-based phylogeny of *Curtobacterium* species curated by the Ribosomal Database Project (18, 19). However, neither yielded an assignment. PhyloPhlAn generated “*Clavibacter michiganensis*” taxonomic assignments for 16 of 17 *Curtobacterium* spp., with available NCBI sequencing data (one reclassified as *Rothia*). Additionally, *Curtobacterium* sp. UCD-KPL2560 did not fall within a well-supported clade of a single species within the genus *Curtobacterium* using 16S rRNA alone.

Accession number(s). This whole-genome shotgun project has been deposited at DDBJ/ENA/GenBank under the accession no. **MCIG00000000**. The version described in this paper is version MCIG01000000.

ACKNOWLEDGMENTS

We thank Tufts University for access to the Gantcher Indoor Track Center facility and promotion of facilities and microbial research. Sequencing was performed at the DNA Technologies Core facility in the Genome Center at the UC Davis, Davis, CA.

This work was funded in part by a grant (to J.A.E.) and postdoctoral

fellowship (to B.A.K.) from the Alfred P. Sloan Foundation as part of their program on the Microbiology of the Built Environment.

FUNDING INFORMATION

This work, including the efforts of Katherine P. Lemon, was funded by HHS | NIH | National Institute of Allergy and Infectious Diseases (NIAID) (R01 AI101018). This work, including the efforts of Brian A. Klein and Jonathan Eisen, was funded by Alfred P. Sloan Foundation.

REFERENCES

1. Yamada K, Komagata K. 1972. Taxonomic studies on coryneform bacteria V. Classification of coryneform bacteria. *J Gen Appl Microbiol* 18: 417–431. <http://dx.doi.org/10.2323/jgam.18.417>.
2. Komagata K, Suzuki K. 1986. Genus *Curtobacterium*, p 1313–1317. In Bergey's manual of systematic bacteriology. Williams & Wilkins, Philadelphia, PA.
3. Kim MK, Kim YJ, Kim HB, Kim SY, Yi TH, Yang DC. 2008. *Curtobacterium ginsengisoli* sp. nov., isolated from soil of a ginseng field. *Int J Syst Evol Microbiol* 58:2393–2397. <http://dx.doi.org/10.1099/ijs.0.65591-0>.
4. Mariita RM, Bhatnagar S, Hanselmann K, Hossain MJ, Korlach J, Boitano M, Roberts RJ, Liles MR, Moss AG, Leadbetter JR, Newman DK, Dawson SC. 2015. Complete genome sequence of *Curtobacterium* sp. strain MR_MD2014, isolated from topsoil in Woods Hole, Massachusetts. *Genome Announc* 3(6):e01504-15. <http://dx.doi.org/10.1128/genomeA.01504-15>.
5. Bulgari D, Minio A, Casati P, Quaglino F, Delledonne M, Bianco PA. 2014. *Curtobacterium* sp. genome sequencing underlines plant growth promotion-related traits. *Genome Announc* 2(4):e00592-14. <http://dx.doi.org/10.1128/genomeA.00592-14>.
6. Didiene R, Defargues C, Callon C, Meylheuc T, Hulin S, Montel MC. 2012. Characteristics of microbial biofilm on wooden vats (“gerles”) in PDO Salers cheese. *Int J Food Microbiol* 156:91–101. <http://dx.doi.org/10.1016/j.ijfoodmicro.2012.03.007>.
7. Flanagan JC, Lang JM, Darling AE, Eisen JA, Coil DA. 2013. Draft genome sequence of *Curtobacterium flaccumfaciens* strain UCD-AKU (phylum *Actinobacteria*). *Genome Announc* 1(3):e00244-13. <http://dx.doi.org/10.1128/genomeA.00244-13>.
8. Agarkova IV, Lambrecht PA, Vidaver AK, Harveson RM. 2012. Genetic diversity among *Curtobacterium flaccumfaciens* pv. *flaccumfaciens* populations in the American high plains. *Can J Microbiol* 58:788–801. <http://dx.doi.org/10.1139/w2012-052>.
9. Francis MJ, Doherty RR, Patel M, Hamblin JF, Ojaimi S, Korman TM. 2011. *Curtobacterium flaccumfaciens* septic arthritis following puncture with a Coxspur hawthorn thorn. *J Clin Microbiol* 49:2759–2760. <http://dx.doi.org/10.1128/JCM.00340-11>.
10. Funke G, Aravena-Roman M, Frodl R. 2005. First description of *Curtobacterium* spp. isolated from human clinical specimens. *J Clin Microbiol* 43:1032–1036. <http://dx.doi.org/10.1128/JCM.43.3.1032-1036.2005>.
11. Tritt A, Eisen JA, Facciotti MT, Darling AE. 2012. An integrated pipeline for *de novo* assembly of microbial genomes. *PLoS One* 7:e42304. <http://dx.doi.org/10.1371/journal.pone.0042304>.
12. Coil D, Jospin G, Darling AE. 2015. A5-miseq: an updated pipeline to assemble microbial genomes from Illumina MiSeq data. *Bioinformatics* 31:587–589. <http://dx.doi.org/10.1093/bioinformatics/btu661>.
13. Darling AE, Jospin G, Lowe E, Matsen FA, Bik HM, Eisen JA. 2014. PhyloSift: phylogenetic analysis of genomes and metagenomes. *PeerJ* 2:e243. <http://dx.doi.org/10.7717/peerj.243>.
14. Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. 2015. CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes *Genome Res* 25:1043–1055. <http://dx.doi.org/10.1101/gr.186072.114>.
15. Overbeek R, Olson R, Pusch GD, Olsen GJ, Davis JJ, Disz T, Edwards RA, Gerdes S, Parrello B, Shukla M, Vonstein V, Wattam AR, Xia F, Stevens R. 2014. The SEED and the rapid annotation of microbial genomes using subsystems technology (RAST). *Nucleic Acids Res* 42:D206–D214. <http://dx.doi.org/10.1093/nar/gkt1226>.
16. Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang Y, Wishart DS. 2016. PHASTER: a better, faster version of the PHAST phage search tool. *Nucleic Acids Res* 44:W16–W21. <http://dx.doi.org/10.1093/nar/gkw387>.
17. Grissa I, Vergnaud G, Pourcel C. 2007. CRISPRFinder: a Web tool to identify clustered regularly interspaced short palindromic repeats. *Nucleic Acids Res* 35:W52–W57. <http://dx.doi.org/10.1093/nar/gkm360>.
18. Segata N, Börnigen D, Morgan XC, Huttenhower C. 2013. PhyloPhlAn is a new method for improved phylogenetic and taxonomic placement of microbes. *Nat Commun* 4:2304. <http://dx.doi.org/10.1038/ncomms3304>.
19. Cole JR, Wang Q, Cardenas E, Fish J, Chai B, Farris RJ, Kulam-Syed-Mohideen AS, McGarrell DM, Marsh T, Garrity GM, Tiedje JM. 2009. The Ribosomal Database Project: improved alignments and new tools for rRNA analysis. *Nucleic Acids Res* 37:D141–D145. <http://dx.doi.org/10.1093/nar/gkn879>.