

Complete Genome Sequence of *Bacillus pumilus* PDSLzg-1, a Hydrocarbon-Degrading Bacterium Isolated from Oil-Contaminated Soil in China

Kun Hao,^b Hongna Li,^a Feng Li,^a Ping Guo^a

Agricultural Clean Watershed Research Group, Institute of Environment and Sustainable Development in Agriculture, Chinese Academy of Agricultural Sciences, Beijing, People's Republic of China^a; State Key Laboratory for Biology of Plant Diseases and Insect Pests, Institute of Plant Protection, Chinese Academy of Agricultural Sciences, Beijing, People's Republic of China^b

***Bacillus pumilus* strain PDSLzg-1, an efficient hydrocarbon-degrading bacterium, was isolated from oil-contaminated soil. Here, we present the complete sequence of its circular chromosome and circular plasmid. The genomic information is essential for the study of degradation of oil by *B. pumilus* PDSLzg-1.**

Received 11 August 2016 Accepted 17 August 2016 Published 6 October 2016

Citation Hao K, Li H, Li F, Guo P. 2016. Complete genome sequence of *Bacillus pumilus* PDSLzg-1, a hydrocarbon-degrading bacterium isolated from oil-contaminated soil in China. *Genome Announc* 4(5):e01079-16. doi:10.1128/genomeA.01079-16.

Copyright © 2016 Hao et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Ping Guo, guoping@caas.cn.

Bacillus pumilus strain PDSLzg-1 is a novel bacterium isolated from oil-contaminated soil of the Shengli oil field (Shandong, People's Republic of China) in 2015. The bacterium is characterized as Gram-positive, aerobic, motile, rod-shaped, and forming endospores. According to 16S rRNA sequencing analysis, the bacterium is identified as a strain of *Bacillus pumilus*. This strain, designated PDSLzg-1, has the ability to effectively degrade petroleum hydrocarbons, especially C8 to C18 alkanes, and was shown to produce at least one bio-surfactant and may be of interest for an industrial application of oil pollution degradation.

B. pumilus PDSLzg-1 genomic DNA was isolated from overnight culture using a Wizard Genomic DNA purification kit (Promega). Total DNA obtained was subjected to quality control by agarose gel electrophoresis and quantified by Qubit. The genome of *B. pumilus* PDSLzg-1 was sequenced by single molecule real-time (SMRT) technology. Sequencing was performed at the Beijing Novogene Bioinformatics Technology Co., Ltd., producing 95,240 reads with 358× coverage. SMRT Analysis 2.3.0 was used to filter low-quality reads and the filtered reads were assembled to generate one contig without gaps. The genome of *B. pumilus* PDSLzg-1 had a 3,698,973-bp circular chromosome and an 11,801-bp circular plasmid with G+C contents of 41.96% and 39.36%, respectively. Gene prediction was performed on the *B. pumilus* PDSLzg-1 genome by GeneMarkS, resulting in the identification of 3,895 genes with an average length of 848 bp (1). 81 tRNA genes were predicted with tRNAscan-SE, 24 rRNA genes were predicted with rRNAmmer, and seven sRNAs were predicted by a BLAST search against the Rfam database (2–4). A whole-genome BLAST search (*E* value less than 1e−5, minimal alignment length percentage larger than 40%) was performed against KEGG, COG, nr, Swiss-Prot, GO, and TrEMBL (5–9). 3,689 (94.7%) genes were annotated to at least one database.

There were a group of genes in genome of *B. pumilus* strain PDSLzg-1 encoding alcohol dehydrogenase, aldehyde dehydrogenase, and monooxygenase, which are probably related to oil deg-

radation (10, 11). This study presents the first fully sequenced and annotated hydrocarbon-degrading strain of *B. pumilus*, which is valuable for studying its evolution, hydrocarbon-degrading mechanism, and industrial applications.

Accession number(s). The complete annotated genome and plasmid sequences of *B. pumilus* strain PDSLzg-1 were deposited in GenBank under the accession numbers CP016784 and CP016785, respectively.

ACKNOWLEDGMENTS

This work was supported by the China Major Science and Technology Program for Water Pollution Control and Treatment, the Standardization and Scale of Application of Rural Drinking Water Security Technology (2015ZX07402003-4).

REFERENCES

- Besemer J, Lomsadze A, Borodovsky M. 2001. GeneMarkS: a self-training method for prediction of gene starts in microbial genomes. Implications for finding sequence motifs in regulatory regions. *Nucleic Acids Res* 29: 2607–2618. <http://dx.doi.org/10.1093/nar/29.12.2607>.
- Lowe TM, Eddy SR. 1997. tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res* 25: 955–964. <http://dx.doi.org/10.1093/nar/25.5.0955>.
- Lagesen K, Hallin P, Rødland EA, Staerfeldt H-H, Rognes T, Ussery DW. 2007. RNAmmer: consistent and rapid annotation of ribosomal RNA genes. *Nucleic Acids Res* 35:3100–3108. <http://dx.doi.org/10.1093/nar/gkm160>.
- Gardner PP, Daub J, Tate JG, Nawrocki EP, Kolbe DL, Lindgreen S, Wilkinson AC, Finn RD, Griffiths-Jones S, Eddy SR, Bateman A. 2009. Rfam: updates to the RNA families database. *Nucleic Acids Res* 37: D136–D140. <http://dx.doi.org/10.1093/nar/gkn766>.
- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. *J Mol Biol* 215:403–410. [http://dx.doi.org/10.1016/S0022-2836\(05\)80360-2](http://dx.doi.org/10.1016/S0022-2836(05)80360-2).
- Kanehisa M, Goto S, Kawashima S, Okuno Y, Hattori M. 2004. The KEGG resource for deciphering the genome. *Nucleic Acids Res* 32: D277–D280. <http://dx.doi.org/10.1093/nar/gkh063>.
- Tatusov RL, Fedorova ND, Jackson JD, Jacobs AR, Kiryutin B, Koonin EV, Krylov DM, Mazumder R, Mekhedov SL, Nikolskaya AN, Rao BS,

- Smirnov S, Sverdlov AV, Vasudevan S, Wolf YI, Yin JJ, Natale DA. 2003. The COG database: an updated version includes eukaryotes. *BMC Bioinformatics* 4:41. <http://dx.doi.org/10.1186/1471-2105-4-41>.
8. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig JT, Harris MA, Hill DP, Issel-Tarver L, Kasarskis A, Lewis S, Matese JC, Richardson JE, Ringwald M, Rubin GM, Sherlock G. 2000. Gene ontology: tool for the unification of biology. *Nat Genet* 25:25–29. <http://dx.doi.org/10.1038/75556>.
 9. Magrane M, UniProt Consortium. 2011. UniProt Knowledgebase: a hub of integrated protein data. *Database* 2011:bar009. <http://dx.doi.org/10.1093/database/bar009>.
 10. Kotani T, Yamamoto T, Yurimoto H, Sakai Y, Kato N. 2003. Propane monooxygenase and NAD⁺-dependent secondary alcohol dehydrogenase in propane metabolism by *Gordonia* sp. strain TY-5. *J Bacteriol* 185: 7120–7128. <http://dx.doi.org/10.1128/JB.185.24.7120-7128.2003>.
 11. Harayama S, Kishira H, Kasai Y, Shutsubo K. 1999. Petroleum biodegradation in marine environments. *J Mol Microbiol Biotechnol* 1:63–70.