

Complete Genome Sequence of *Magnetospirillum* sp. Strain XM-1, Isolated from the Xi'an City Moat, China

Yinzhao Wang,^{a,b,c} Tongwei Zhang,^{a,b} Wei Lin,^{a,b} Bingfang Zhang,^{a,b} Yao Cai,^{a,b} Caiyun Yang,^{a,b} Jinhua Li,^{a,b} Huangtao Xu,^{a,b} Yongxin Pan^{a,b,c}

PGL, Key Laboratory of Earth and Planetary Physics, Institute of Geology and Geophysics, Chinese Academy of Sciences, Beijing, China^a; France-China Bio-Mineralization and Nano-Structures Laboratory, Institute of Geology and Geophysics, Chinese Academy of Sciences, Beijing, China^b; University of Chinese Academy of Sciences, Beijing, China^c

The magnetotactic bacterium *Magnetospirillum* sp. strain XM-1 was recently isolated from the Xi'an City moat, China. It belongs to the *Rhodospirillaceae* family in the *Alphaproteobacteria* class. Here, we report the complete genome sequence of XM-1. The genome contains a single circular chromosome of 4,825,187 bp and a plasmid of 167,290 bp.

Received 26 August 2016 Accepted 26 August 2016 Published 20 October 2016

Citation Wang Y, Zhang T, Lin W, Zhang B, Cai Y, Yang C, Li J, Xu H, Pan Y. 2016. Complete genome sequence of *Magnetospirillum* sp. strain XM-1, isolated from the Xi'an City moat, China. *Genome Announc* 4(5):e01171-16. doi:10.1128/genomeA.01171-16.

Copyright © 2016 Wang et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Yongxin Pan, yxpan@mail.iggcas.ac.cn.

Magnetotactic bacteria are able to synthesize intracellular nanosized magnetic crystals ($\text{Fe}_3\text{O}_4/\text{Fe}_3\text{S}_4$) usually arranged into a chain structure, known as magnetosomes, that help the cells navigate along the geomagnetic field in aquatic environments (1, 2). *Magnetospirillum* sp. strain XM-1, belonging to the *Alphaproteobacteria* was isolated from the eutrophic city moat in Xi'an, China (3), and synthesizes about 10 magnetite magnetosomes per cell (4).

The complete genome of *Magnetospirillum* sp. strain XM-1 was sequenced using Illumina HiSeq 2000 sequencing platforms (BGI, China). Short-insert (~500-bp) and long-insert (~6,000-bp) size DNA libraries were constructed, and a total of 121,032,268 reads were obtained, corresponding to 1,089 Mb of sequence data. *De novo* and reference-based assemblies were performed by SOAPdenovo version 2.04 (5). Genome of *Magnetospirillum magneticum* AMB-1 (6) was used as a reference strain. For highly complex region, PCRs were conducted to fill all the gaps.

The complete genome sequence of strain XM-1 comprised one circular chromosome of 4,825,187 bp and one plasmid of 167,290 bp. The G+C contents of the chromosome and plasmid are 65.64% and 66.48%, respectively. Gene annotation and analysis were performed using the MaGe Genoscope Web-based system (7). All predicted open reading frames were manually checked. The chromosome contains 4,766 coding sequences (CDSs), 53 tRNAs, two sets of rRNA operons, and 11 miscellaneous RNAs. The average CDS length is 940 bp. Of 4,766 CDSs, 74.26% could be classified as at least one category of the Clusters of Orthologous Groups (COG), in which genes related to signal transduction mechanisms (10.93%) account for dominant COG categories. Inorganic ion transport and metabolism correspond to 5.6% of the gene products in COGs. The plasmid contains 186 CDSs, and their average length is 817 bp. Compared with the chromosome, only 38.17% of plasmid CDSs could be classified as COG groups. Genes associated with replication, recombination,

and repair (8.6%) rank first in COG groups in plasmids while making up only 5.7% in the chromosome.

Magnetosome formation in magnetotactic bacteria is controlled by a cluster of conserved genes (8). In *Magnetospirillum* sp. strain XM-1, we found that these essential genes are organized in *mamGFDC*, *mms6*, *mamAB*, and *mamXY* clusters in an approximately 100-kb region. A total of 39 genes are predicted related to magnetosome synthesis, including *fur*, *feoAB*, and *ftsZ*-like genes, and 24 genes were annotated as transposases or mobile elements (9).

Strain XM-1 possesses a megaplasmid containing three clusters of heavy metal efflux genes, one *czcD* gene, and *merR*-like heavy metal regulation genes. These outer membrane heavy metal efflux genes can help bacteria export the toxic heavy metal (10). Five DNA methylase-related genes, one putative type II restriction enzyme gene, a resolvase domain-containing protein gene, and a putative DNA topoisomerase are present in the plasmid, which may help bacteria avoid foreign DNA invasion (11, 12).

Accession number(s). The complete genome sequence of *Magnetospirillum* sp. strain XM-1 has been deposited at DDBJ/ENA/GenBank under the accession numbers [LN997848](https://www.ncbi.nlm.nih.gov/nuccore/LN997848) and [LN997849](https://www.ncbi.nlm.nih.gov/nuccore/LN997849).

FUNDING INFORMATION

This work, including the efforts of Yongxin Pan, was funded by National Natural Science Foundation of China (NSFC) (41330104). This work, including the efforts of Wei Lin, was funded by the Youth Innovation Promotion Association of the CAS and the State Key Laboratory of Biogeology and Environmental Geology (GBL21402).

REFERENCES

- Bazylinski DA, Lefèvre C, Schüler D. 2013. Magnetotactic bacteria, p 453–494. In Rosenberg E, DeLong E, Lory S, Stackebrandt E, Thompson F (ed), *The prokaryotes*. Springer Verlag, Berlin, Germany Heidelberg.
- Lin W, Bazylinski DA, Xiao T, Wu L-F, Pan Y. 2014. Life with compass: diversity and biogeography of magnetotactic bacteria. *Environ Microbiol* 16:2646–2658. <http://dx.doi.org/10.1111/1462-2920.12313>.

3. Wang Y, Lin W, Li J, Pan Y. 2013. High diversity of magnetotactic *Deltaproteobacteria* in a freshwater niche. *Appl Environ Microbiol* 79: 2813–2817. <http://dx.doi.org/10.1128/AEM.03635-12>.
4. Wang Y, Lin W, Li J, Zhang T, Li Y, Tian J, Gu L, Vander Heyden YV, Pan Y. 2015. Characterizing and optimizing magnetosome production of *Magnetospirillum* sp. XM-1 isolated from Xi'an City Moat, China. *FEMS Microbiol Lett* 362: <http://dx.doi.org/10.1093/femsle/fnv167>.
5. Luo R, Liu B, Xie Y, Li Z, Huang W, Yuan J, He G, Chen Y, Pan Q, Liu Y, Tang J, Wu G, Zhang H, Shi Y, Liu Y, Yu C, Wang B, Lu Y, Han C, Cheung DW, Yiu S-M, Peng S, Zhu X, Liu G, Liao X, Li Y, Yang H, Wang J, Lam T-W, Wang J. 2012. SOAPdenovo2: an empirically improved memory-efficient short-read *de novo* assembler. *GigaScience* 1:18. <http://dx.doi.org/10.1186/2047-217X-1-18>.
6. Matsunaga T, Okamura Y, Fukuda Y, Wahyudi AT, Murase Y, Takeyama H. 2005. Complete genome sequence of the facultative anaerobic magnetotactic bacterium *Magnetospirillum* sp. strain AMB-1. *DNA Res* 12:157–166. <http://dx.doi.org/10.1093/dnares/dsi002>.
7. Vallenet D, Engelen S, Mornico D, Cruveiller S, Fleury L, Lajus A, Rouy Z, Roche D, Salvignol G, Scarpelli C, Médigue C. 2009. Microscope: a platform for microbial genome annotation and comparative genomics. *Database* 2009: <http://dx.doi.org/10.1093/database/bap021>.
8. Komeili A. 2012. Molecular mechanisms of compartmentalization and biomineralization in magnetotactic bacteria. *FEMS Microbiol Rev* 36: 232–255. <http://dx.doi.org/10.1111/j.1574-6976.2011.00315.x>.
9. Kolinko I, Jogler C, Katzmann E, Schüler D. 2011. Frequent mutations within the genomic magnetosome island of *Magnetospirillum gryphiswaldense* are mediated by RecA. *J Bacteriol* 193:5328–5334. <http://dx.doi.org/10.1128/JB.05491-11>.
10. Delmar JA, Su C-C, Yu EW. 2014. Bacterial multidrug efflux transporters. *Annu Rev Biophys* 43:93–117. <http://dx.doi.org/10.1146/annurev-biophys-051013-022855>.
11. Clark TA, Murray IA, Morgan RD, Kislyuk AO, Spittle KE, Boitano M, Fomenkov A, Roberts RJ, Korlach J. 2012. Characterization of DNA methyltransferase specificities using single-molecule, real-time DNA sequencing. *Nucleic Acids Res* 40:e29. <http://dx.doi.org/10.1093/nar/gkr1146>.
12. Downen RH, Pelizzola M, Schmitz RJ, Lister R, Downen JM, Nery JR, Dixon JE, Ecker JR. 2012. Widespread dynamic DNA methylation in response to biotic stress. *Proc Natl Acad Sci U S A* 109:E2183–E2191. <http://dx.doi.org/10.1073/pnas.1209329109>.