

Complete Genome Sequences of Three Important Methicillin-Resistant Clinical Isolates of *Staphylococcus pseudintermedius*

Matthew C. Riley,^a Vincent Perreten,^b David A. Bemis,^a Stephen A. Kania^a

Department of Biomedical and Diagnostic Sciences, Knoxville, Tennessee, USA^a; Institute of Veterinary Bacteriology, Vetsuisse Faculty, University of Bern, Bern, Switzerland^b

We report the first complete genome sequences of three predominant clones (ST68, ST71, and ST84) of methicillin-resistant *Staphylococcus pseudintermedius* in North America. All strains were isolated from canine infections and have different SCCmec elements and antibiotic resistance gene patterns.

Received 31 August 2016 Accepted 2 September 2016 Published 20 October 2016

Citation Riley MC, Perreten V, Bemis DA, Kania SA. 2016. Complete genome sequences of three important methicillin-resistant clinical isolates of *Staphylococcus pseudintermedius*. *Genome Announc* 4(5):e01194-16. doi:10.1128/genomeA.01194-16.

Copyright © 2016 Riley et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Stephen A. Kania, skania@utk.edu.

Staphylococcus pseudintermedius is a Gram-positive opportunistic pathogen (1) that primarily causes infections in canines but is also relevant to human medicine (2–4), particularly with the worldwide expansion of methicillin-resistant clonal lineages (5–8). While a complete genome is publically available for a European methicillin-resistant *S. pseudintermedius* (MRSP) isolate (9), no complete MRSP genomes from dominant clonal lineages in North America are available. Here, we present the complete circular chromosomes of MRSP strains NA45, 081661, and 063228, which were isolated from canine infections in 2006 and 2008 and represent three dominant sequence types (ST) in North America, namely, ST84, ST71, and ST68, respectively (10).

All isolates were sequenced using Roche/454 (Roche Diagnostics, Switzerland), Illumina MiSeq (Illumina, Inc., USA), and Ion Torrent technologies (Thermo, Fisher Scientific, USA); isolates NA45 and 063228 were additionally sequenced using PacBio single-molecule real-time sequencing technology (Pacific Biosciences, USA). All genomes were mapped using the Argus Whole Genome Mapping System (OpGen, Inc., USA). *De novo* assemblies were individually produced and merged using Geneious version 9.1.3 (11) and CLC Genomics Workbench version 9.0 (<https://www.qiagenbioinformatics.com>). PacBio reads and whole-genome maps were used for scaffolding and genome closure (12). Automated annotation for strain 063228 was performed using the RAST server (<http://rast.nmpdr.org>), while the NCBI Prokaryote Genome Annotation Pipeline (http://www.ncbi.nlm.nih.gov/genome/annotation_prok) was used for NA45 and 081661.

The total number of high-quality reads for strain NA45 were 29,463 (PacBio), 583,182 (Illumina), 279,674 (Roche), and 4,660,374 (Ion Torrent), resulting in >250-fold overall coverage. The NA45 genome is 2,841,212 bp with a 37.3% GC content, 2,665 predicted coding sequences, and 78 predicted RNAs. High-quality reads for strain 081661 were 456,358 (Illumina), 129,593 (Roche), and 2,578,704 (Ion Torrent), resulting in >250-fold overall coverage. The 081661 genome is 2,731,109 bp with a 37.5% GC content, 2,610 predicted coding sequences, and 87 predicted RNAs.

High-quality reads for strain 063228 were 24,585 (PacBio), 15,886,636 (Illumina), 113,288 (Roche), and 3,864,512 (Ion Torrent), resulting in >250-fold overall coverage. The 063228 genome is 2,766,566 bp with a 37.4% GC content, 2,734 predicted coding sequences, and 77 predicted RNAs.

The 081661 genome shared 99% identity over 96% of the published ST71 European isolate E140, with the major differences resulting from prophage composition (9). In addition to the methicillin resistance gene *mecA*, all strains contained the beta-lactamase gene *blaZ*, the kanamycin and streptomycin resistance genes *aph(3')-III* and *ant(6)-Ia*. Strains 063228 and 081661 also harbor genes conferring resistance to gentamicin-kanamycin [*aac(6')-Ie-aph(2')-Ia*], macrolides-lincosamides-streptogramins B [*erm(B)*], while strain 063228 had additional lincosamide and tetracycline resistance genes *lnu(A)* and *tet(M)* (13). The methicillin resistance gene *mecA* was found on the staphylococcal cassette chromosome *mec* (SCCmec) SCCmecV in 063228, on SCCmecII-III in strain 081661, and on a novel SCCmec element in strain NA45. This 43,922-bp cassette has *mecA* integrated in the opposite direction compared to all other SCCmec elements (14) and contains the recombinase gene *ccrC6*.

The complete genomes of three strains belonging to three predominant clones causing infections in dogs in the United States permits further comparative genomic analyses and gives new insights into the molecular epidemiology and biological characteristics of *S. pseudintermedius*.

Accession number(s). These whole-genome projects have been deposited in DDBJ/ENA/GenBank under the accession numbers CP016072, CP016073, and CP015626. The versions described in this paper are the first versions, CP016072.1, CP016073.1, and CP015626.1.

ACKNOWLEDGMENTS

We acknowledge the University of Tennessee, College of Veterinary Medicine Bacteriology Lab for isolating strains 081661 and 063228 and for technical assistance and susceptibility testing, as well as Faye Hartmann at the Wisconsin School of Veterinary Medicine for providing strain NA45.

We also thank the Canine Health Foundation and the Center of Excellence in Livestock Diseases and Human Health for funding and support.

FUNDING INFORMATION

This work, including the efforts of Stephen Anthony Kania, was funded by the Canine Health Foundation. This work, including the efforts of Stephen Anthony Kania, was funded by the Center of Excellence in Livestock Diseases and Human Health.

The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

REFERENCES

- Devriese LA, Vancanneyt M, Baele M, Vaneechoutte M, De Graef E, Snauwaert C, Cleenwerck I, Dawyndt P, Swings J, Decostere A, Haesebrouck F. 2005. *Staphylococcus pseudintermedius* sp. nov., a coagulase-positive species from animals. *Int J Syst Evol Microbiol* 55:1569–1573. <http://dx.doi.org/10.1099/ijs.0.63413-0>.
- Van Duijkeren E, Cattri B, Greko C, Moreno MA, Pomba MC, Pyörälä S, Ruzauskas M, Sanders P, Threlfall EJ, Torren-Edo J, Törneke K, Scientific Advisory Group on Antimicrobials (SAGAM). 2011. Review on methicillin-resistant *Staphylococcus pseudintermedius*. *J Antimicrob Chemother* 66:2705–2714. <http://dx.doi.org/10.1093/jac/dkr367>.
- Somayaji R, Priyantha MA, Rubin JE, Church D. 2016. Human infections due to *Staphylococcus pseudintermedius*, an emerging zoonosis of canine origin: report of 24 cases. *Diagn Microbiol Infect Dis* 85:471–476. <http://dx.doi.org/10.1016/j.diagmicrobio.2016.05.008>.
- Kuan EC, Yoon AJ, Vijayan T, Humphries RM, Suh JD. 2016. Canine *Staphylococcus pseudintermedius* sinonasal infection in human hosts. *Int Forum Allergy Rhinol* 6:710–715. <http://dx.doi.org/10.1002/alr.21732>.
- Jones RD, Kania SA, Rohrbach BW, Frank LA, Bemis DA. 2007. Prevalence of oxacillin- and multidrug-resistant staphylococci in clinical samples from dogs: 1,772 samples (2001–2005). *J Am Vet Med Assoc* 230:221–227. <http://dx.doi.org/10.2460/javma.230.2.221>.
- Bemis DA, Jones RD, Frank LA, Kania SA. 2009. Evaluation of susceptibility test breakpoints used to predict *mecA*-mediated resistance in *Staphylococcus pseudintermedius* isolated from dogs. *J Vet Diagn Invest* 21:53–58. <http://dx.doi.org/10.1177/104063870902100108>.
- Perreten V, Kadlec K, Schwarz S, Grönlund Andersson U, Finn M, Greko C, Moodley A, Kania SA, Frank LA, Bemis DA, Franco A, Iurescia M, Battisti A, Duim B, Wagenaar JA, van Duijkeren E, Weese JS, Fitzgerald JR, Rossano A, Guardabassi L, Jr. 2010. Clonal spread of methicillin-resistant *Staphylococcus pseudintermedius* in Europe and North America: an international multicentre study. *J Antimicrob Chemother* 65:1145–1154. <http://dx.doi.org/10.1093/jac/dkq078>.
- Börjesson S, Gómez-Sanz E, Ekström K, Torres C, Grönlund U. 2015. *Staphylococcus pseudintermedius* can be misdiagnosed as *Staphylococcus aureus* in humans with dog bite wounds. *Eur J Clin Microbiol Infect Dis* 34:839–844. <http://dx.doi.org/10.1007/s10096-014-2300-y>.
- Moodley A, Riley MC, Kania SA, Guardabassi L. 2013. Genome sequence of *Staphylococcus pseudintermedius* strain E140, an ST71 European-associated methicillin-resistant isolate. *Genome Announc* 1(2):e0020712. <http://dx.doi.org/10.1128/genomeA.00207-12>.
- Solyman SM, Black CC, Duim B, Perreten V, van Duijkeren E, Wagenaar JA, Eberlein LC, Sadeghi LN, Videla R, Bemis DA, Kania SA. 2013. Multilocus sequence typing for characterization of *Staphylococcus pseudintermedius*. *J Clin Microbiol* 51:306–310. <http://dx.doi.org/10.1128/JCM.02421-12>.
- Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Duran C, Thierer T, Ashton B, Meintjes P, Drummond A. 2012. Geneious basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 28:1647–1649. <http://dx.doi.org/10.1093/bioinformatics/bts199>.
- Onmus-Leone F, Hang J, Clifford RJ, Yang Y, Riley MC, Kuschner RA, Waterman PE, Lesho EP. 2012. Enhanced de novo assembly of high throughput pyrosequencing data using whole genome mapping. *PLoS One* 8:e61762. <http://dx.doi.org/10.1371/journal.pone.0061762>.
- Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, Aarestrup FM, Larsen MV. 2012. Identification of acquired antimicrobial resistance genes. *J Antimicrob Chemother* 67:2640–2644. <http://dx.doi.org/10.1093/jac/dks261>.
- International Working Group on the Classification of Staphylococcal Cassette Chromosome Elements (IWG-SCC). 2009. Classification of staphylococcal cassette chromosome *mec* (SCC*mec*): guidelines for reporting novel SCC*mec* elements. *Antimicrob Agents Chemother* 53:4961–4967. <http://dx.doi.org/10.1128/AAC.00579-09>.