




Draft Genome Sequence of *Streptomyces* sp. Strain IB2014011-1, Isolated from *Trichoptera* sp. Larvae of Lake Baikal

 Denis V. Axenov-Gribanov,^{a,b} Bogdan T. Tokovenko,^c Yuriy V. Rebets,^c Irina V. Voytsekhovskaya,^{a,b} Zhanna M. Shatilina,^{a,b} Eugenii S. Protasov,^a Andriy N. Luzhetskyy,^{c,d} Maxim A. Timofeyev^a

Irkutsk State University, Institute of Biology, Irkutsk, Russia^a; Baikal Research Centre, Irkutsk, Russia^b; Helmholtz Center for Infection Research (HZI), Helmholtz Institute for Pharmaceutical Research Saarland (HIPS), Saarbrücken, Germany^c; Universität des Saarlandes, Saarbrücken, Germany^d

ABSTRACT Unique ecosystems with specific environmental conditions have been proven to be a promising source for isolation of new actinobacterial strains. Ancient Lake Baikal is one of the greatest examples of an ecosystem with high species biodiversity and endemism caused by long-lasting isolated evolution and stable environmental conditions. Herein we report the draft genome sequence of *Streptomyces* sp. strain IB2014011-1, which was isolated from insect *Trichoptera* sp. larvae collected at the bottom of Lake Baikal.

Actinobacteria are high-GC Gram-positive bacteria with high ability to produce secondary metabolites (1). It has been previously shown that actinobacteria isolated from unusual ecosystems often produce new biologically active compounds (2). Recently, we reported the isolation of new actinobacterial strains from areas of endemism such as Lake Baikal (3) and caves (4).

Genomic DNA was extracted from *Streptomyces* sp. strain IB2014011-1, which was isolated from insect *Trichoptera* sp. larvae (3). Standard protocol was used to prepare a paired-end library. The library had reads 100 bp long, insert size 259.07 ($\sigma = 85.71$), and mean coverage 720 \times , as determined by postassembly mapping with bwa v 0.7.13-r1126 (mem alignment algorithm) (5). The raw sequencing data were obtained using Illumina HiSeq 2500 technology. The genome was assembled using SPAdes v 3.7 (6). A total of 73 contigs and 68 scaffolds were assembled. Scaffolding, performed using SSPACE 2.1 Premium (7), resulted in 43 scaffolds, of which 31 passed coverage and length (at least 1 kbp) thresholds. Genome annotation was performed using Prokka (8) and antiSMASH v.3 (9), followed by manual presubmission curation.

The genome of *Streptomyces* sp. IB2014011-1 is large: all scaffolds together are 8,195,763 bp long. The G+C content (71.5%) and the number of protein-coding (7,323) and tRNA (78) genes are in accordance with those of other *Streptomyces* strains. The average gene length is 988 bp and the average coding density is 88.3%. The genome of *Streptomyces* sp. IB2014 011-1 contains at least four rRNA gene clusters, judging by the coverage of the assembled rRNA gene cluster. In addition, one *ssrA* transfer-messenger RNA (*tmRNA*) and 206 TTA codons within predicted coding sequences (CDS) were found.

Functional annotation of *Streptomyces* sp. IB2014011-1 genes within the bactNOG subset of the eggNOG v 4 database (performed using protein BLAST with an expectation value cutoff 0.001) (10) showed that 4,610 (63%) out of 7,323 protein-coding genes had at least some biological function assigned, with some of the genes assigned to more than one category. Of the remainder, 194 CDS (12.6%) had no hits against bactNOGs, and 1,857 CDS (25.4%) had hits but were not assigned to functional

Received 24 January 2017 Accepted 2 March 2017 Published 27 April 2017

Citation Axenov-Gribanov DV, Tokovenko BT, Rebets YV, Voytsekhovskaya IV, Shatilina ZM, Protasov ES, Luzhetskyy AN, Timofeyev MA. 2017. Draft genome sequence of *Streptomyces* sp. strain IB2014011-1, isolated from *Trichoptera* sp. larvae of Lake Baikal. *Genome Announc* 5:e00062-17. <https://doi.org/10.1128/genomeA.00062-17>.

Copyright © 2017 Axenov-Gribanov et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Denis V. Axenov-Gribanov, denis.axengri@gmail.com.

D.V.A.-G. and B.T.T. contributed equally to this work.

categories. Among the genes with functional assignment, 1,979 (27%) are implicated in metabolism, including 142 (1.9%) putatively involved in secondary metabolism.

The potential of this strain to produce secondary metabolites was analyzed with the search tool antiSMASH v.3 (9). Thirty putative secondary metabolism gene clusters were found with this method, and 55 more were predicted with the ClusterFinder algorithm.

A gene cluster of 50 is predicted to be involved in the assembly of the compound grixazone B (11) (accurate mass is 417.0616 m/z), which was found in the culture of the strain. All 13 *gri* genes of the originally reported gene cluster are present, with an average nucleotide identity of 88% across the 15-kbp fragment.

Thus, the genome information provided by the draft sequence of *Streptomyces* sp. IB2014011-1 has importance for basic as well as applied microbial genomic research.

Accession number(s). The genome sequence of *Streptomyces* sp. IB2014011-1 has been deposited at DDBJ/ENA/GenBank under the accession number [LZQS00000000](https://www.ncbi.nlm.nih.gov/nuclseq/LZQS00000000). The version described in this paper is LZQS00000000.1.

ACKNOWLEDGMENTS

This study was supported by the Ministry of Education and Science of the Russian Federation as a part of the Goszadanie project (6.1387.2017, 6.9654.2017), the Russian Science Foundation (project N 15-14-10008), the Russian Foundation for Basic Research (projects N 16-34-00686, 16-34-60060), and grants from the Irkutsk State University for Young Researchers and Deutscher Akademischer Austauschdienst.

REFERENCES

1. Das S, Ward LR, Burke C. 2008. Prospects of using marine actinobacteria as probiotics in aquaculture. *Appl Microbiol Biotechnol* 81:419–429. <https://doi.org/10.1007/s00253-008-1731-8>.
2. Jenke-Kodama H, Dittmann E. 2009. Evolution of metabolic diversity: insights from microbial polyketide synthases. *Phytochemistry* 70:1858–1866. <https://doi.org/10.1016/j.phytochem.2009.05.021>.
3. Axenov-Gribanov D, Rebets Y, Tokovenko B, Voytsekhovskaya I, Timofeyev M, Luzhetsky A. 2016. The isolation and characterization of actinobacteria from dominant benthic macroinvertebrates endemic to Lake Baikal. *Folia Microbiol (Praga)* 61:159–168. <https://doi.org/10.1007/s12223-015-0421-z>.
4. Axenov-Gribanov DV, Voytsekhovskaya IV, Tokovenko BT, Protasov ES, Gamaiunov SV, Rebets YV, Luzhetsky AN, Timofeyev MA. 2016. Correction: actinobacteria isolated from an underground lake and moonmilk speleothem from the biggest conglomeratic karstic cave in Siberia as sources of novel biologically active compounds. *PLoS One* 11:e0152957. <https://doi.org/10.1371/journal.pone.0152957>.
5. Li H. 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. *arXiv arXiv:1303.3997 [q-bio.GN]*. 1303.3997v2.
6. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
7. Boetzer M, Henkel CV, Jansen HJ, Butler D, Pirovano W. 2011. Scaffolding pre-assembled contigs using SSPACE. *Bioinformatics* 27:578–579. <https://doi.org/10.1093/bioinformatics/btq683>.
8. Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068–2069. <https://doi.org/10.1093/bioinformatics/btu153>.
9. Weber T, Blin K, Duddela S, Krug D, Kim HU, Brucoleri R, Lee SY, Fischbach MA, Müller R, Wohlleben W, Breitling R, Takano E, Medema MH. 2015. AntiSMASH 3.0—a comprehensive resource for the genome mining of biosynthetic gene clusters. *Nucleic Acids Res* 43:W237–W243. <https://doi.org/10.1093/nar/gkv437>.
10. Powell S, Forslund K, Szklarczyk D, Trachana K, Roth A, Huerta-Cepas J, Gabaldón T, Rattei T, Creevey C, Kuhn M, Jensen LJ, von Mering C, Bork P. 2014. eggNOG v4.0: nested orthology inference across 3686 organisms. *Nucleic Acids Res* 42:D231–D239. <https://doi.org/10.1093/nar/gkt1253>.
11. Ohnishi Y, Furusho Y, Higashi T, Chun HK, Furihata K, Sakuda S, Horinouchi S. 2004. Structures of grixazone A and B, A-factor-dependent yellow pigments produced under phosphate depletion by *Streptomyces griseus*. *J Antibiot (Tokyo)* 57:218–223. <https://doi.org/10.7164/antibiotics.57.218>.