



Sequence of Reston Virus Isolate AZ-1435, an Ebolavirus Isolate Obtained during the 1989–1990 Reston Virus Epizootic in the United States

Joseph P. Cornish,^a Larissa Diaz,^a Stacy M. Ricklefs,^b Kishore Kanakabandi,^b Jennifer Sword,^c  Peter B. Jahrling,^{a,c}  Jens H. Kuhn,^c Stephen F. Porcella,^b  Reed F. Johnson^a

Emerging Viral Pathogens Section, Division of Intramural Research, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Fort Detrick, Frederick, Maryland, USA^a; Genomics Unit, Research Technologies Section, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Rocky Mountain Laboratories, Hamilton, Montana, USA^b; Integrated Research Facility at Fort Detrick, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Fort Detrick, Frederick, Maryland, USA^c

ABSTRACT Reston virus (RESTV) was discovered in 1989–1990 during three connected epizootics of highly lethal viral hemorrhagic fever among captive macaques in primate housing facilities in the United States and Philippines. Currently, only one RESTV isolate from that outbreak (named Pennsylvania) has been sequenced. Here, we report the sequence of a second isolate, Reston virus/M.fascicularis-tc/USA/1990/Philippines89-AZ1435.

Reston virus (RESTV; *Filoviridae*: *Ebolavirus*: *Reston ebolavirus* [1]) was discovered in 1989–1990, during a highly lethal viral hemorrhagic fever epizootic among crab-eating macaques (*Macaca fascicularis*) in primate-holding facilities in Reston, Virginia (1989 and 1990), Philadelphia, Pennsylvania (1989), and Alice, Texas (1989), USA. The affected animals were traced to a single primate export facility in the Philippines (2–5). RESTV reemerged in crab-eating macaques imported from the same facility in 1992 in Siena, Italy (6, 7), and in 1996 in Alice, Texas, USA (8–11), and was found in sick domestic pigs (*Sus scrofa*) and unspecified nonhuman primates in the Philippines in 2008 (12) and 2015 (13), respectively.

RESTV isolate sequence data are sparse (14), and only a single complete (15) genome (Reston virus/M.fascicularis-tc/USA/1989/Philippines89-Pennsylvania; RefSeq no. NC_004161 [16]) is available from the 1989–1990 epizootic. We sequenced RESTV AZ-1435, an isolate obtained from the spleen of a crab-eating macaque with signs of respiratory infection sampled during the 1990 Reston, Virginia, USA, episode of the 1989–1990 outbreak (17, 18). RESTV AZ-1435 was isolated by inoculating MA-104 cells with 10% spleen homogenate, followed by passage of MA-104 cell supernatant onto Vero E6 cells (17). We further passaged this sample by infecting Vero E6 cells and harvesting supernatant.

The virus was inactivated with Trizol LS (Thermo Fisher Scientific), and RNA was extracted as described previously (19). rRNA removal (low-input Ribo-Zero, Illumina) was performed prior to cDNA synthesis. cDNA libraries for sequencing were prepared using the Ovation RNA-Seq version 2 system (NuGEN Technologies) and the Ultralow DR (directional read) multiplex system (NuGEN Technologies) following the manufacturers' recommendations. After single-primer isothermal amplification (SPIA, NuGEN Technologies), cDNA was purified using Agencourt RNAClean XP beads (Beckman Coulter, Inc. Life Sciences), sheared, and sequenced via paired-end (2 × 100 bp) in rapid chemistry mode on a HiSeq 2500 sequencing system (Illumina). SMARTer rapid ampli-

Received 31 October 2016 **Accepted** 8 November 2016 **Published** 12 January 2017

Citation Cornish JP, Diaz L, Ricklefs SM, Kanakabandi K, Sword J, Jahrling PB, Kuhn JH, Porcella SF, Johnson RF. 2017. Sequence of Reston virus isolate AZ-1435, an ebolavirus isolate obtained during the 1989–1990 Reston virus epizootic in the United States. *Genome Announc* 5:e01448-16. <https://doi.org/10.1128/genomeA.01448-16>.

Copyright © 2017 Cornish et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Reed F. Johnson, johnsonreed@niaid.nih.gov.

fication of cDNA ends (RACE) 5'/3' kit (Clontech Laboratories, Inc.) was used to amplify the 5' end of RNA. The 3' end was amplified by adding adapter oligonucleotides with T4 RNA ligase I (New England BioLabs) and reverse transcriptase (RT)-PCR. RT-PCR products were cloned into pGEM T-easy vectors (Promega) for sequencing.

Trimmomatic (20), was used to trim reads for quality and length prior to assembly. Read quality was assessed pre- and post-trimming using open source FastQC version 0.11.5 software (Babraham Bioinformatics, <http://www.bioinformatics.babraham.ac.uk/projects/fastqc>). Reads were mapped to the RESTV reference genome (RefSeq no. NC_004161) using the Burrows–Wheeler aligner MEM (BWA-MEM) algorithm (21). Variants were identified using genome analysis toolkit (GATK) best practices (22, 23) and were manually inspected with Integrative Genomics Viewer (IGV) (24, 25). A final sequence was identified by comparison to NC_004161.

The RESTV AZ-1435 genome differs from the Pennsylvania genome at few locations, supporting previous investigations indicating that the 1989–1990 U.S. episodes were directly connected. Our data support the hypothesis that both RESTV isolates share an ancestor virus that was exported to all affected U.S. locations.

Accession number(s). The GenBank accession number of RESTV AZ-1435, now designated Reston virus/M.fascicularis-tc/USA/1989/Philippines89-AZ1435 based on current filovirus nomenclature standards (26), is [KY008770](https://www.ncbi.nlm.nih.gov/nuccore/KY008770).

ACKNOWLEDGMENTS

We are grateful to Laura Bollinger (IRF-Frederick) for editing the manuscript.

This work was supported by the Division of Intramural Research and the Division of Clinical Research, both of the National Institute of Allergy and Infectious Diseases (NIAID), National Institutes of Health, United States. The content of this publication does not necessarily reflect the views or policies of the U.S. Department of Health and Human Services or of the institutions and companies affiliated with the authors. This work was supported in part through Battelle Memorial Institute's prime contract with NIAID under contract no. HHSN2722007000161. A subcontractor to Battelle Memorial Institute who performed this work is J.H.K., an employee of Tunnell Government Services, Inc.

REFERENCES

- Kuhn JH, Becker S, Ebihara H, Geisbert TW, Jahrling PB, Kawaoka Y, Netesov SV, Nichol ST, Peters CJ, Volchkov VE, Ksiazek TG. 2011. Family *Filoviridae*, p 665–671. In King AMQ, Adams MJ, Carstens EB, Lefkowitz EJ (ed.), *Virus taxonomy—ninth report of the International Committee on taxonomy of viruses*. Elsevier/Academic Press, London, United Kingdom.
- Geisbert TW, Jahrling PB. 1990. Use of immunoelectron microscopy to show Ebola virus during the 1989 United States epizootic. *J Clin Pathol* 43:813–816. <https://doi.org/10.1136/jcp.43.10.813>.
- Jahrling PB, Geisbert TW, Dalgard DW, Johnson ED, Ksiazek TG, Hall WC, Peters CJ. 1990. Preliminary report: isolation of Ebola virus from monkeys imported to USA. *Lancet* 335:502–505. [https://doi.org/10.1016/0140-6736\(90\)90737-P](https://doi.org/10.1016/0140-6736(90)90737-P).
- Hayes CG, Burans JP, Ksiazek TG, Del Rosario RA, Miranda ME, Manaloto CR, Barrientos AB, Robles CG, Dayrit MM, Peters CJ. 1992. Outbreak of fatal illness among captive macaques in Philippines caused by an Ebola-related filovirus. *Am J Trop Med Hyg* 46:664–671.
- Miranda ME, Miranda NL. 2011. Reston ebolavirus in humans and animals in Philippines: a review. *J Infect Dis* 204(suppl 3):S757–S760. <https://doi.org/10.1093/infdis/jir296>.
- Ciorba A, Matteucci G, Perini L, Caristo ME, Brown D. 1997. Infezione da virus Ebola nella scimmia: reperti clinici et anatomoistopatologici. Osservati nel corso del primo episodio verificatosi in Europa. *Veterinaria (Cremona)* 11:109–112.
- Perini L. 2000. Outbreak of Ebola virus in Italy. *Folia Primatol* 71:280.
- Centers for Disease Control and Prevention. 1996. Ebola-Reston virus infection among quarantined nonhuman primates—Texas, 1996. *MMWR Morb Mortal Wkly Rep* 45:314–316.
- Miranda ME, Ksiazek TG, Retuya TJ, Khan AS, Sanchez A, Fulhorst CF, Rollin PE, Calaor AB, Manalo DL, Roces MC, Dayrit MM, Peters CJ. 1999. Epidemiology of Ebola (subtype Reston) virus in Philippines, 1996. *J Infect Dis* 179(suppl 1):S115–S119. <https://doi.org/10.1086/514314>.
- Miranda ME, Yoshikawa Y, Manalo DL, Calaor AB, Miranda NL, Cho F, Ikegami T, Ksiazek TG. 2002. Chronological and spatial analysis of the 1996 Ebola Reston virus outbreak in a monkey breeding facility in Philippines. *Exp Anim* 51:173–179. <https://doi.org/10.1538/expanim.51.173>.
- Rollin PE, Williams RJ, Bressler DS, Pearson S, Cottingham M, Pucak G, Sanchez A, Trappier SG, Peters RL, Greer PW, Zaki S, Demarcus T, Hendricks K, Kelley M, Simpson D, Geisbert TW, Jahrling PB, Peters CJ, Ksiazek TG. 1999. Ebola (subtype Reston) virus among quarantined nonhuman primates recently imported from the Philippines to the United States. *J Infect Dis* 179(suppl 1):S108–S114. <https://doi.org/10.1086/514303>.
- Barrette RW, Metwally SA, Rowland JM, Xu L, Zaki SR, Nichol ST, Rollin PE, Towner JS, Shieh WJ, Batten B, Sealy TK, Carrillo C, Moran KE, Bracht AJ, Mayr GA, Sirios-Cruz M, Catbagan DP, Lautner EA, Ksiazek TG, White WR, McIntosh MT. 2009. Discovery of swine as a host for the Reston ebolavirus. *Science* 325:204–206. <https://doi.org/10.1126/science.1172705>.
- Geronimo JY. 2015. Reston virus—Philippines (02): monkeys. ProMED-mail, archive no. 20150907.3629141. <http://www.promedmail.org>.
- Burk R, Bollinger L, Johnson JC, Wada J, Radoshitzky SR, Palacios G, Bavari S, Jahrling PB, Kuhn JH. 2016. Neglected filoviruses. *FEMS Microbiol Rev* 40:494–519. <https://doi.org/10.1093/femsre/fuw010>.
- Ladner JT, Kuhn JH, Palacios G. 2016. Standard finishing categories for high-throughput sequencing of viral genomes. *Rev Sci Tech* 35:43–52. <https://doi.org/10.20506/rst.35.1.2416>.
- Kuhn JH, Andersen KG, Bao Y, Bavari S, Becker S, Bennett RS, Bergman NH, Blinkova O, Bradfute S, Brister JR, Bukreyev A, Chandran K, Chep-

- urnov AA, Davey RA, Dietzgen RG, Doggett NA, Dolnik O, Dye JM, Enterlein S, Fenimore PW, Formenty P, Freiberg AN, Garry RF, Garza NL, Gire SK, Gonzalez JP, Griffiths A, Happi CT, Hensley LE, Herbert AS, Hevey MC, Hoenen T, Honko AN, Ignatyev GM, Jahrling PB, Johnson JC, Johnson KM, Kindrachuk J, Klenk HD, Kobinger G, Kochel TJ, Lackemeyer MG, Lackner DF, Leroy EM, Lever MS, Mühlberger E, Netesov SV, Olinger GG, Omilabu SA, Palacios G, Panchal RG, Park DJ, Patterson JL, Paweska JT, Peters CJ, Pettitt J, Pitt L, Radoshitzky SR, Ryabchikova EI, Saphire EO, Sabeti PC, Sealfon R, Shestopalov AM, Smither SJ, Sullivan NJ, Swanepoel R, Takada A, Towner JS, van der Groen G, Volchkov VE, Volchkova VA, Wahl-Jensen V, Warren TK, Warfield KL, Weidmann M, Nichol ST. 2014. Filovirus RefSeq entries: evaluation and selection of filovirus type variants, type sequences, and names. *Viruses* 6:3663–3682. <https://doi.org/10.3390/v6093663>.
17. Jahrling PB, Geisbert TW, Jaax NK, Hanes MA, Ksiazek TG, Peters CJ. 1996. Experimental infection of cynomolgus macaques with Ebola-Reston filoviruses from the 1989–1990 U.S. epizootic, p 115–134. *In* Schwarz TF, Siegl G (ed.), *Imported virus infections*, vol. 11. Springer, Vienna, Austria.
 18. Dalgard DW, Hardy RJ, Pearson SL, Pucak GJ, Quander RV, Zack PM, Peters CJ, Jahrling PB. 1992. Combined simian hemorrhagic fever and Ebola virus infection in cynomolgus monkeys. *Lab Anim Sci* 42:152–157.
 19. Matsuno K, Weisend C, Travassos da Rosa APA, Anzick SL, Dahlstrom E, Porcella SF, Dorward DW, Yu XJ, Tesh RB, Ebihara H. 2013. Characterization of the Bhanja serogroup viruses (*Bunyaviridae*): a novel species of the genus *phlebovirus* and its relationship with other emerging tick-borne phleboviruses. *J Virol* 87:3719–3728. <https://doi.org/10.1128/JVI.02845-12>.
 20. Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
 21. Li H, Durbin R. 2010. Fast and accurate long-read alignment with Burrows–Wheeler transform. *Bioinformatics* 26:589–595. <https://doi.org/10.1093/bioinformatics/btp698>.
 22. DePristo MA, Banks E, Poplin R, Garimella KV, Maguire JR, Hartl C, Philippakis AA, del Angel G, Rivas MA, Hanna M, McKenna A, Fennell TJ, Kernytsky AM, Sivachenko AY, Cibulskis K, Gabriel SB, Altshuler D, Daly MJ. 2011. A framework for variation discovery and genotyping using next-generation DNA sequencing data. *Nat Genet* 43:491–498. <https://doi.org/10.1038/ng.806>.
 23. McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, Garimella K, Altshuler D, Gabriel S, Daly M, DePristo MA. 2010. The genome analysis toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res* 20:1297–1303. <https://doi.org/10.1101/gr.107524.110>.
 24. Robinson JT, Thorvaldsdóttir H, Winckler W, Guttman M, Lander ES, Getz G, Mesirov JP. 2011. Integrative genomics viewer. *Nat Biotechnol* 29:24–26. <https://doi.org/10.1038/nbt.1754>.
 25. Thorvaldsdóttir H, Robinson JT, Mesirov JP. 2013. Integrative genomics viewer (IGV): high-performance genomics data visualization and exploration. *Brief Bioinform* 14:178–192. <https://doi.org/10.1093/bib/bbs017>.
 26. Kuhn JH, Bao Y, Bavari S, Becker S, Bradfute S, Brister JR, Bukreyev AA, Chandran K, Davey RA, Dolnik O, Dye JM, Enterlein S, Hensley LE, Honko AN, Jahrling PB, Johnson KM, Kobinger G, Leroy EM, Lever MS, Mühlberger E, Netesov SV, Olinger GG, Palacios G, Patterson JL, Paweska JT, Pitt L, Radoshitzky SR, Saphire EO, Smither SJ, Swanepoel R, Towner JS, van der Groen G, Volchkov VE, Wahl-Jensen V, Warren TK, Weidmann M, Nichol ST. 2013. Virus nomenclature below the species level: a standardized nomenclature for natural variants of viruses assigned to the family *Filoviridae*. *Arch Virol* 158:301–311. <https://doi.org/10.1007/s00705-012-1454-0>.