



Draft Genome Sequence of the Luminescent Strain *Vibrio campbellii* LB102, Isolated from a Black Tiger Shrimp (*Penaeus monodon*) Broodstock Rearing System

Sujeet Kumar,^a Ashok Kumar Jangam,^a V. Akhil,^a Vidya Rajendran,^a Vinaya Kumar Katneni,^a J. Joseph Sahaya Rajan,^a Monendra Grover,^b Viswas Konasagara Nagaleekar,^c S. V. Alavandi,^a K. K. Vijayan^a

ICAR-Central Institute of Brackishwater Aquaculture, Chennai, India^a; Centre for Agricultural Bioinformatics, ICAR-Indian Agricultural Statistics Research Institute, New Delhi, India^b; ICAR-Indian Veterinary Research Institute, Izatnagar, Bareilly, India^c

ABSTRACT We report here the genome sequence of *Vibrio campbellii* LB102, isolated from the broodstock rearing system of a shrimp hatchery in India. Sequence analysis revealed the presence of effector toxins of the type III (YopT, sharing 39% identity with *Yersinia pestis*) and type VI (VgrG-3 and hemolysin coregulated protein of *V. cholerae*) secretion systems.

Vibrio campbellii is an emerging bacterial pathogen that affects the mysis and early postlarval stages of penaeid shrimp. It closely resembles the marine pathogen *V. harveyi* and is almost indistinguishable from it based on phenotypic characteristics, 16S rRNA gene sequence analysis, and DNA-DNA homology (1). In the past, *V. campbellii* was often misidentified and classified as *V. harveyi* (2). Development of multilocus sequence typing using housekeeping genes and whole-genome sequencing, however, have paved the way for their differentiation (2). While investigating luminescent bacterial disease, *V. campbellii* LB102 was isolated in 2006 from a broodstock rearing system at a tiger shrimp hatchery on the southeast coast of India (3). The strain was luminescent, was confirmed as *V. campbellii* using *rpoD* gene sequence analysis, and was found to be virulent in an experimental challenge trial on postlarvae.

The genomic DNA from LB102 was extracted using a Qiagen genomic DNA extraction kit. The 350-bp genomic library was sequenced using the Illumina HiSeq2500 platform, generating 59,844,642 paired-end reads (100 bp) with a coverage of 997×. The reads were assembled *de novo* using SPAdes version 3.10.0 (*k*-mer = 55) (4). The assembly contained 90 scaffolds (107 contigs) of ≥200 bp, covering 5,588,138 bp. The size of the genome was 5.59 Mb with a 45.5% G+C content. The three scaffolds accounted for 50% of the assembly (L_{50}), with the largest scaffold size being 1,047,310 bp. Average nucleotide identity of the LB102 whole genome was calculated using Pyani (5), a Python module, and found to be 97.9% for *V. campbellii* LMB29 strain. Gene prediction and annotation were performed using the Rapid Annotations using Subsystems Technology (RAST) server (6) and the NCBI Prokaryotic Genome Annotation Pipeline (7). The LB102 genome had total 5198 genes; 4920 protein coding, 15 rRNA, 121 tRNA, 4 other RNA and 138 pseudogenes.

The LB102 genome contained RTX toxin (cytolysin), vibriolysin (extracellular zinc protease), aerolysin (pore forming toxin), SpvB (Actin-ADP-ribosyltransferase), thermolabile hemolysin, chitinase, zinc metalloprotease, alkaline serine protease, phospholipase C, DNase, and collagenase. Secretion systems play a crucial role in the virulence

Received 21 March 2017 Accepted 27 March 2017 Published 18 May 2017

Citation Kumar S, Jangam AK, Akhil V, Rajendran V, Katneni VK, Sahaya Rajan JJ, Grover M, Nagaleekar VK, Alavandi SV, Vijayan KK. 2017. Draft genome sequence of the luminescent strain *Vibrio campbellii* LB102, isolated from a black tiger shrimp (*Penaeus monodon*) broodstock rearing system. Genome Announc 5:e00342-17. <https://doi.org/10.1128/genomeA.00342-17>.

Copyright © 2017 Kumar et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Sujeet Kumar, sujeetmicrobiol@gmail.com.

of bacterial pathogens. The genome of *V. campbellii* LB102 had secretion systems of types I, II, III, and VI. Four genes encoding homologs of *V. cholerae* toxins—VgrG-3, hemolysin coregulated protein (hcp), transcriptional activator ToxR, and transmembrane regulatory protein ToxS—were present in the genome. Both VgrG-3 and hcp are effector proteins secreted by a type VI secretion system in *V. cholerae* (8). The genome also contained an effector protein of the type III secretion system, YopT, which is homologous to the effector protein of *Yersinia pestis* (9). YopT is known to cause cytotoxicity through the depolymerization of actin filaments (9). Sequence data showed the presence of various colonization factors, such as mannose-sensitive hemagglutinin (MSHA), type IV pilin (PilA), V10 pilin, and fimbrial protein, in LB102 genome. In most of the Gram-negative bacterial pathogens, type III and type VI secretion systems are known to play major roles in their pathogenesis. Further investigation is required to understand the role of these putative effector proteins and colonization factors to understand the virulence mechanism of *V. campbellii* in shrimp.

Accession number(s). This whole-genome shotgun project has been deposited at DDBJ/ENA/GenBank under the accession number [MWRX00000000](https://doi.org/10.1093/mbe/mwz000). The version described in this paper is the first version, MWRX01000000.

ACKNOWLEDGMENT

We thank the Indian Council of Agricultural Research (ICAR), New Delhi, under the “Network Project on Agricultural Bioinformatics and Computational Biology” for financial support of this work.

REFERENCES

- Gomez-Gil B, Soto-Rodríguez S, García-Gasca A, Roque A, Vazquez-Juarez R, Thompson FL, Swings J. 2004. Molecular identification of *Vibrio harveyi*-related isolates associated with diseased aquatic organisms. *Microbiology* 150:1769–1777. <https://doi.org/10.1099/mic.0.26797-0>.
- Lin B, Wang Z, Malanoski AP, O'grady EA, Wimpee CF, Vuddhakul V, Alves N Jr, Thompson FL, Gomez-Gil B, Vora GJ. 2010. Comparative genomic analyses identify the *Vibrio harveyi* genome sequenced strains BAA-1116 and HY01 as *Vibrio campbellii*. *Environ Microbiol Rep* 2:81–89. <https://doi.org/10.1111/j.1758-2229.2009.00100.x>.
- Chrisolite B, Thiyagarajan S, Alavandi SV, Abhilash EC, Kalaimani N, Vijayan KK, Santiago TC. 2008. Distribution of luminescent *Vibrio harveyi* and their bacteriophages in a commercial shrimp hatchery in south India. *Aquaculture* 275:13–19. <https://doi.org/10.1016/j.aquaculture.2007.12.016>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
- Arahal DR. 2014. Whole-genome analyses: average nucleotide identity, p 103–122. In Goodfellow M, Sutcliffe I, Chun J (ed), *Methods in microbiology: new approaches to prokaryotic systematics*. Elsevier, Amsterdam, Netherlands. <https://doi.org/10.1016/bs.mim.2014.07.002>.
- Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: rapid annotations using subsystems technology. *BMC Genomics* 9:75. <https://doi.org/10.1186/1471-2164-9-75>.
- Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI prokaryotic genome annotation pipeline. *Nucleic Acids Res* 44: 6614–6624. <https://doi.org/10.1093/nar/gkw569>.
- Pukatzki S, Ma AT, Sturtevant D, Krastins B, Sarracino D, Nelson WC, Heidelberg JF, Mekalanos JJ. 2006. Identification of a conserved bacterial protein secretion system in *Vibrio cholerae* using the *Dictyostelium* host model system. *Proc Natl Acad Sci U S A* 103:1528–1533. <https://doi.org/10.1073/pnas.0510322103>.
- Iriarte M, Cornelis GR. 1998. YopT, a new *Yersinia* Yop effector protein, affects the cytoskeleton of host cells. *Mol Microbiol* 29:915–929. <https://doi.org/10.1046/j.1365-2958.1998.00992.x>.