



Complete Genome Sequence of an Emergent Recombinant GII.P16-GII.2 Norovirus Strain Associated with an Epidemic Spread in the Winter of 2016-2017 in Hong Kong, China

 Martin C. W. Chan,^a Kirsty Kwok,^a Tin-Nok Hung,^a Lok-Yi Chan,^a Paul K. S. Chan^{a,b}

Department of Microbiology, Faculty of Medicine, Chinese University of Hong Kong, Hong Kong, China^a; Stanley Ho Centre for Emerging Infectious Diseases, Faculty of Medicine, Chinese University of Hong Kong, Hong Kong, China^b

ABSTRACT A new recombinant norovirus, GII.P16-GII.2, emerged in the winter of 2016-2017. Here, we report the complete genome of this strain (Hu/GII/HK/2016/GII.P16-GII.2/CUHK-NS-1082), which was collected from a patient hospitalized with gastroenteritis in September 2016 in Hong Kong, China, and sequenced using next-generation sequencing. This genome had a 95.2% nucleotide identity to the closest sequence in GenBank.

Human noroviruses comprise a diverse group of enteric RNA viruses that cause a majority of acute gastroenteritis in all age groups worldwide (1). These viruses are genetically very diverse and are currently classified into at least 30 genotypes based on viral capsid protein sequences (2). Genogroup II genotype 4 (GII.4) strains have been predominant since 2002, and new epidemic variants have emerged every 2 to 3 years (3). In the past 3 years, novel non-GII.4 viruses are continuing to emerge, including a new GII.17 Kawasaki variant in the winter of 2014/2015 and a recombinant GII.P16-GII.2 variant in the winter of 2016-2017 (4). Here, we report the complete genome of a GII.P16-GII.2 strain that caused an epidemic spread in the winter of 2016-2017 in Hong Kong, China.

This recombinant norovirus strain (GII/Hu/HK/2016/GII.P16-GII.2/CUHK-NS-1082) was detected as part of our norovirus surveillance (5) in a stool specimen collected from an 8-year-old boy hospitalized for acute gastroenteritis in September 2016 in Hong Kong, China. Preliminary virus capsid genotyping was performed as previously described (6). The complete genome of the virus was determined using a metagenomic next-generation sequencing approach. Briefly, 10% (wt/vol) stool suspension in phosphate-buffered saline was filtered and digested with a cocktail of DNase, RNase, and benzonase to remove cell-free, unprotected host and microbial DNA and RNA. Viral RNA protected in intact virions was extracted and purified using the Viral RNA mini kit (Qiagen). Reverse transcription was performed using tagged random octamers and SuperScript III reverse transcriptase (Thermo Fisher). Second-strand cDNA was synthesized by Klenow fragment (TaKaRa), followed by random cDNA amplification using high-fidelity Phusion Hot Start II DNA polymerase (Thermo Fisher). The library for next-generation sequencing was prepared using the Nextera XT DNA library prep kit (Illumina). Purification between steps was performed using AMPure magnetic beads (Beckman Coulter, Inc.). Paired-end 2 × 75-bp sequencing was performed on the MiSeq system (Illumina). Protocols are available upon request.

Received 21 March 2017 **Accepted** 30 March 2017 **Published** 18 May 2017

Citation Chan MCW, Kwok K, Hung T-N, Chan L-Y, Chan PKS. 2017. Complete genome sequence of an emergent recombinant GII.P16-GII.2 norovirus strain associated with an epidemic spread in the winter of 2016-2017 in Hong Kong, China. *Genome Announc* 5: e00343-17. <https://doi.org/10.1128/genomeA.00343-17>.

Copyright © 2017 Chan et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Paul K. S. Chan, paulkschan@cuhk.edu.hk.

A total of 1,025,446 reads were generated. Quality trimming of reads was performed using Trimmomatic version 0.36, and the survival rate was 97.4%. Reference sequence mapping to a recent GII.2 strain (Hu/GII.2/Miyahi1_2012_JP; GenBank accession no. LC145787) was performed on Geneious version 9.1.6, and the percentage of on-target reads was 66.7%. The mean sequencing depth was $6,762\times$ (range 1 to $38,477\times$). The terminal 5' and 3' ends of the virus genome with low coverage ($<3\times$) were validated by Sanger sequencing.

The complete genome of the recombinant GII.P16-GII.2 was 7,536 nucleotides in length, excluding the poly(A) tail. The best BLAST hit in GenBank was a norovirus GII.2 strain collected in 2011 from the United States (Hu/GII.2/HS255/2011/USA; KJ407074) with a 95.2% identity on the nucleotide level. Interestingly, open reading frame 1 (ORF1) of this new GII.2 variant had the highest pairwise identity (98.5% on the nucleotide level and 99.4% on the amino acid level) to a circulating recombinant GII.P16-GII.4 Sydney 2012 strain (Hu/USA/2015/GII.P16_GII.4_Sydney/CA3477; KX907727). It would be interesting to postulate that this emergent GII.2 originated from a recombination event between a GII.4 and a GII.2 virus. Furthermore, this new GII.2 strain contained a one-to-two-codon insertion in the N-terminal protein of ORF1 compared with older GII.2 strains. The functional importance of these insertions merits further investigation.

Accession number(s). The complete genome sequence of the recombinant norovirus GII.P16-GII.2 strain (GII/Hu/HK/2016/GII.P16-GII.2/CUHK-NS-1082) has been deposited in GenBank under the accession number [KY771081](https://www.ncbi.nlm.nih.gov/nuccore/KY771081).

ACKNOWLEDGMENTS

We thank the Core Utilities of Cancer Genomics and Pathobiology (CUHK) for providing the facilities and assistance in support of this research on next-generation sequencing. M.C.W.C. and P.K.S.C. conceived the study. M.C.W.C. designed and supervised the study. K.K., T.-N.H., and L.-Y.C. performed experiments. M.C.W.C. analyzed data and drafted the manuscript. All authors critically reviewed the manuscript and approved the final revision. M.C.W.C. has access to all data and is responsible for the scientific integrity of this work.

This study was supported in part by departmental research funds and by the Commissioned Health and Medical Research Fund (phase 3) of the Food and Health Bureau of the HKSAR Government (to M.C.W.C.; reference no. CU-15-C2).

REFERENCES

- Ahmed SM, Hall AJ, Robinson AE, Verhoef L, Premkumar P, Parashar UD, Koopmans M, Lopman BA. 2014. Global prevalence of norovirus in cases of gastroenteritis: a systematic review and meta-analysis. *Lancet Infect Dis* 14:725–730. [https://doi.org/10.1016/S1473-3099\(14\)70767-4](https://doi.org/10.1016/S1473-3099(14)70767-4).
- Vinje J. 2015. Advances in laboratory methods for detection and typing of norovirus. *J Clin Microbiol* 53:373–381. <https://doi.org/10.1128/JCM.01535-14>.
- Pringle K, Lopman B, Vega E, Vinje J, Parashar UD, Hall AJ. 2015. Noroviruses: epidemiology, immunity and prospects for prevention. *Future Microbiol* 10:53–67. <https://doi.org/10.2217/fmb.14.102>.
- Niendorf S, Jacobsen S, Faber M, Eis-Hubinger AM, Hofmann J, Zimmermann O, Hohne M, Bock CT. 2017. Steep rise in norovirus cases and emergence of a new recombinant strain GII.P16-GII.2, Germany, winter 2016. *Euro Surveill* 22:30447. <https://doi.org/10.2807/1560-7917.ES.2017.22.4.30447>.
- Chan MC, Lee N, Hung TN, Kwok K, Cheung K, Tin EK, Lai RW, Nelson EA, Leung TF, Chan PK. 2015. Rapid emergence and predominance of a broadly recognizing and fast-evolving norovirus GII.17 variant in late 2014. *Nat Commun* 6:10061. <https://doi.org/10.1038/ncomms10061>.
- Chan MC, Leung TF, Chung TW, Kwok AK, Nelson EA, Lee N, Chan PK. 2015. Virus genotype distribution and virus burden in children and adults hospitalized for norovirus gastroenteritis, 2012–2014, Hong Kong. *Sci Rep* 5:11507. <https://doi.org/10.1038/srep11507>.