



# Draft Genome Sequence of *Byssochlamys* sp. Isolate BYSS01, a Filamentous Fungus Adapted to the Fuel Environment

Osman Radwan,<sup>a</sup> Thusitha S. Gunasekera,<sup>a</sup>  Oscar N. Ruiz<sup>b</sup>

<sup>a</sup>Environmental Microbiology Group, University of Dayton Research Institute, Dayton, Ohio, USA

<sup>b</sup>Fuels and Energy Branch, Aerospace Systems Directorate, Air Force Research Laboratory, Wright-Patterson AFB, Ohio, USA

**ABSTRACT** *Byssochlamys* sp. isolate BYSS01 (anamorph, *Paecilomyces* sp.), which was isolated from jet fuel, is highly adapted to grow in hydrocarbons, having predicted genes involved in degradation of *n*-alkanes, branched alkanes, and aromatic compounds. The draft genome size is estimated at 29 Mb, containing 8,509 proteins.

The filamentous fungus *Byssochlamys* sp. isolate BYSS01 was recovered from Jet A fuel and identified based on morphological characteristics and sequence similarity (94%) to the 18S rRNA gene of *Byssochlamys fulva*. *Byssochlamys* sp. (teleomorph) belongs to the genus *Paecilomyces*, which comprises formaldehyde-resistant and heat-resistant food spoilage fungi (1–4). It was confirmed that the BYSS01 isolate metabolizes alkanes and aromatic hydrocarbons efficiently; this is the first report of a *Byssochlamys* species adapted to kerosene fuel. Therefore, to gain a deeper understanding of the adaptive mechanisms in isolate BYSS01, its genome was sequenced.

A whole-genome shotgun (WGS) approach on a HiSeq 2000 platform was used to generate TruSeq paired-end libraries, resulting in 47,245,046 paired-end reads (150× sequence coverage), with a read length of 100 bp (~4.72 Gb). The raw sequences were trimmed using Trimmomatic (5), and reads shorter than 50 bp were removed. The sequence reads were *de novo* assembled with SPAdes software (6), generating a draft assembly comprised of 394 scaffolds greater than 500 bp, with an  $N_{50}$  value of 463,366, an  $L_{50}$  value of 20 scaffolds, and a G+C content of 49.3%. BUSCO (7) identified 1,425 out of 1,438 ultraconserved eukaryotic genes in *Byssochlamys* sp. isolate BYSS01 (99.1%). Repetitive sequences (3.09%) were masked using the RepeatMasker program (8), and the masked genome was used for gene prediction by Augustus 2.5.5 (9) with an option set for *Aspergillus oryzae* parameters, resulting in 8,509 protein-encoding genes. The average gene density is one gene per 1.67 kb, with an average of 3.45 exons per gene. Approximately 21,244 introns, ranging from 34 to 6,725 bp in length, are present in the genome, with 140 bp as an average intron size and 2.5 as the average number of introns per open reading frame.

A BLASTP search against UniProt and the annotated genomes of *Hormoconis resinae*, *Aspergillus fumigatus*, and *Byssochlamys spectabilis* (4) revealed significant matches (E value,  $1 \times 10^{-5}$ ) for 6,646, 7,578, 7,808, and 7,508 proteins, respectively. Using the Carbohydrate-Active Enzymes (CAZY) database (10), a total of 334 fungal enzymes involved in carbohydrate metabolism and assimilation were identified (E value,  $1 \times 10^{-4}$ ), including glycoside hydrolases ( $n = 113$ ), carbohydrate esterases ( $n = 77$ ), glycosyl transferases ( $n = 76$ ), enzymes involved in auxiliary activities ( $n = 46$ ), carbohydrate-binding modules ( $n = 21$ ), and polysaccharide lyases ( $n = 1$ ). The Transporter Classification Database (TCDB; <http://www.tcdb.org/>) identified 283 major facilitator superfamily (MSF) transporters with an E value of  $1 \times 10^{-5}$ , reflecting the ability of the BYSS01 isolate to extrude toxic compounds; efflux pumps have been shown to aid in adaptation to hydrocarbons in bacteria (11, 12).

Received 6 February 2018 Accepted 13 February 2018 Published 8 March 2018

**Citation** Radwan O, Gunasekera TS, Ruiz ON. 2018. Draft genome sequence of *Byssochlamys* sp. isolate BYSS01, a filamentous fungus adapted to the fuel environment. Genome Announc 6:e00164-18. <https://doi.org/10.1128/genomeA.00164-18>.

**Copyright** © 2018 Radwan et al. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International license.

Address correspondence to Oscar N. Ruiz, [oscar.ruiz@us.af.mil](mailto:oscar.ruiz@us.af.mil).

The Kyoto Encyclopedia of Genes and Genomes (KEGG) database and BLASTP searches identified important proteins involved in carbon and energy metabolism ( $n = 49$ ). In agreement with the ability of BYSS01 to grow in fuel, genes involved in the degradation of aromatics and  $n$ -alkanes, including cytochrome P450 alkane hydroxylase, cytochrome P450 monooxygenase, aromatic ring-opening dioxygenase, salicylate hydroxylase, 2-haloacid dehalogenase, benzyl alcohol dehydrogenase, benzoate 4-monooxygenase, and dimethylsulfide monooxygenase, were found. This genome sequence will help to understand the adaptive mechanisms employed by fungi to survive and proliferate in hydrocarbon-based fuels.

**Accession number(s).** This whole-genome shotgun project has been deposited at DDBJ/ENA/GenBank under the accession number [NIXA00000000](https://www.ncbi.nlm.nih.gov/nuclseq/NIXA00000000).

## ACKNOWLEDGMENT

This material is based on research sponsored by AFRL/RQTF under agreement number FA8650-16-2-2605.

The U.S. Government is authorized to reproduce and distribute reprints for Governmental purposes notwithstanding any copyright notation thereon. The views and conclusions contained herein are those of the authors and should not be interpreted as necessarily representing the official policies or endorsements, either expressed or implied, of AFRL/RQTF or the U.S. Government.

## REFERENCES

- Kotzekidou P. 1997. Heat resistance of *Byssoschlamys nivea*, *Byssoschlamys fulva* and *Neosartorya fischeri* isolated from canned tomato paste. *J Food Science* 62:410–413. <https://doi.org/10.1111/j.1365-2621.1997.tb04014.x>.
- Iwahara M, Fukuda R, Nakahara K, Tanaka T, Nomura Y. 2001. Isolation and properties of *Paecilomyces* sp. no. 5 capable of degrading high concentrations of formaldehyde. *BioControl Sci* 7:107–110.
- Fukuda R, Nagahama K, Fukuda K, Ekino K, Oka T, Nomura Y. 2012. Purification and properties of *S*-hydroxymethylglutathione dehydrogenase of *Paecilomyces variotii* no. 5, a formaldehyde-degrading fungus. *AMB Express* 2:32. <https://doi.org/10.1186/2191-0855-2-32>.
- Oka T, Ekino K, Fukuda K, Nomura Y. 2014. Draft genome sequence of the formaldehyde-resistant fungus *Byssoschlamys spectabilis* no. 5 (anamorph *Paecilomyces variotii* no. 5) (NBRC109023). *Genome Announc* 2:e01162-13. <https://doi.org/10.1128/genomeA.01162-13>.
- Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
- Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. 2015. BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics* 31:3210–3212. <https://doi.org/10.1093/bioinformatics/btv351>.
- Smit AFA, Hubley R, Green P. RepeatMasker Open-3.0. 1996–2010. <http://www.repeatmasker.org>.
- Stanke M, Steinkamp R, Waack S, Morgenstern B. 2004. AUGUSTUS: a web server for gene finding in eukaryotes. *Nucleic Acids Res* 32:W309–W312. <https://doi.org/10.1093/nar/gkh379>.
- Lombard V, Ramulu HG, Drula E, Coutinho PM, Henrissat B. 2014. The carbohydrate-active enzymes database (CAZy) in 2013. *Nucleic Acids Res* 42:D490–D495. <https://doi.org/10.1093/nar/gkt1178>.
- Gunasekera TS, Bowen LL, Zhou CE, Howard-Byerly SC, Foley WS, Striebich RC, Dugan LC, Ruiz ON. 2017. Transcriptomic analyses elucidate adaptive differences of closely related strains of *Pseudomonas aeruginosa* in fuel. *Appl Environ Microbiol* 83:e03249-16. <https://doi.org/10.1128/AEM.03249-16>.
- Ruiz ON, Brown LM, Striebich RC, Smart CE, Bowen LL, Lee JS, Little BJ, Mueller SS, Gunasekera TS. 2016. Effect of conventional and alternative fuels on a marine bacterial community and the significance to bioremediation. *Energy Fuels* 30:434–444. <https://doi.org/10.1021/acs.energyfuels.5b02439>.