



Partial Mitochondrial Genome Sequences of Two Abyssal Sponges (Porifera: Hexactinellida), *Bathydorus laniger* and *Docosaccus maculatus*

 Amanda S. Kahn,^{a*} Jonathan B. Geller^a

^aMoss Landing Marine Laboratories, Moss Landing, California, USA

ABSTRACT We announce the nearly complete mitochondrial genome sequences of two hexactinellid sponges, *Bathydorus laniger* and *Docosaccus maculatus*. A contiguous region of over 15,000 bp was sequenced from each genome. An uncommon structural element was identified as a series of repetitive elements with sequences matching *cob* in the genome of *D. maculatus*.

We sequenced the nearly complete mitochondrial genomes of two abyssal sponges (phylum Porifera), bringing the total coverage to 13 hexactinellid genomes. Tissue samples from the holotypes of *Bathydorus laniger* Kahn et al. 2013 and *Docosaccus maculatus* Kahn et al. 2013 were collected in 2007 from Station M, 200 km west of Point Conception, CA (4,100 m depth [1]). Tissue samples were frozen in liquid nitrogen and stored at -80°C . DNA was extracted under liquid nitrogen according to DNeasy animal and blood tissue extraction kit protocols (Qiagen, USA). Mitochondrial DNA (mtDNA) was sequenced using primer walking with standard PCR, long PCR (2), and restriction digestion and cloning. Sequences were assembled using Geneious version 5.3 (3). Coding regions were identified using BLAST searches (GenBank), followed by a comparison of gene translations in Geneious. tRNA sequences were predicted using tRNAscan-SE (4).

We sequenced 15,704 bp for *Bathydorus laniger*, 14,709 bp of which (93.7%) represents coding regions; the total A+T content was 71.3%. This included 29 genes, with 12 protein-coding genes (three units of cytochrome oxidase [*cox1*, *cox2*, and *cox3*], six subunits of NADH dehydrogenase [*nad1*, *nad2*, *nad3*, *nad4*, *nad4L*, and *nad5*], cytochrome *b* [*cob*], and two subunits of ATP synthase [*atp6* and *atp9*]), small (*rns*) and large (*rnl*) subunit rRNAs, and 15 tRNAs. The largest noncoding space was 262 bp long, with no notable elements.

The coverage of the mitochondrial genome of *Docosaccus maculatus* was similar, with a single contiguous sequence of 17,143 bp, of which 15,212 bp (88.7%) represents coding regions; the A+T content was 70.9%. There were 32 genes, with the same 13 protein-coding genes as *B. laniger* plus NADH dehydrogenase subunit 6 (*nad6*), *rns*, and *rnl*, and 17 tRNAs.

In both species, most genes used ATG as a start codon, but the start codon for *atp9* was ATA. The stop codons were either TAA or TAG. In *D. maculatus*, a +1 frameshift occurred in *nad2* (amino acid position 63). The insertion was preceded by the UGG codon for tryptophan, which was present only there and once in *cox1* of *B. laniger*; all other tryptophan residues were encoded by the more common UGA codon. Interestingly, the UGG codon has been associated with a +1 frameshift in coding regions of other glass sponges (5–7).

Noncoding mtDNA for *D. maculatus* was concentrated in a 1,439-bp region between *nad1* and *cob*. It contained two stuttering repeats of *trnY* and the 5' end of a partial *cob* sequence upstream of a functional *cob* gene. The region also included two nonfunc-

Received 27 February 2018 Accepted 22 March 2018 Published 19 April 2018

Citation Kahn AS, Geller JB. 2018. Partial mitochondrial genome sequences of two abyssal sponges (Porifera: Hexactinellida), *Bathydorus laniger* and *Docosaccus maculatus*. Genome Announc 6:e00234-18. <https://doi.org/10.1128/genomeA.00234-18>.

Copyright © 2018 Kahn and Geller. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Amanda S. Kahn, kahn@ualberta.ca.

* Present address: Amanda S. Kahn, Department of Biological Sciences, University of Alberta, Edmonton, Alberta, Canada.

tional copies of *trnY (gua)* and a third functional gene (predicted using tRNAscan-SE [4]) which differed by only a few base pairs from the other two. Repetitive regions have been found in demosponges as palindromic repeats (8–10) but have not been observed in hexactinellid species, and stuttering repeats of protein-coding genes have not been observed in any other sponge genomes; the feature is so far unique to *D. maculatus*.

Accession number(s). Sequences have been deposited in DDBJ/ENA/GenBank under the accession numbers [KJ634155](#) (*Bathydorus laniger*) and [KJ634156](#) (*Docosaccus maculatus*).

ACKNOWLEDGMENTS

K. L. Smith, H. A. Ruhl, J. Ellena, M. Vardaro, the crew of the research vehicle (R/V) *Western Flyer*, and pilots of the remotely operated vehicles (ROVs) *Tiburon* and *Doc Ricketts* were responsible for the collection of sponge specimens. We thank N. B. Webster, G. Cailliet, K. L. Smith, and H. Hawk for reviewing the manuscript and providing constructive feedback.

Funding came from student support from the Marine Technology Society, the Dr. James Nybakken Scholarship administered by the Friends of MLML, and the David and Lucille Packard Foundation.

The funders had no role in the study design, data collection and interpretation, or the decision to submit the work for publication.

REFERENCES

- Smith KL, Jr, Druffel ERM. 1998. Long time-series monitoring of an abyssal site in the NE Pacific: an introduction. *Deep Sea Res II* 45: 573–586. [https://doi.org/10.1016/S0967-0645\(97\)00094-5](https://doi.org/10.1016/S0967-0645(97)00094-5).
- Burger G, Lavrov DV, Forget L, Lang BF. 2007. Sequencing complete mitochondrial and plastid genomes. *Nat Protoc* 2:603–614. <https://doi.org/10.1038/nprot.2007.59>.
- Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Thierer T, Ashton B, Meintjes P, Drummond A. 2012. Geneious basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 28:1647–1649. <https://doi.org/10.1093/bioinformatics/bts199>.
- Lowe TM, Eddy SR. 1997. tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res* 25:955–964.
- Rosengarten RD, Sperling EA, Moreno MA, Leys SP, Dellaporta SL. 2008. The mitochondrial genome of the hexactinellid sponge *Aphrocallistes vastus*: evidence for programmed translational frameshifting. *BMC Genomics* 9:33. <https://doi.org/10.1186/1471-2164-9-33>.
- Haen KM, Lang BF, Pomponi SA, Lavrov DV. 2007. Glass sponges and bilaterian animals share derived mitochondrial genomic features: a common ancestry or parallel evolution? *Mol Biol Evol* 24:1518–1527. <https://doi.org/10.1093/molbev/msm070>.
- Haen KM, Pett W, Lavrov DV. 2013. Eight new mtDNA sequences of glass sponges reveal an extensive usage of +1 frameshifting in mitochondrial translation. *Gene* 535:336–344. <https://doi.org/10.1016/j.gene.2013.10.041>.
- Wang X, Lavrov DV. 2007. Mitochondrial genome of the homoscleromorph *Oscarella carmela* (Porifera, Demospongiae) reveals unexpected complexity in the common ancestor of sponges and other animals. *Mol Biol Evol* 24:363–373. <https://doi.org/10.1093/molbev/msl167>.
- Lukić-Bilela L, Brandt D, Pojskić N, Wiens M, Gamulin V, Müller WEG. 2008. Mitochondrial genome of *Suberites domuncula*: palindromes and inverted repeats are abundant in non-coding regions. *Gene* 412:1–11. <https://doi.org/10.1016/j.gene.2008.01.001>.
- Lavrov DV. 2010. Rapid proliferation of repetitive palindromic elements in mtDNA of the endemic Baikalian sponge *Lubomirskia baicalensis*. *Mol Biol Evol* 27:757–760. <https://doi.org/10.1093/molbev/msp317>.