



# Complete Chromosome and Plasmid Sequences of Two Plant Pathogens, *Dickeya solani* Strains D s0432-1 and PPO 9019

Slimane Khayi,<sup>a,b</sup> Pauline Blin,<sup>a</sup> Teik Min Chong,<sup>c</sup>  Kok-Gan Chan,<sup>d,e</sup> Denis Faure<sup>a</sup>

<sup>a</sup>Institute for Integrative Biology of the Cell (I2BC), CEA CNRS Université Paris-Sud, Université Paris-Saclay, Gif-sur-Yvette, France

<sup>b</sup>Biotechnology Research Unit, National Institute for Agronomic Research (INRA), Rabat, Morocco

<sup>c</sup>Division of Genetics and Molecular Biology, Institute of Biological Sciences, Faculty of Science, University of Malaya, Kuala Lumpur, Malaysia

<sup>d</sup>International Genome Centre, Jiangsu University, Zhenjiang, China

<sup>e</sup>ISB, Faculty of Science, University of Malaya, Kuala Lumpur, Malaysia

**ABSTRACT** *Dickeya solani* species are emerging bacterial pathogens of *Solanum tuberosum*. Here, we announce the complete genome sequences of two strains, *Dickeya solani* D s0432-1 and PPO 9019. Strain PPO 9019 represents the first described member of the genus *Dickeya* with an extrachromosomal genetic element.

The genera *Dickeya* and *Pectobacterium* encompass several bacterial species causing blackleg and soft-rot diseases on economically important crops and ornamental plants around the world (1). In Europe, *Pectobacterium* spp. and *Dickeya dianthicola* are the most prevalent pathogens associated with the soft-rot and blackleg symptoms of potato cultures. However, since the 2000s, a novel species named *Dickeya solani* has emerged. Its emergence was associated with an increased incidence of soft-rot and blackleg diseases in several European countries and in the Middle East (2).

The whole genomes of *D. solani* strains D s0432-1 and PPO 9019 were subjected to Pacific Bioscience (PacBio) sequencing technology to highlight their complete nucleotide sequences. Draft genome sequences of these two strains, obtained using other DNA sequencing technologies, were published previously (3, 4). The genomic DNA was extracted using the MasterPure DNA purification kit (Epicentre, Inc., USA) following the manufacturer's recommendations. Quantification and quality control of the DNA were completed using a NanoDrop device (ND 1000), a Qubit 2.0 fluorometer, and agarose gel electrophoresis at 1.0% (wt/vol). The library insert size was 10 kbp. Before the assembly step, short reads that were less than 500 bp were filtered, and the minimum polymerase read quality used for mapping the subreads from a single zero-mode waveguide was set to 0.75. Totals of 340,825 filtered reads ( $N_{50}$ , 4,976 bp) for *D. solani* PPO 9019 and 287,091 filtered reads ( $N_{50}$ , 4,319 bp) for *D. solani* D s0432-1 were used for genome assembly.

The assembly was sequenced using RS\_HGAP\_Assembly software v2.0. The cutoff lengths of seed bases were set at 10,117 bp for strain PPO 9019 and 8,999 bp for strain D s0432-1 in order to serve as a reference for the recruitment of shorter reads for preassembly. The resulting consensus accuracy based on multiple sequence alignment of the subreads was at 99.99%. Polished assembly resulted in one contig for *D. solani* D s0432-1 with a length of 4,919,812 bp and 160 $\times$  coverage. Two contigs were obtained from the assembly for the *D. solani* strain PPO 9019, measuring 4,918,850 bp for the chromosome and 43,580 bp for the plasmid, with an average coverage of 220 $\times$ .

The genome sequences for both strains were submitted for annotation to the NCBI Prokaryotic Genome Annotation Pipeline (released 2013).

Received 27 February 2018 Accepted 22 March 2018 Published 26 April 2018

**Citation** Khayi S, Blin P, Chong TM, Chan K-G, Faure D. 2018. Complete chromosome and plasmid sequences of two plant pathogens, *Dickeya solani* strains D s0432-1 and PPO 9019. Genome Announc 6:e00233-18. <https://doi.org/10.1128/genomeA.00233-18>.

**Copyright** © 2018 Khayi et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Denis Faure, [denis.faire@i2bc.paris-saclay.fr](mailto:denis.faire@i2bc.paris-saclay.fr).

**Accession number(s).** The complete genome sequences for these bacteria have been deposited at DDBJ/EMBL/GenBank under the accession numbers [CP017454](#) and [CP017455](#) (*Dickeya solani* strain PPO 9019) and [CP017453](#) (*Dickeya solani* strain D s0432-1).

## ACKNOWLEDGMENTS

This work is supported by CNRS (grant I2BC-SB2015), Agence Nationale de la Recherche (project ANR-15-CE21-0003), a High Impact Research Grant (UM.C/625/1/ HIR/MOHE/CHAN/14/01, grant A-000001-50001), and the French-Malaysian exchange program awarded by the French Embassy of Malaysia.

## REFERENCES

1. Samson R, Legendre JB, Christen R, Fischer-Le Saux M, Achouak W, Gardan L. 2005. Transfer of *Pectobacterium chrysanthemi* (Burkholder et al. 1953) Brenner et al. 1973 and *Brenneria paradisiaca* to the genus *Dickeya* gen. nov. as *Dickeya chrysanthemi* comb. nov. and *Dickeya paradisiaca* comb. nov. and delineation of four novel species, *Dickeya dadantii* sp. nov., *Dickeya dianthicola* sp. nov., *Dickeya dieffenbachiae* sp. nov. and *Dickeya zea* sp. nov. *Int J Syst Evol Microb* 55:1415–1427. <https://doi.org/10.1099/ijs.0.02791-0>.
2. Toth IK, van der Wolf JM, Saddler G, Lojkowska E, Hélias V, Pirhonen M, Tsrör Lahkim L, Elphinstone JG. 2011. *Dickeya* species: an emerging problem for potato production in Europe. *Plant Pathol* 60:385–399. <https://doi.org/10.1111/j.1365-3059.2011.02427.x>.
3. Garland L, Koskinen P, Rouhiainen L, Laine P, Paulin L, Auvinen P, Holm L, Pirhonen M. 2013. Genome sequence of *Dickeya solani*, a new soft rot pathogen of potato, suggests its emergence may be related to a novel combination of non-ribosomal peptide/polyketide synthetase clusters. *Diversity* 5:824–842. <https://doi.org/10.3390/d5040824>.
4. Khayi S, Blin P, Pédrón J, Chong TM, Chan KG, Moumni M, Hélias V, Van Gijsegem F, Faure D. 2015. Population genomics reveals additive and replacing horizontal gene transfers in the emerging pathogen *Dickeya solani*. *BMC Genomics* 16:788. <https://doi.org/10.1186/s12864-015-1997-z>.