



Draft Genome Sequence of *Tuber borchii* Vittad., a Whitish Edible Truffle

Claude Murat,^a Alan Kuo,^b Kerrie W. Barry,^b Alicia Clum,^b Rhyan B. Dockter,^b Laure Fauchery,^a Mirco Iotti,^c Annegret Kohler,^a Kurt LaButti,^b Erika A. Lindquist,^b Anna Lipzen,^b Emmanuelle Morin,^a Mei Wang,^b Igor V. Grigoriev,^b Alessandra Zambonelli,^d Francis M. Martin^a

^aUMR1136 Interactions Arbres-Microorganismes, Laboratoire d'Excellence ARBRE, INRA, Université de Lorraine, Champenoux, France

^bU.S. Department of Energy Joint Genome Institute, Walnut Creek, California, USA

^cDepartment of Life, Health and Environmental Sciences, University of L'Aquila, L'Aquila, Italy

^dDepartment of Agricultural Science, University of Bologna, Bologna, Italy

ABSTRACT The ascomycete *Tuber borchii* (Pezizomycetes) is a whitish edible truffle that establishes ectomycorrhizal symbiosis with trees and shrubs. This fungus is ubiquitous in Europe and is also cultivated outside Europe. Here, we present the draft genome sequence of *T. borchii* strain Tbo3840 (97.18 Mb in 969 scaffolds, with 12,346 predicted protein-coding genes).

The whitish truffle (*Tuber borchii* Vittad.) is an ectomycorrhizal ascomycete ubiquitous in Europe growing from cold temperate to Mediterranean climates (1). *T. borchii* is found in subalkaline and acid soils associated with a wide range of hosts, including oak, poplar, strawberry, and pine trees (1, 2). Seedlings inoculated with *T. borchii* have been available since 1969 (3), and this species is also cultivated outside Europe in Australia, the United States, and New Zealand (4). Before the release of the genome sequence of the black truffle of Périgord (*Tuber melanosporum*) (5), *T. borchii* was the most studied truffle species, as it is amenable to laboratory manipulations (6).

Tuber borchii strain Tbo3840 was isolated from a fruiting body harvested in April 2010 at Guzzano (Pianoro) in central Italy. For the genome sequencing, the vegetative haploid mycelium was grown in liquid modified Woody plant medium (7) for 2 months at 22°C in the dark. DNA was extracted from 2 g of mycelium by using a modified cetyltrimethylammonium bromide (CTAB) protocol (8). The genome sequencing was performed using Pacific Biosciences RS II version C4 chemistry. The genome assembly was generated with Falcon version 0.4.2 (9), improved with FinisherSC version 2.0 (10), and polished with Quiver (11). Contigs shorter than 1,000 bp were excluded from the assembly. RNA was extracted from 40 mg of mycelium ground in liquid nitrogen with the RNeasy plant minikit (Qiagen, Hilden, German), according to the provider's recommendations, and using the RLC lysis buffer. The RNA quality and quantity were checked with the Experion RNA StdSens analysis kit (Bio-Rad, Hercules, CA). Three micrograms of RNA was used for RNA sequencing using the Illumina HiSeq 2500 platform and *de novo* assembled using Rnnotator (12). The genome was annotated using the JGI annotation pipeline (13, 14).

The nuclear genome assembly is 97.18 Mb long in 969 scaffolds (N_{50} , 0.19 Mbp), with 47.12% of repeated sequences (<http://genome.jgi.doe.gov/Tubbor1>). *T. borchii* has a reduced genome size compared to other truffle species, such as *Tuber aestivum* (145 Mb) (15), *Tuber magnatum* (192 Mb) (15), and *T. melanosporum* (125 Mb) (5). A total of 12,346 protein-coding genes were predicted, with an average length of 1,521 bp and 379 amino acids for genes and proteins, respectively. This number of gene models is higher than that in *T. melanosporum* (7,496 genes annotation version 1.0) (5) but similar

Received 11 May 2018 Accepted 13 May 2018 Published 21 June 2018

Citation Murat C, Kuo A, Barry KW, Clum A, Dockter RB, Fauchery L, Iotti M, Kohler A, LaButti K, Lindquist EA, Lipzen A, Morin E, Wang M, Grigoriev IV, Zambonelli A, Martin FM. 2018. Draft genome sequence of *Tuber borchii* Vittad., a whitish edible truffle. Genome Announc 6: e00537-18. <https://doi.org/10.1128/genomeA.00537-18>.

This is a work of the U.S. Government and is not subject to copyright protection in the United States. Foreign copyrights may apply.

Address correspondence to Claude Murat, claude.murat@inra.fr, or Francis M. Martin, francis.martin@inra.fr.

to that of the closely related peizomycete *Pyronema confluens* (13,367 genes) (16). As found in *T. melanosporum*, glycoside hydrolase 6 (GH6) and GH7 are absent from the *T. borchii* genome, confirming that ectomycorrhizal species have a reduced content of plant cell wall-degrading enzymes (5, 8). The haploid genome of *T. borchii* Tbo3840 exhibits the mating type locus MAT1-2-1 with HMG box (protein identification [ID] 1092840); the other mating type locus was not detected, as expected for a heterothallic species (6, 17).

Accession number(s). This whole-genome shotgun project has been deposited at DDBJ/ENA/GenBank under the accession number [NESQ00000000](https://www.ncbi.nlm.nih.gov/nuccore/NESQ00000000). The version described in this paper is version NESQ01000000. All of the sequences are also available at the JGI-DOE MycoCosm portal (13) (<http://genome.jgi.doe.gov/Tubbor1>).

ACKNOWLEDGMENTS

The work conducted by the U.S. Department of Energy Joint Genome Institute, a DOE Office of Science User Facility, was supported by the Office of Science of the U.S. Department of Energy under contract DE-AC02-05CH11231. The UMR1136 is supported by a grant overseen by the French National Research Agency (ANR) as part of the “Investissements d’Avenir” program (ANR-11-LABX-0002-01, Lab of Excellence ARBRE).

REFERENCES

- Lancellotti E, Franceschini A. 2013. Studies on the ectomycorrhizal community in a declining *Quercus suber* L. stand. *Mycorrhiza* 23:533–542. <https://doi.org/10.1007/s00572-013-0493-z>.
- Lancellotti E, Iotti M, Zambonelli A, Franceschini A. 2016. The *Puberulum* group sensu lato (whitish truffles), p 105–124. In Zambonelli A, Iotti M, Murat C (ed), True truffle (*Tuber* spp.) in the world. Springer International Publishing, Cham, Switzerland.
- Fontana A, Palenzona M. 1969. Sintesi micorrizica di tuber albidum in coltura pura, con *Pinus strobus* e pioppo euroamericano. *Allionia* 15: 100–104.
- Zambonelli A, Iotti M, Hall I. 2015. Current status of truffle cultivation: recent results and future perspectives. *Ital J Mycol* 44:31–40. <https://doi.org/10.6092/issn.2465-311X/5593>.
- Martin F, Kohler A, Murat C, Balestrini R, Coutinho PM, Jaillon O, Montanini B, Morin E, Noel B, Percudani R, Porcel B, Rubini A, Amicucci A, Amsellem J, Anthouard V, Arcioni S, Artiguenave F, Aury J-M, Ballario P, Bolchi A, Brenna A, Brun A, Buée M, Cantarel B, Chevalier G, Couloux A, Da Silva C, Denoeud F, Duplessis S, Ghignone S, Hilselberger B, Iotti M, Marçais B, Mello A, Miranda M, Pacioni G, Quesneville H, Riccioni C, Ruotolo R, Splivallo R, Stocchi V, Tisserant E, Viscomi AR, Zambonelli A, Zampieri E, Henrissat B, Lebrun M-H, Paolocci F, Bonfante P, Ottonello S, et al. 2010. Périgord black truffle genome uncovers evolutionary origins and mechanisms of symbiosis. *Nature* 464:1033–1038. <https://doi.org/10.1038/nature08867>.
- Belfiori B, Riccioni C, Paolocci F, Rubini A. 2016. Characterization of the reproductive mode and life cycle of the whitish truffle *T. borchii*. *Mycorrhiza* 26:515–527. <https://doi.org/10.1007/s00572-016-0689-0>.
- Iotti M, Barbieri E, Stocchi V, Zambonelli A. 2005. Morphological and molecular characterisation of mycelia of ectomycorrhizal fungi in pure culture. *Fungal Divers* 19:51–68.
- Kohler A, Kuo A, Nagy LG, Morin E, Barry KW, Buscot F, Canbäck B, Choi C, Cichocki N, Clum A, Colpaert J, Copeland A, Costa MD, Doré J, Floudas D, Gay G, Giralanda M, Henrissat B, Herrmann S, Hess J, Högberg N, Johansson T, Khouja H-R, LaButti K, Lahrman U, Levasseur A, Lindquist EA, Lipzen A, Marmeisse R, Martino E, Murat C, Ngan CY, Nehls U, Plett JM, Pringle A, Ohm RA, Perotto S, Peter M, Riley R, Rineau F, Ruytinx J, Salamov A, Shah F, Sun H, Tarkka M, Tritt A, Veneault-Fourrey C, Zuccaro A, Tunlid A, Grigoriev IV, Hibbett DS, Martin F. 2015. Convergent losses of decay mechanisms and rapid turnover of symbiosis genes in mycorrhizal mutualists. *Nat Genet* 47:410–415. <https://doi.org/10.1038/ng.3223>.
- Chin C-S, Peluso P, Sedlazeck FJ, Nattestad M, Concepcion GT, Clum A, Dunn C, O’Malley R, Figueroa-Balderas R, Morales-Cruz A, Cramer GR, Delledonne M, Luo C, Ecker JR, Cantu D, Rank DR, Schatz MC. 2016. Phased diploid genome assembly with single-molecule real-time sequencing. *Nat Methods* 13:1050–1054. <https://doi.org/10.1038/nmeth.4035>.
- Lam K-K, LaButti K, Khalak A, Tse D. 2015. FinisherSC: a repeat-aware tool for upgrading *de novo* assembly using long reads. *Bioinformatics* 31: 3207–3209. <https://doi.org/10.1093/bioinformatics/btv280>.
- Chin C-S, Alexander DH, Marks P, Klammer AA, Drake J, Heiner C, Clum A, Copeland A, Huddleston J, Eichler EE, Turner SW, Korlach J. 2013. Nonhybrid, finished microbial genome assemblies from long-read SMRT sequencing data. *Nat Methods* 10:563–569. <https://doi.org/10.1038/nmeth.2474>.
- Martin J, Bruno VM, Fang Z, Meng X, Blow M, Zhang T, Sherlock G, Snyder M, Wang Z. 2010. Rnnotator: an automated *de novo* transcriptome assembly pipeline from stranded RNA-seq reads. *BMC Genomics* 11:663. <https://doi.org/10.1186/1471-2164-11-663>.
- Grigoriev IV, Nikitin R, Haridas S, Kuo A, Ohm R, Otilar R, Riley R, Salamov A, Zhao X, Korzeniewski F, Smirnova T, Nordberg H, Dubchak I, Shabalov I. 2014. MycoCosm portal: gearing up for 1000 fungal genomes. *Nucleic Acids Res* 42:D699–D704. <https://doi.org/10.1093/nar/gkt1183>.
- Kuo A, Bushnell B, Grigoriev IV. 2014. Fungal genomics: sequencing and annotation, p 1–52. In Martin F (ed), *Fungi. Advances in botanical research*. Elsevier Academic Press, Cambridge, United Kingdom.
- Murat C, Martin F. 2016. Truffle genomics: investigating an early diverging lineage of *Pezizomycotina*, p 137–149. In Zambonelli A, Iotti M, Murat C (ed), True truffle (*Tuber* spp.) in the world. Springer International Publishing, Cham, Switzerland.
- Traeger S, Altegoer F, Freitag M, Gabaldon T, Kempken F, Kumar A, Marcet-Houben M, Pöggeler S, Stajich JE, Nowrousian M. 2013. The genome and development-dependent transcriptomes of *Pyronema confluens*: a window into fungal evolution. *PLoS Genet* 9:e1003820. <https://doi.org/10.1371/journal.pgen.1003820>.
- Debuchy R, Berteaux-Lecellier V, Silar P. 2010. Mating systems and sexual morphogenesis in ascomycetes, p 501–535. In Borkovich KA, Ebbole DJ (ed), *Cellular and molecular biology of filamentous fungi*. ASM Press, Washington, DC.