



Complete Genome Sequence of *Rice hoja blanca tenuivirus* Isolated from a Susceptible Rice Cultivar in Colombia

Jenyfer Jimenez,^{a,b} Monica Carvajal-Yepes,^a Ana Maria Leiva,^a Maribel Cruz,^c Luz Elena Romero,^a Carmen A. Bolaños,^a Ivan Lozano,^a Wilmer J. Cuellar^a

^aVirology Laboratory, Agrobiodiversity Research Area (AgBio), International Center for Tropical Agriculture (CIAT), Cali, Colombia

^bFacultad de Ciencias Agropecuarias, Universidad Nacional de Colombia (UNAL), Palmira, Colombia

^cLatin American Fund for Irrigated Rice (FLAR), Cali, Colombia

ABSTRACT We describe here the complete genome of *Rice hoja blanca tenuivirus*. The sequenced isolate was obtained by insect vector transmission from a symptomatic rice sample grown in Colombia. Sequence data from the four RNA components were obtained by deep sequencing (Illumina), and infections were confirmed by enzyme-linked immunosorbent assay and Sanger sequencing.

Rice hoja blanca tenuivirus (RHBV) is a multipartite single-stranded, negative-sense RNA virus belonging to the genus *Tenuivirus*. It causes hoja blanca disease (HBD) in rice (*Oryza sativa* L.), which has been reported as affecting rice yield in Colombia since 1935. The virus was first described in 1983 (1) and partially characterized between 1992 and 1993 (2, 3). Since then, HBD has been reported in countries of major rice production in Latin America, the Caribbean, and the southern United States. The vector of RHBV is *Tagosodes orizicolus* Müir (Hemiptera: Delphacidae), which can also be infected by the virus (4). Infected rice plants show symptoms that include chlorotic striping and mottling on the leaves, plant dwarfing, panicle sterility, premature wilting, and necrosis, causing serious yield losses (25 to 75%; [4]). Until now, a complete genome sequence of RHBV has not been available.

To obtain the complete genome of RHBV, the rice cultivar Bluebonnet 50 was infected with an isolate from Tolima (central Colombia) using *T. orizicolus* as a vector. Total RNA extraction was carried out, using TRIzol (5), from 50 mg of dried leaf tissue. cDNA libraries were prepared from small interfering RNAs (siRNAs) as previously described (6), and deep sequencing was performed using the Illumina (San Diego, CA, USA) platform, according to the manufacturer's instructions. Removal of the 3' end adapters resulted in 1,676,078 reads between 21 and 24 nucleotides (nt) in length that generated 162 contigs; the conditions for contig assembly were a minimal overlapping length *k*-mer of 15 and minimum contig length of 50 nt. Genome assembly was achieved using Velvet version 1.2.03 for *de novo* sequence assembly (7) and MAQ version 0.7.1 software (<http://maq.sourceforge.net>). Compared to the GenBank nonredundant database using BLASTx and BLASTn searches, 33.33% of the total number of contigs showed significant similarity with different viral sequences reported for RHBV. RHBV encodes a typical tenuivirus genome arrangement compared to *Rice stripe tenuivirus* (RSV), the type species of the group (8). It is composed of four RNA components, where RNA2 to RNA4 encode two open reading frames (ORFs), the second of which is encoded by the complementary strand. All RNA components present a conserved complementary 5' and 3' terminal nucleotide sequence (5' ACACAAAGTC 3'). RHBV RNA1 (8,999 bp) encodes the RNA-dependent RNA polymerase (RdRp; 73%). RNA2 (3,620 bp) has an ambisense translation strategy encoding a nonstructural (NS2; 60%) protein in the 5' proximal end and the membrane glycoprotein (68%) at the 3'

Received 30 November 2017 Accepted 17 January 2018 Published 15 February 2018

Citation Jimenez J, Carvajal-Yepes M, Leiva AM, Cruz M, Romero LE, Bolaños CA, Lozano I, Cuellar WJ. 2018. Complete genome sequence of *Rice hoja blanca tenuivirus* isolated from a susceptible rice cultivar in Colombia. *Genome Announc* 6:e01490-17. <https://doi.org/10.1128/genomeA.01490-17>.

Copyright © 2018 Jimenez et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Wilmer J. Cuellar, w.cuellar@cgiar.org.

J.J. and M.C.-Y. contributed equally to this work.

proximal ORF. RNA3 (2,299 bp) also encodes two ORFs, the NS3 protein (67%) in the 5' proximal ORF and the nucleocapsid protein (65%) in the 3' proximal ORF. RNA4 (1,998 bp) encodes the major nonstructural protein (NCP; 74%) at the 5' proximal end and the NS4 protein (75%) at the 3' proximal end. The percentage values indicate the amino acid sequence similarity to RSV for each RHBV protein. Specific primers and Sanger sequencing (Macrogen, South Korea) were used to confirm the assembled sequences and filling in of gapped regions.

Accession number(s). The complete genome sequences reported here have been deposited at GenBank under the accession no. [MG566074](#) (RNA1), [MG566075](#) (RNA2), [MG566076](#) (RNA3), and [MG566077](#) (RNA4).

ACKNOWLEDGMENTS

We thank Joe Tohme, Edgar Torres, Fernando Correa, and Mathias Lorieux for supporting the study.

This work was supported by the Federal Ministry for Economic Cooperation and Development (GIZ-BMZ) of Germany and the CIAT Rice Program.

REFERENCES

- Morales FJ, Niessen AI. 1983. Association of spiral filamentous virus-like particles with rice hoja blanca. *Phytopathology* 73:971–974. <https://doi.org/10.1094/Phyto-73-971>.
- Ramirez BC, Macaya G, Calvert LA, Haenni AL. 1992. Rice hoja blanca virus genome characterization and expression in vitro. *J Gen Virol* 73: 1457–1464. <https://doi.org/10.1099/0022-1317-73-6-1457>.
- Ramirez BC, Lozano I, Constantino LM, Haenni AL, Calvert LA. 1993. Complete nucleotide sequence and coding strategy of rice hoja blanca virus RNA4. *J Gen Virol* 74:2463–2468. <https://doi.org/10.1099/0022-1317-74-11-2463>.
- Morales FJ, Jennings PR. 2010. Rice hoja blanca: a complex plant–virus–vector pathosystem. *Plant Sci Rev* 5:043.
- Chomczynski P, Sacchi N. 1987. Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal Biochem* 162:156–159. [https://doi.org/10.1016/0003-2697\(87\)90021-2](https://doi.org/10.1016/0003-2697(87)90021-2).
- Chen YR, Zheng Y, Liu B, Zhong S, Giovannoni J, Fei Z. 2012. A cost-effective method for Illumina small RNA-Seq library preparation using T4 RNA ligase 1 adenylated adapters. *Plant Methods* 8:41. <https://doi.org/10.1186/1746-4811-8-41>.
- Zerbino DR, Birney E. 2008. Velvet: algorithms for de novo short read assembly using de Bruijn graphs. *Genome Res* 18:821–829. <https://doi.org/10.1101/gr.074492.107>.
- Lefkowitz EJ, Davison AJ, Sabanadzovic S, Siddell SG, Simmonds P (ed). 2017. Classification and nomenclature of viruses: online (10th) report of the International Committee on Taxonomy of Viruses. https://talk.ictvonline.org/ictv-reports/ictv_online_report/.