



# Draft Genome Sequence of the Type Strain *Streptomyces armeniacus* ATCC 15676

Puneet Labana,<sup>a</sup> Jessica T. Gosse,<sup>a</sup> Christopher N. Boddy<sup>a</sup>

<sup>a</sup>Department of Chemistry and Biomolecular Sciences, University of Ottawa, Ottawa, ON, Canada

**ABSTRACT** The *Streptomyces* genus represents a prolific and significant source for secondary metabolite discovery. Here, we report a *de novo* assembly and draft genome sequence for the type strain *Streptomyces armeniacus* ATCC 15676.

*Streptomyces* spp. are known producers of a diverse array of bioactive natural products, including antibiotics, antifungals, and anticancer agents (1). As such, considerable effort has been spent examining and sequencing environmental isolates to characterize their ability to produce these important molecules (2, 3). However, to complement this effort, it is essential to also adequately characterize *Streptomyces* type strains, as they establish the points of reference that other *Streptomyces* strains are compared with (4). Here, we sequenced the complete genome of *Streptomyces armeniacus* ATCC 15676 (NBRC 12555, BCRC 16847, RIA 807, 26A-32, KCC A-0070, NCIB 10179) type strain, isolated from soil of Armenian origin (5, 6), and assembled a high-quality draft genome sequence.

The *Streptomyces* strain was revived per ATCC guidelines and cultivated in ISP-2 medium at 26°C for 1 week (7, 8). The genomic DNA (gDNA) was then extracted using a Wizard genomic DNA purification kit (Promega). Purified gDNA was sent to Dalhousie University (Halifax, NS, Canada) for sequencing at the Centre for Comparative Genomics and Evolutionary Bioinformatics (CGEB). Sequencing results were obtained using an Illumina MiSeq DNA sequencer. The library was prepared with a Nextera DNA sample preparation kit (Illumina), following the manufacturer's instructions, and sequenced using a MiSeq 600-cycle v3 reagent kit (Illumina).

*De novo* assembly of paired-end files was performed using a SPAdes protocol in the PATRIC genome assembly service (version 3.5.18) (9). Quality control of reads was set to a minimum contig coverage of 5× and a minimum contig length of 300 bp. The resulting draft genome sequence consists of 12 contigs with a total genome size of 8,073,679 bp and a G+C content of 72.4%. The average read length is 3,354,904 bp with an  $N_{50}$  value of 2,357,927 and coverage of 65×.

Subsequently, an automated annotation was performed with Rapid Annotations using Subsystem Technology (RAST) version 2.0 (10). A total of 7,080 coding sequences and 68 RNAs were predicted. Biosynthetic gene clusters (BGCs) were identified through antiSMASH version 4.2.0 (11). In total, 36 BGCs related to multiple different secondary metabolites were predicted. Mixed pathway products, along with the type 1 polyketide synthases (PKSs) and nonribosomal peptide synthetases (NRPSs) were most abundant, with 10, 7, and 6 BGCs, respectively. Following were 4 BGCs encoding ribosomal natural products and 4 BGCs encoding terpene pathways. A single prediction each for a siderophore, type 2 PKS, and type 3 PKS BGC were also identified. It is noteworthy that while this type strain is not documented to produce any secondary metabolites, it clearly possesses significant genetic potential, which further validates the utility of type strain sequencing.

**Data availability.** This whole-genome sequence has been deposited at DDBJ/ENA/GenBank under the accession number [CP031320](https://doi.org/10.1128/MRA.01107-18). The version described in this paper is the first version.

Received 8 August 2018 Accepted 23 August 2018 Published 20 September 2018

**Citation** Labana P, Gosse JT, Boddy CN. 2018. Draft genome sequence of the type strain *Streptomyces armeniacus* ATCC 15676. *Microbiol Resour Announc* 7:e01107-18. <https://doi.org/10.1128/MRA.01107-18>.

**Editor** John J. Dennehy, Queens College

**Copyright** © 2018 Labana et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Christopher N. Boddy, [cboddy@uottawa.ca](mailto:cboddy@uottawa.ca).

## REFERENCES

- de Lima Procópio RE, da Silva IR, Martins MK, de Azevedo JL, de Araújo JM. 2012. Antibiotics produced by *Streptomyces*. *Brazilian J Infect Dis* 16:466–471. <https://doi.org/10.1016/j.bjid.2012.08.014>.
- Hill P, Heberlig GW, Boddy CN. 2017. Sampling terrestrial environments for bacterial polyketides. *Molecules* 22:707. <https://doi.org/10.3390/molecules22050707>.
- Gosse JT, Hill P, Dowd SE, Boddy CN. 2015. Draft genome sequence of *Streptomyces* sp. strain PBH53, isolated from an urban environment. *Genome Announc* 3:e00859-15. <https://doi.org/10.1128/genomeA.00859-15>.
- Wu L, McCluskey K, Desmeth P, Liu S, Hideaki S, Yin Y, Moriya O, Itoh T, Kim C, Lee J-S, Zhou Y, Kawasaki H, Hazbón MH, Robert V, Boekhout T, Lima N, Evtushenko L, Boundy-Mills K, Bunk B, Moore ERB, Eurwilaichitr L, Ingsriswang S, Shah H, Yao S, Jin T, Huang J, Shi W, Sun Q, Fan G, Li W, Li X, Kurtböke I, Ma J. 2018. The global catalogue of microorganisms 10K type strain sequencing project: closing the genomic gaps for the validly published prokaryotic and fungi species. *GigaScience* 7:1–4. <https://doi.org/10.1093/gigascience/giy026>.
- Kalakoutskii LV, Kusnetsov VD. 1964. A new species of the Actinoplanes—*A. armeniacus* and some peculiarities of its mode of spore formation. *Mikrobiologiya* 33:613–621.
- Wellington EMH, Williams ST. 1981. Transfer of *Actinoplanes armeniacus* Kalakoutskii and Kusnetsov to *Streptomyces*: *Streptomyces armeniacus* (Kalakoutskii and Kusnetsov) comb. nov. *International J Systematic Bacteriology* 31:77–81. <https://doi.org/10.1099/00207713-31-1-77>.
- American Type Culture Collection. 2015. ATCC bacterial culture guide. American Type Culture Collection, Manassas, VA. [https://www.atcc.org/~media/PDFs/Culture%20Guides/ATCC\\_Bacterial\\_Culture\\_Guide.ashx](https://www.atcc.org/~media/PDFs/Culture%20Guides/ATCC_Bacterial_Culture_Guide.ashx).
- Zimbro MJ, Power DA, Miller SM, Wilson GE, Johnson JA. 2009. ISP medium 1, ISP medium 2, ISP medium 4, p 2–3. *In* Difco & BBL manual: manual of microbiological culture media. BD Diagnostics—Diagnostic Systems, Sparks, MD. [http://www.bd.com/europe/regulatory/assets/ifu/difco\\_bbl/277210.pdf](http://www.bd.com/europe/regulatory/assets/ifu/difco_bbl/277210.pdf).
- Wattam AR, Davis JJ, Assaf R, Boisvert S, Brettin T, Bun C, Conrad N, Dietrich EM, Disz T, Gabbard JL, Gerdes S, Henry CS, Kenyon RW, Machi D, Mao C, Nordberg EK, Olsen GJ, Murphy-Olson DE, Olson R, Overbeek R, Parrello B, Pusch GD, Shukla M, Vonstein V, Warren A, Xia F, Yoo H, Stevens RL. 2017. Improvements to PATRIC, the all-bacterial bioinformatics database and analysis resource center. *Nucleic Acids Res* 45:D535–D542. <https://doi.org/10.1093/nar/gkw1017>.
- Overbeek R, Olson R, Pusch GD, Olsen GJ, Davis JJ, Disz T, Edwards RA, Gerdes S, Parrello B, Shukla M, Vonstein V, Wattam AR, Xia F, Stevens R. 2014. The SEED and the Rapid Annotation of microbial genomes using Subsystems Technology (RAST). *Nucleic Acids Res* 42:D206–D214. <https://doi.org/10.1093/nar/gkt1226>.
- Blin K, Wolf T, Chevrette MG, Lu X, Schwalen CJ, Kautsar SA, Suarez Duran HG, de Los Santos ELC, Kim HU, Nave M, Dickschat JS, Mitchell DA, Shelest E, Breitling R, Takano E, Lee SY, Weber T, Medema MH. 2017. antiSMASH 4.0—improvements in chemistry prediction and gene cluster boundary identification. *Nucleic Acids Res* 45:W36–W41. <https://doi.org/10.1093/nar/gkx319>.