



Complete Genomic Sequence of the Potyvirus *Mashua Virus Y*, Obtained from a 33-Year-Old Mashua (*Tropaeolum tuberosum*) Sample

Ian P. Adams,^a Adrian Fox,^a Neil Boonham,^{a,b} Roger A. C. Jones^{c,d}

^aFera, Sand Hutton, York, United Kingdom

^bSchool of Natural and Environmental Sciences, Newcastle University, Newcastle upon Tyne, United Kingdom

^cInstitute of Agriculture, Faculty of Science, University of Western Australia, Crawley, Western Australia, Australia

^dDepartment of Primary Industries and Regional Development, South Perth, Western Australia, Australia

ABSTRACT We present the complete genomic sequence of a new potyvirus we tentatively call *Mashua virus Y* (MasVY), first isolated in 1984 from a plant of the Andean tuber crop mashua (*Tropaeolum tuberosum*, family *Tropaeolaceae*). There was a 70% nucleotide identity between MasVY and a genomic sequence of *Verbena virus Y*.

In 1984, a tuber sample of mashua (*Tropaeolum tuberosum*, family *Tropaeolaceae*) (1) was obtained from an Andean root and tuber species collection at the Department of Applied Biology at the University of Cambridge in England. After the tuber was planted, a virus was transmitted from the plant grown to *Nicotiana clelandii* plants. Both original mashua and inoculated *N. clelandii* foliage developed mosaics. Electron microscopy of infected plant sap from *N. clelandii* revealed approximately 750-nm-long flexuous filamentous potyvirus particles. In 1985, infected *N. clelandii* leaf samples were freeze-dried in glass vials, which now are kept in the “Fera plant virus collection.” Mashua plants showing mosaic symptoms were first recorded in 1977 in Bolivia (2). In 1990, two potyviruses named Tropaeolum 1 and 2 potyviruses were reported to cause leaf chlorosis in plants grown from mashua tubers imported to England from Peru (3). In 1998, a potyvirus infecting mashua imported to the United States from Ecuador was reported, and it was named Tropaeolum mosaic potyvirus (TropMV) (4). Whether TropMV resembles either of the Tropaeolum 1 and 2 potyviruses is unclear. Subsequently, TropMV from Peruvian mashua was shown to reduce tuber yield (5). None of these mashua potyviruses were ever sequenced.

In 2017, using an RNAeasy kit (Qiagen, UK) including optional DNase treatment, total RNA was extracted from infected leaf material from *N. clelandii* freeze-dried in 1985. An indexed plant ribosome-subtracted sequencing library was produced using the ScriptSeq complete plant leaf kit (Illumina, USA) following the manufacturer’s instructions. The indexed library was sequenced on a MiSeq instrument (Illumina) with a 600-cycle v3 kit. The resulting 748,695 paired reads were 3’ trimmed to a quality score of 20 with Sickle in paired-end mode (6) and assembled with Trinity v2 with the maximum memory allocation set to 99 GB of RAM, and the process allocated 64 central processing units (CPUs) (7), and the resulting contigs were compared to the GenBank nonredundant (nr) and nucleotide databases with BLAST+ (8). Reads of viral origin were extracted with the extract reads function in MEGAN (9). A 9,769-nucleotide (nt) contig was assembled by comparison with other genomes and constituted a complete potyvirus genome with a complete potyvirus coding region and typical 5’ and 3’ untranslated regions (UTRs). The sequence contains a complete potyvirus coding region and typical 5’ and 3’ UTRs. There was a 70% nt identity between the new

Received 14 August 2018 Accepted 24 September 2018 Published 18 October 2018

Citation Adams IP, Fox A, Boonham N, Jones RAC. 2018. Complete genomic sequence of the potyvirus *Mashua virus Y*, obtained from a 33-year-old mashua (*Tropaeolum tuberosum*) sample. *Microbiol Resour Announc* 7:e01064-18. <https://doi.org/10.1128/MRA.01064-18>.

Editor Catherine Putonti, Loyola University Chicago

Copyright © 2018 Adams et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Roger A. C. Jones, roger.jones@uwa.edu.au.

genome and the nearest genomic sequence, *Verbena virus Y* (VVY) (GenBank accession number [NC_010735](#)), from the ornamental plant *Verbena* × *hybrid* (family *Verbenaceae*) reported from the United States (10). This <76% nt identity falls outside the species discrimination limit for *Potyviridae* (11). Establishing the relationship between the new potyvirus found in this study, which we tentatively name *Mashua virus Y* (MasVY), and the three mashua potyviruses named previously (3–5) awaits future studies. VVY belongs to an American subclade also containing *Pepper mottle virus* and *Potato virus Y* (10), so MasVY probably belongs with them. In 2018, to assess the likelihood of virus dissemination via Internet trading in unregulated plant species, RNA extracted from a mashua tuber obtained from a European country was sequenced as described above. The 1,335-nt potyvirus sequence recovered had 91% (nucleotide) and 96% (amino acid) coat protein identity with MasVY, which suggests MasVY's presence in Internet-traded mashua tubers.

Data availability. The sequences described here were deposited in GenBank under accession numbers [MH680824](#) (genome) and [MH680823](#) (partial sequence). Raw data were deposited in the SRA under BioSample number [SAMN10081141](#), which is part of BioProject [PRJNA491634](#).

ACKNOWLEDGMENTS

This work was supported by the UK Government's Department of Environment, Food and Rural Affairs (Defra) under the Defra-Fera long-term services agreement. Sequencing work was carried out with funding from Defra.

REFERENCES

- Hawkes JG, Harris DR, Hillman GC. 1989. The domestication of roots and tubers in the American tropics, p 481–503. *In* Harris DR, Hillman GC (ed), *Foraging and farming: the evolution of plant exploitation*. Unwin Hyman, Boston, MA.
- Delhey R, Monasterios T. 1977. A mosaic disease of isañu (*Tropaeolum tuberosum*) from Bolivia. *J Plant Dis Protect* 84:224–231.
- Brunt AA, Crabtree K, Gibbs AJ (ed). 1999. *Viruses of tropical plants: descriptions and lists from the VIDE database*. CAB International, Wallingford, United Kingdom. <https://www.cabdirect.org/cabdirect/abstract/19912304303#>.
- Soria SL, Vega R, Damsteegt VD, McDaniel LL, Kitto SL, Evans TA. 1998. Occurrence and partial characterization of a new mechanically transmissible virus in mashua from the Ecuadorian highlands. *Plant Dis* 82:69–73. <https://doi.org/10.1094/PDIS.1998.82.1.69>.
- Guimarães RL, Flores HE. 2005. *Tropaeolum* mosaic potyvirus (TropMV) reduces yield of Andean mashua (*Tropaeolum tuberosum*) accessions. *HortScience* 40:1405–1407.
- Najoshi. 2011. Sickle—a windowed adaptive trimming tool for FASTQ files using quality. <https://github.com/najoshi/sickle>.
- Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, Adiconis X, Fan L, Raychowdhury R, Zeng Q, Chen Z, Muceli E, Hacohen N, Gnirke A, Rhind N, di Palma F, Birren BW, Nusbaum C, Lindblad-Toh K, Friedman N, Regev A. 2011. Full-length transcriptome assembly from RNA-Seq data without a reference genome. *Nat Biotechnol* 29:644–652. <https://doi.org/10.1038/nbt.1883>.
- Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009. BLAST+: architecture and applications. *BMC Bioinformatics* 10:421. <https://doi.org/10.1186/1471-2105-10-421>.
- Huson DH, Beier S, Flade I, Górska A, El-Hadidi M, Mitra S, Ruscheweyh H-J, Tappu R. 2016. MEGAN community edition—interactive exploration and analysis of large-scale microbiome sequencing data. *PLoS Comput Biol* 12:e1004957. <https://doi.org/10.1371/journal.pcbi.1004957>.
- Kraus J, Cleveland S, Putnam ML, Keller KE, Martin RR, Tzanetakis IE. 2010. A new *Potyvirus* sp. infects verbena exhibiting leaf mottling symptoms. *Plant Dis* 94:1132–1136. <https://doi.org/10.1094/PDIS-94-9-1132>.
- Adams MJ, Antoniw JF, Fauquet CM. 2005. Molecular criteria for genus and species discrimination within the family *Potyviridae*. *Arch Virol* 150:459–479. <https://doi.org/10.1007/s00705-004-0440-6>.