



Draft Genome Sequence of *Acidobacteria* Group 1 *Acidipila* sp. Strain EB88, Isolated from Forest Soil

Luiz A. Domeignoz-Horta,^a Kristen M. DeAngelis,^a Grace Pold^b

^aDepartment of Microbiology, University of Massachusetts, Amherst, Massachusetts, USA

^bGraduate Program in Organismic and Evolutionary Biology, University of Massachusetts, Amherst, Massachusetts, USA

ABSTRACT Here, we present the genome sequence of a member of the group I *Acidobacteria*, *Acidipila* sp. strain EB88, which was isolated from temperate forest soil. Like many other members of its class, its genome contains evidence of the potential to utilize a broad range of sugars.

Group I *Acidobacteria* are abundant members of acidic-soil microbial communities (1), typically characterized by slow growth, heterotrophy, and the production of copious extracellular polysaccharides (2). Within this group, members of the genus *Acidipila* are characterized as acidophilic heterotrophs that use sugars as favored growth substrates (3, 4).

Acidipila sp. strain EB88 was isolated from the <0.8- μ m size fraction of a deciduous forest mineral soil slurry plated onto VL45 plus glucose-yeast extract (GYE) medium (5). It was selected for sequencing based on the paucity of cultivated group I *Acidobacteria*, its peculiar bright-red waxy colonies on solid medium, and its undetectable growth in liquid medium unless a solid substrate, such as sand, is added.

DNA was prepared for sequencing by repeatedly freeze-thawing 9-day-old biomass scraped from pH 5 R2A medium and extracted using the Qiagen genomic DNA protocol. Sequencing at UMass Worcester using the PacBio RS II sequencer resulted in 476,266 filtered reads, with a mean length of 3,208 bp. The genome was assembled using sprai version 0.9.9.23 (<http://zombie.cb.k.u-tokyo.ac.jp/sprai/index.html>) and Canu version 1.5 (6). Default parameters were used for all tools unless otherwise specified. This yielded a 4.48-Mbp genome spanning 5 contigs, estimated as 100% complete and 0.86% contaminated using CheckM version 1.0.8 (7) in KBASE (8). Its GC content is 62.6%. Gene calling and annotation using the U.S. Department of Energy Joint Genome Institute's annotation pipeline (9) identified 48 tRNA genes, 2 rRNA operons, and 3,593 protein-coding genes. TCDB (10) and TransAAP (11) annotated 392 of these genes as transporters.

The genome of EB88 is most closely related to the publicly available *Acidipila rosea* genome, with which it shares 94.5% 16S rRNA identity and 72.1% average nucleotide identity, according to IMG, and it has 60.9% mean amino acid identity using the average amino acid identity tool (<http://enve-omics.ce.gatech.edu/aai/>). The genome of EB88 is deficient in genes for organic acid uptake but contains genes for amino acid, ammonium, and nitrate uptake. It is capable of growth under both high- and low-oxygen conditions, and its genome contains five high-affinity *cbb*₃ terminal oxidases and one low-affinity group A heme-copper oxygen oxidase (12). As is typical for the group I soil *Acidobacteria*, the genome of EB88 is rich in glycolytic enzymes and was found to contain 85 glycoside hydrolases in 48 families using HMMer searches in dbCAN (13). dbCAN also identified genes for capsule biosynthesis and export, including cellulose (14). However, rather than forming a copious hydrophilic capsule as is common for the group (2), EB88 has a waxy composition when grown on solid agar medium. It shares

Citation Domeignoz-Horta LA, DeAngelis KM, Pold G. 2019. Draft genome sequence of *Acidobacteria* group 1 *Acidipila* sp. strain EB88, isolated from forest soil. *Microbiol Resour Announc* 8:e01464-18. <https://doi.org/10.1128/MRA.01464-18>.

Editor David A. Baltus, University of Arizona

Copyright © 2019 Domeignoz-Horta et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Kristen M. DeAngelis, deangelis@microbio.umass.edu.

Received 1 November 2018

Accepted 19 November 2018

Published 3 January 2019

most of its lipid and fatty acid biosynthesis genes with other acidobacteria, so we posit that the hydrophobic nature of its colonies is driven by deficiencies in fatty acid degradation genes (EC 1.3.8.1, 2.3.1.9, and 2.3.1.16).

Together, these results indicate that the genome of *Acidipila* sp. EB88 is typical of many group I *Acidobacteria* but possibly differs in genes involved in extracellular associations. Its niche in forest soils is expected to include the use of sugars and amino acids under microaerobic to aerobic conditions.

Data availability. The complete genome sequence is available in GenBank under accession number [QWEV00000000](https://www.ncbi.nlm.nih.gov/nuccore/QWEV00000000). The version described in this paper is the first version, QWEV01000000.

ACKNOWLEDGMENTS

Funding for this work came from Department of Energy contract DE-SC0016590 to K.M.D. G.P. is funded by an American Fellowship from AAUW.

REFERENCES

- Jones RT, Robeson MS, Lauber CL, Hamady M, Knight R, Fierer N. 2009. A comprehensive survey of soil acidobacterial diversity using pyrosequencing and clone library analyses. *ISME J* 3:442. <https://doi.org/10.1038/ismej.2008.127>.
- Eichorst SA, Trojan D, Roux S, Herbold C, Rattei T, Wobken D. 2018. Genomic insights into the *Acidobacteria* reveal strategies for their success in terrestrial environments. *Environ Microbiol* 20:1041–1063. <https://doi.org/10.1111/1462-2920.14043>.
- Jiang YW, Wang J, Chen MH, Lv YY, Qiu LH. 2016. *Acidipila dinghuensis* sp. nov., an acidobacterium isolated from forest soil. *Int J Syst Evol Microbiol* 66:76–83. <https://doi.org/10.1099/ijsem.0.000676>.
- Okamura K, Kawai A, Yamada T, Hiraishi A. 2011. *Acidipila rosea* gen. nov., sp. nov., an acidophilic chemoorganotrophic bacterium belonging to the phylum *Acidobacteria*. *FEMS Microbiol Lett* 317:138–142. <https://doi.org/10.1111/j.1574-6968.2011.02224.x>.
- Lladó S, Žifčáková L, Větrovský T, Eichlerová I, Baldrian P. 2016. Functional screening of abundant bacteria from acidic forest soil indicates the metabolic potential of *Acidobacteria* subdivision 1 for polysaccharide decomposition. *Biol Fertil Soils* 52:251–260. <https://doi.org/10.1007/s00374-015-1072-6>.
- Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM. 2017. Canu: scalable and accurate long-read assembly via adaptive k-mer weighting and repeat separation. *Genome Res* 27:722–736. <https://doi.org/10.1101/gr.215087.116>.
- Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. 2015. CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res* 14:1043–1055.
- Arkin AP, Cottingham RW, Henry CS, Harris NL, Stevens RL, Maslov S, Dehal P, Ware D, Perez F, Canon S, Sneddon MW, Henderson ML, Riehl WJ, Murphy-Olson D, Chan SY, Kamimura RT, Kumari S, Drake MM, Brettin TS, Glass EM, Chivian D, Gunter D, Weston DJ, Allen BH, Baumohl J, Best AA, Bowen B, Brenner SE, Bun CC, Chandonia JM, Chia JM, Colasanti R, Conrad N, Davis JJ, Davison BH, DeJongh M, Dvoid S, Dietrich E, Dubchak I, Edirisinghe JN, Fang G, Faria JP, Frybarger PM, Gerlach W, Gerstein M, Greiner A, Gurtowski J, Haun HL, He F, Jain R, et al. 2018. KBase: the United States Department of Energy Systems Biology Knowledgebase. *Nat Biotechnol* 36:566. <https://doi.org/10.1038/nbt.4163>.
- Huntemann M, Ivanova NN, Mavromatis K, Tripp HJ, Paez-Espino D, Palaniappan K, Szeto E, Pillay M, Chen I-MA, Pati A, Nielsen T, Markowitz VM, Kyrpides NC. 2015. The standard operating procedure of the DOE-JGI Microbial Genome Annotation Pipeline (MGAP v. 4). *Stand Genomic Sci* 10:86. <https://doi.org/10.1186/s40793-015-0077-y>.
- Saier MH, Jr, Reddy VS, Tsu BV, Ahmed MS, Li C, Moreno-Hagelsieb G. 2016. The Transporter Classification Database (TCDB): recent advances. *Nucleic Acids Res* 44:D372–D379. <https://doi.org/10.1093/nar/gkv1103>.
- Elbourne LD, Tetu SG, Hassan KA, Paulsen IT. 2017. TransportDB 2.0: a database for exploring membrane transporters in sequenced genomes from all domains of life. *Nucleic Acids Res* 45:D320–D324. <https://doi.org/10.1093/nar/gkw1068>.
- Morris RL, Schmidt TM. 2013. Shallow breathing: bacterial life at low O₂. *Nat Rev Microbiol* 11:205. <https://doi.org/10.1038/nrmicro2970>.
- Zhang H, Yohe T, Huang L, Entwistle S, Wu P, Yang Z, Busk PK, Xu Y, Yin Y. 2018. dbCAN2: a meta server for automated carbohydrate-active enzyme annotation. *Nucleic Acids Res* 46:W95–W101. <https://doi.org/10.1093/nar/gky418>.
- Schmid J, Sieber V, Rehm B. 2015. Bacterial exopolysaccharides: biosynthesis pathways and engineering strategies. *Front Microbiol* 6:496.