




Draft Genome Sequence of *Agrobacterium deltaense* Strain CNPSo 3391, Isolated from a Soybean Nodule in Mozambique

Anderson José Scherer,^{a,b} Jakeline Renata Marçon Delamuta,^{a,c} Renan Augusto Ribeiro,^c Amaral Machaculeha Chibeba,^d Stephen Kyei-Boahen,^d Marco Antonio Nogueira,^{a,c}  Mariangela Hungria^{a,b,c}

^aEmbrapa Soja, Soil Biotechnology Laboratory, Londrina, Paraná, Brazil

^bDepartment of Microbiology, Universidade Estadual de Londrina, Londrina, Paraná, Brazil

^cCNPq, SHIS QI 1 Conjunto B, Blocos A, B, C e D, Lago Sul, Brasília, Federal District, Brazil

^dInternational Institute of Tropical Agriculture (IITA), Nampula, Mozambique

ABSTRACT *Agrobacterium deltaense* strain CNPSo 3391 was isolated from a soybean nodule in Mozambique. Its genome size was estimated at 4,926,588 bp. This isolate carries several coding sequences for stress tolerance, but no identifiable nodulation or virulence genes. Possible ecological roles of bacteria isolated from legume nodules and closely related to *Agrobacterium* are discussed.

For 2 decades our group has reported the isolation of agrobacteria from root nodules of soybean (*Glycine max*) (1–3), common bean (*Phaseolus vulgaris*) (4–6), and other legumes (7–9). However, the ability to reestablish nodulation with the host legume is usually not confirmed. Another example is the *Agrobacterium deltaense* type strain YIC4121, isolated from a root nodule of *Sesbania cannabina* in China; the ability to nodulate seven legumes was not confirmed (10). Here, we report the draft genome sequence of strain CNPSo 3391 (= Moz59, = 9 J1), isolated from a plant grown in Mutequelesse, Gurué District, Zambézia Province, Mozambique, showing no symptoms of N deficiency. Preliminary genetic characterization based on the 16S rRNA and three housekeeping genes positioned the strain in a *Rhizobium* (*Agrobacterium*) clade (3).

Growth conditions for CNPSo 3391 were the same as those reported for its isolation (3), and DNA extraction and paired-end sequencing on the MiSeq platform (Illumina) were performed as described before (11), resulting in 630,975,648 bp. Shotgun sequences were assembled with the A5-MiSeq pipeline (*de novo*) v.20140604 with 128-fold genome coverage assembled in 50 contigs with an N_{50} of 177,127 bp. The genome was estimated at 4,926,588 bp, with G+C content of 59.9 mol%, confirmed with RAST v.2.0 (12) and QUAST v.2.0 (13), using default parameters. Average nucleotide identity (ANI) (ANI calculator [14]) indicated highest similarity (97.68%) with *Agrobacterium deltaense* YIC4121^T. Compared to the genomes of *A. deltaense* at the NCBI (strains NCPPB 1641, RV3, Zutra 3-1, and YIC4121^T), CNPSo 3391 is slightly smaller than YIC4121^T (5.02 Mb), but within the same G+C range of all strains.

A total of 4,765 DNA coding sequences (CDSs) were identified in RAST (12), with 49% classified in 475 subsystems; this annotation is the public version available at GenBank. Similarly to *A. deltaense* YIC4121^T, CNPSo 3391 carries no nodulation genes or *nif* and *fix* operons. However, CNPSo 3391 also carries no genes related to virulence, and we were not able to find sequences coding for *telA*, related to the speciation of some *Agrobacterium* (15). The environmental adaptability of CNPSo 3391 might be explained by genes such as 63 CDSs related to resistance to antibiotic and toxic compounds, 51 to iron acquisition and metabolism, 104 to motility and chemotaxis, and 162 to stress response.

Isolation of agrobacteria from legume root nodules seems to occur worldwide, with

Citation Scherer AJ, Delamuta JRM, Ribeiro RA, Chibeba AM, Kyei-Boahen S, Nogueira MA, Hungria M. 2019. Draft genome sequence of *Agrobacterium deltaense* strain CNPSo 3391, isolated from a soybean nodule in Mozambique. Microbiol Resour Announc 8:e01675-18. <https://doi.org/10.1128/MRA.01675-18>.

Editor David A. Baltus, University of Arizona

Copyright © 2019 Scherer et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Mariangela Hungria, mariangela.hungria@embrapa.br.

Received 11 December 2018

Accepted 5 February 2019

Published 7 March 2019

reports in Brazil (2, 4, 6–9), China (10), Ecuador (5), Mozambique (3), and Paraguay (1), among other countries. As these bacteria apparently do not carry nodulation genes, Yan et al. (10) suggested that they might be endophytes, but we cannot discard the hypothesis of a temporary acquisition of a symbiotic plasmid from another rhizobia. However, the role of agrobacteria in symbiosis deserves further investigation; interestingly, 3 decades ago *in vivo* results suggested that *Agrobacterium* spp. might produce extracellular “signals” that would supplement the ability of rhizobia to induce root nodulation in the host legume (16).

Data availability. This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under the GenBank accession number [RRZI00000000](https://doi.org/10.1007/s11274-009-0300-3), BioProject number [PRJNA507793](https://doi.org/10.1007/s11274-009-0300-3), BioSample number [SAMN10506010](https://doi.org/10.1007/s11274-009-0300-3), and organism number [RRZI00000000](https://doi.org/10.1007/s11274-009-0300-3); the version described in this paper is [RRZI01000000](https://doi.org/10.1007/s11274-009-0300-3).

ACKNOWLEDGMENTS

Funding was provided by Embrapa, INCT–Plant Growth-Promoting Microorganisms for Agricultural Sustainability and Environmental Responsibility (CNPq 465133/2014-2, Fundação Araucária–STI, CAPES), CNPq-Universal (400468/2016-6), and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior–Brasil (CAPES, Código de Financiamento 001).

REFERENCES

- Chen LS, Figueredo A, Pedrosa FO, Hungria M. 2000. Genetic characterization of soybean rhizobia in Paraguay. *Appl Environ Microbiol* 66: 5099–5103. <https://doi.org/10.1128/AEM.66.11.5099-5103.2000>.
- Hungria M, Chueire LMO, Megias M, Lamrabet Y, Probanza A, Guttierrez-Mañero FJ, Campo RJ. 2006. Genetic diversity of indigenous tropical fast-growing rhizobia isolated from soybean nodules. *Plant Soil* 288: 343–356. <https://doi.org/10.1007/s11104-006-9125-0>.
- Chibeba AM, Kyei-Boahen S, Guimarães MF, Nogueira MA, Hungria M. 2017. Isolation, characterization and selection of indigenous *Bradyrhizobium* strains with outstanding symbiotic performance to increase soybean yields in Mozambique. *Agric Ecosyst Environ* 246:291–305. <https://doi.org/10.1016/j.agee.2017.06.017>.
- Grange L, Hungria M. 2004. Genetic diversity of indigenous common bean (*Phaseolus vulgaris*) rhizobia in two Brazilian ecosystems. *Soil Biol Biochem* 36:1389–1398. <https://doi.org/10.1016/j.soilbio.2004.03.005>.
- Ribeiro RA, Ormeño-Orrillo E, Dall’Agnol RF, Graham PH, Martínez-Romero E, Hungria M. 2013. Novel *Rhizobium* lineages isolated from root nodules of common bean (*Phaseolus vulgaris* L.) in Andean and Mesoamerican areas. *Res Microbiol* 164:740–748. <https://doi.org/10.1016/j.resmic.2013.05.002>.
- Costa MR, Chibeba AM, Mercante FM, Hungria M. 2018. Polyphasic characterization of rhizobia microsymbionts of common bean [*Phaseolus vulgaris* (L.)] isolated in Mato Grosso do Sul, a hotspot of Brazilian biodiversity. *Symbiosis* 76:163–176. <https://doi.org/10.1007/s13199-018-0543-6>.
- Menna P, Hungria M, Barcellos FG, Bangel EV, Hess PN, Martínez-Romero E. 2006. Molecular phylogeny based on the 16S rRNA gene of elite rhizobial strains used in Brazilian commercial inoculants. *Syst Appl Microbiol* 29: 315–332. <https://doi.org/10.1016/j.syapm.2005.12.002>.
- Binde DR, Menna P, Bangel EV, Barcellos FG, Hungria M. 2009. rep-PCR fingerprinting and taxonomy based on the sequencing of the 16S rRNA gene of fifty-four elite commercial rhizobial strains. *Appl Microbiol Biotechnol* 83:897–908. <https://doi.org/10.1007/s00253-009-1927-6>.
- Roma Neto I, Ribeiro RA, Hungria M. 2010. Genetic diversity of elite rhizobial strains of subtropical and tropical legumes based on the 16S rRNA and *glnII* genes. *World J Microbiol Biotechnol* 22:1291–1302. <https://doi.org/10.1007/s11274-009-0300-3>.
- Yan J, Li Y, Han XZ, Chen WF, Zou WX, Xie Z, Li M. 2017. *Agrobacterium deltaense* sp. nov., an endophytic bacteria isolated from nodule of *Sesbania cannabina*. *Arch Microbiol* 199:1003–1009. <https://doi.org/10.1007/s00203-017-1367-0>.
- Megias E, Reis Junior FB, Ribeiro RA, Ollero FJ, Megias M, Hungria M. 2017. Genome sequence of *Pantoea ananatis* strain AMG 501, a plant-growth promoting bacterium isolated from rice leaves grown in paddies of southern Spain. *Genome Announc* 5:e00848-17. <https://doi.org/10.1128/genomeA.00848-17>.
- Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75. <https://doi.org/10.1186/1471-2164-9-75>.
- Gurevich A, Saveliev V, Vyahhi N, Tesler G. 2013. QUASt: quality assessment tool for genome assemblies. *Bioinformatics* 29:1072–1075. <https://doi.org/10.1093/bioinformatics/btt086>.
- Rodríguez-R LM, Konstantinidis KT. 2016. The enveomics collection: a toolbox for specialized analyses of microbial genomes and metagenomes. *PeerJ Prepr* 4:e1900v1. <https://doi.org/10.7287/peerj.preprints.1900v1>.
- Ramírez-Bahena MH, Vial L, Lassalle F, Diel B, Chapulliot D, Daubin V, Nesme X, Muller D. 2014. Single acquisition of protelomerase gave rise to speciation of a large and diverse clade within the *Agrobacterium/Rhizobium* supercluster characterized by the presence of a linear chromid. *Mol Phylogenet Evol* 73:202–207. <https://doi.org/10.1016/j.ympev.2014.01.005>.
- Caetano-Anollés G, Bauer WD. 1988. Enhanced nodule initiation on alfalfa by wild-type *Rhizobium meliloti* co-inoculated with *nod* gene mutants and other bacteria. *Planta* 174:385–395. <https://doi.org/10.1007/BF00959525>.