



Draft Genome Sequence of *Halobacillus trueperi* SS1, Isolated from Lunsu, a Saltwater Body in the Northwest Himalayas

Sonika Gupta,^a Parul Sharma,^a Kamal Dev,^{a,b} David J. Baumler,^{b,c,d} Anuradha Sourirajan^a

^aFaculty of Biotechnology, Shoolini University of Biotechnology and Management Sciences, Solan, Himachal Pradesh, India

^bDepartment of Food Science and Nutrition, University of Minnesota Twin Cities St. Paul, Minneapolis-St. Paul, Minnesota, USA

^cMicrobial and Plant Genomic Institute, University of Minnesota Twin Cities St. Paul, Minneapolis-St. Paul, Minnesota, USA

^dBiotechnology Institute, University of Minnesota Twin Cities St. Paul, Minneapolis-St. Paul, Minnesota, USA

ABSTRACT We report here the genome sequence of halophilic *Halobacillus trueperi* SS1, isolated from the Lunsu saltwater body in India. The bacteria are Gram positive and rod shaped. The genome of *H. trueperi* SS1 has 4.14 Mbp, with 4,329 coding sequences, 35 RNA genes (29 tRNAs, 2 rRNAs, and 4 noncoding RNAs), and 42.15% G+C content.

Saltwater lakes and salt mines are found across the Himalayas, yet their unique flora and fauna largely remain unexplored. *Halobacillus trueperi* SS1 (16S rRNA gene sequence submitted under GenBank accession no. [KM260166](https://www.ncbi.nlm.nih.gov/nuccore/KM260166)) was isolated from the soil sediments of Lunsu, a saltwater body located in Himachal Pradesh in the foothills of the northwestern Himalayas (1). *Halobacillus trueperi* SS1 is a strict halophile requiring at least 3.8% NaCl for growth, exhibits optimum growth at 11.6% NaCl, and tolerates up to 26.1% NaCl (1). It forms yellow-orange-pigmented colonies and produces an array of halozymes (1, 2). Despite the widespread reports of several halophiles, the mechanisms of salt tolerance have not been completely elucidated in all known halophiles. *H. trueperi* DSM10404 has been reported to accumulate glycine, betaine, and glutamate as compatible solutes for salt tolerance (3). We reported for the first time that *H. trueperi* SS1 utilizes a combination of a salt-in strategy and compatible solutes like proline, glycine betaine, and glutamate for survival under hypersaline conditions (4). To explore the salt-inducible regulons and biotechnological potential of *H. trueperi* SS1 (2, 4), we sequenced the entire genome of *H. trueperi* SS1. The *H. trueperi* SS1 bacterial strain was cultured in nutrient broth (NB) medium to an A_{600} of ~ 1.0 under optimal growth conditions (1), and the cells were harvested by centrifugation at $12,000 \times g$ for 5 min. Genomic DNA from the bacterial cell pellet was isolated as described by Sambrook et al. (5) and analyzed by agarose gel electrophoresis. The genomic DNA (200 ng) was used to prepare the paired-end sequencing library with the Illumina TruSeq Nano DNA high-throughput (HT) library preparation kit. The PCR-amplified library was analyzed in a Bioanalyzer 2100 (Agilent Technologies) using the high-sensitivity (HS) DNA chip according to the manufacturer's instructions and loaded onto the Illumina NextSeq 500 platform for cluster generation and sequencing. A total of 1,725,613 paired-end (PE) reads with 517,683,900 bp were produced from the sequencing run. A total of 1,725,613 paired-end (PE) reads with 517,683,900 bp were produced from the sequencing run. The *de novo* genome assembly of high-quality (phred score ≥ 30) PE reads and scaffolding were accomplished using SOAPdenovo version 2 (6), with a genome coverage of $130.0\times$. The assembled genome sequence of *H. trueperi* SS1 yielded 4,258,559 bp in the form of 113 scaffolds. The G+C content was found to be 42.15%. The coding sequences (CDS), RNA, and repeat regions were predicted using the National Center for Biological Information (NCBI) Prokaryotic

Citation Gupta S, Sharma P, Dev K, Baumler DJ, Sourirajan A. 2019. Draft genome sequence of *Halobacillus trueperi* SS1, isolated from Lunsu, a saltwater body in the Northwest Himalayas. *Microbiol Resour Announc* 8:e01710-18. <https://doi.org/10.1128/MRA.01710-18>.

Editor Catherine Putonti, Loyola University Chicago

Copyright © 2019 Gupta et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to David J. Baumler, dbaumler@umn.edu, or Anuradha Sourirajan, asourirajan@gmail.com.

Received 20 December 2018

Accepted 5 February 2019

Published 7 March 2019

TABLE 1 Global statistics of *Halobacillus trueperi* SS1 genome

Statistic	Annotation data
Total sequence length (bp)	4,258,559
No. of total genes	4,364
No. of total CDS	4,329
No. of coding genes	3,925
No. of coding CDS	3,925
No. of RNA genes	35
No. of rRNAs	2 (5S)
No. of complete rRNAs	1 (5S)
No. of partial rRNAs	1 (5S)
No. of tRNAs	29
No. of ncRNAs	4
No. of scaffolds	113
Scaffold N_{50} (bp)	81,490
Scaffold L_{50} (bp)	18
No. of contigs	1,533
Contig N_{50} (bp)	6,527
Contig L_{50} (bp)	203

Genome Annotation Pipeline and best-placed reference protein set of the GeneMarkS+ annotation software (version 4.6), as described previously (7, 8). A total of 4,364 CDS and 35 RNA genes (29 tRNAs, 2 rRNAs, and 4 noncoding RNAs [ncRNAs]) were predicted. One dinucleotide [(TA)₆] simple sequence repeat (SSR) was also identified using the MicroSatellite (MISA) identification tool, as described previously (9). The genome annotations of the *H. trueperi* SS1 genome provided by the NCBI are summarized in Table 1.

Data availability. This whole-genome shotgun project has been deposited in DDBJ/ENA/GenBank under the accession no. [QTLCO0000000](https://doi.org/10.1093/nar/gkw569). Raw sequence reads are available under SRA accession no. [SRR8351973](https://doi.org/10.1093/nar/gkw569).

ACKNOWLEDGMENTS

We acknowledge Prem Kumar Khosla, Vice Chancellor, and the members of yeast biology laboratory (YBL), Shoolini University, for supporting this project, and Guy Plunkett III for assistance with short-read submission to the NCBI.

This work was funded by the Shoolini University to A.S. to promote research under the Center for Omics and Biodiversity Research.

REFERENCES

- Gupta S, Sharma P, Dev K, Srivastava M, Sourirajan A. 2015. A diverse group of halophilic bacteria exist in Lunsu, a natural salt water body of Himachal Pradesh, India. *SpringerPlus* 4:274. <https://doi.org/10.1186/s40064-015-1028-1>.
- Gupta S, Sharma P, Dev K, Sourirajan A. 2016. Halophilic bacteria of Lunsu produce an array of industrially important enzymes with salt-tolerant activity. *Biochem Res Int* 2016:9237418. <https://doi.org/10.1155/2016/9237418>.
- Lu W, Zhao B, Feng D, Yang S. 2004. Cloning and characterization of the *Halobacillus trueperi betH* gene, encoding the transport system for the compatible solute glycine betaine. *FEMS Microbiol Lett* 235:393–399. <https://doi.org/10.1016/j.femsle.2004.05.015>.
- Vaidya S, Dev K, Sourirajan A. 2018. Distinct osmoadaptation strategies in the strict halophilic and halotolerant bacteria isolated from Lunsu salt water body of North West Himalayas. *Curr Microbiol* 75:888–895. <https://doi.org/10.1007/s00284-018-1462-8>.
- Sambrook J, Fritsch FF, Maniatis T. 1989. *Molecular cloning: a laboratory manual*, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Luo R, Liu B, Xie Y, Li Z, Huang W, Yuan J, He G, Chen Y, Pan Q, Liu Y, Tang J, Wu G, Zhang H, Shi Y, Liu Y, Yu C, Wang B, Lu Y, Han C, Cheung DW, Yiu S-M, Peng S, Xiaoqian Z, Liu G, Liao X, Li Y, Yang H, Wang J, Lam T-W, Wang J. 2012. SOAPdenovo2: an empirically improved memory-efficient short-read *de novo* assembler. *GigaScience* 1:18. <https://doi.org/10.1186/2047-217X-1-18>.
- Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44: 6614–6624. <https://doi.org/10.1093/nar/gkw569>.
- Haft DH, DiCuccio M, Badretdin A, Brover V, Chetvernin V, O'Neill K, Li W, Chitsaz F, Derbyshire MK, Gonzales NR, Gwadz M, Lu F, Marchler GH, Song JS, Thanki N, Yamashita RA, Zheng C, Thibaud-Nissen F, Geer LY, Marchler-Bauer A, Pruitt KD. 2018. RefSeq: an update on prokaryotic genome annotation and curation. *Nucleic Acids Res* 46:D851–D860. <https://doi.org/10.1093/nar/gkx1068>.
- Beier S, Thiel T, Münch T, Scholz U, Mascher M. 2017. MISA-web: a Web server for microsatellite prediction. *Bioinformatics* 33:2583–2585. <https://doi.org/10.1093/bioinformatics/btx198>.