



# Draft Genome Sequences of 38 *Serratia marcescens* Isolates Associated with Acroporid Serratosiis

Nicole C. Elledge,<sup>a</sup> Ron I. Eytan,<sup>b</sup> Lee J. Pinnell,<sup>a</sup> Reaveilyn Pray,<sup>a</sup> Jessica L. Joyner,<sup>c,d\*</sup> John P. Wares,<sup>c,e</sup> Kathryn P. Sutherland,<sup>f</sup> Erin K. Lipp,<sup>d</sup> Jeffrey W. Turner<sup>a</sup>

<sup>a</sup>Department of Life Sciences, Texas A&M University at Corpus Christi, Corpus Christi, Texas, USA

<sup>b</sup>Department of Marine Biology, Texas A&M University at Galveston, Galveston, Texas, USA

<sup>c</sup>Odum School of Ecology, University of Georgia, Athens, Georgia, USA

<sup>d</sup>Department of Environmental Health Science, University of Georgia, Athens, Georgia, USA

<sup>e</sup>Department of Genetics, University of Georgia, Athens, Georgia, USA

<sup>f</sup>Department of Biology, Rollins College, Winter Park, Florida, USA

**ABSTRACT** *Serratia marcescens* is a Gram-negative bacterium causally linked to acroporid serratosiis, a form of white pox disease implicated in the decline of elkhorn corals. We report draft genomes of 38 *S. marcescens* isolates collected from host and nonhost sources. The availability of these genomes will aid future analyses of acroporid serratosiis.

*Serratia marcescens* is a widely distributed Gram-negative bacillus within the *Enterobacteriaceae* family (1). The species has long been recognized as an important pathogen of humans (1, 2), insects (3–5), and plants (6). Two ecotypes (pulsed-field gel electrophoresis [PFGE] types PDL100 and PDR60) were identified as causative agents of acroporid serratosiis (a form of white pox disease) in reef-building *Acropora palmata* corals (7, 8). Given the ecological importance of *A. palmata*, it is imperative to gain a better understanding of the genetic mechanisms underlying acroporid serratosiis and what sets the PDL100 and PDR60 ecotypes apart from other pathogenic strains.

Thirty-five *S. marcescens* PDR60 isolates were collected from a range of host (*A. palmata*) and nonhost (*Siderastrea siderea* and *Solenastrea bournoni* corals, corallivorous snail *Coralliophila abbreviata*, and wastewater) sources throughout the Florida Keys National Marine Sanctuary (Table 1) (8–10). The WWI31 isolate (obtained from wastewater influent) was virulent against *A. palmata* but had a novel PFGE pattern (8). The PDL100 isolate was obtained previously from diseased *A. palmata* in 1999 (7), and the ATCC 13880 isolate was obtained by others from pond water in the Czech Republic and deposited to ATCC in 1961. Starting with glycerol stocks, each isolate was streaked onto Trypticase soy agar plates (37°C overnight), and isolated colonies were grown in lysogeny broth (37°C overnight with shaking). Total DNA for Illumina sequencing was isolated using a Qiagen DNeasy blood and tissue kit (Qiagen, Valencia, CA, USA) per the manufacturer's instructions. Genomic DNA for PacBio sequencing was isolated using a cetyltrimethylammonium bromide protocol (11). Illumina sequencing libraries were prepared using a PCR-free TrueSeq DNA kit (Illumina, San Diego, CA, USA) and sequenced at Hudson Alpha Institute for Biotechnology (Huntsville, AL, USA) using paired-end chemistry with 250-bp (EL1 and EL119), 150-bp (EL1, EL119, and KS10), or 100-bp (remaining 35 isolates) read lengths. Additionally, six isolates (EL1, EL116, EL119, EL41, EL60, and KS10) were also sequenced on three PacBio single-molecule real-time (SMRT) cells at the Interdisciplinary Center for Biotechnology Research (University of Florida, Gainesville, FL,

**Citation** Elledge NC, Eytan RI, Pinnell LJ, Pray R, Joyner JL, Wares JP, Sutherland KP, Lipp EK, Turner JW. 2019. Draft genome sequences of 38 *Serratia marcescens* isolates associated with acroporid serratosiis. *Microbiol Resour Announc* 8:e00194-19. <https://doi.org/10.1128/MRA.00194-19>.

**Editor** J. Cameron Thrash, University of Southern California

**Copyright** © 2019 Elledge et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Jeffrey W. Turner, [jeffrey.turner@tamucc.edu](mailto:jeffrey.turner@tamucc.edu).

\* Present address: Jessica L. Joyner, Department of Biology, Georgia State University, Atlanta, Georgia, USA.

N.C.E. and R.I.E. contributed equally to this work.

**Received** 19 February 2019

**Accepted** 4 March 2019

**Published** 4 April 2019

**TABLE 1** Accession numbers, genome assembly metrics, and sources for the 38 *S. marcescens* isolates<sup>a</sup>

Isolate	Accession no.	k-mer size	No. of contigs	$N_{50}$ value	G+C content (%)	Genome size (bp)	Coverage (x)	No. of genes	Source
EL1 <sup>b</sup>	CP027796	77	2	5,201,691	59.46	5,240,588	776	4,887	<i>S. siderea</i>
EL116 <sup>b</sup>	PXZP00000000	51	6	1,493,212	59.47	5,254,956	87	4,958	<i>A. palmata</i>
EL119 <sup>b</sup>	PXZQ00000000	77	3	4,696,493	59.45	5,250,706	712	4,895	<i>A. palmata</i>
EL41 <sup>b</sup>	PXZR00000000	51	7	1,445,005	59.46	5,263,255	81	4,998	Wastewater influent
EL60 <sup>b</sup>	PXZS00000000	51	12	621,937	59.45	5,223,299	317	4,957	<i>A. palmata</i>
KS10 <sup>b</sup>	CP027798	77	2	5,199,459	59.45	5,238,337	152	4,882	<i>C. abbreviata</i>
EL3	RCEP00000000	29	300	43,061	59.34	5,261,854	20	5,110	<i>S. siderea</i>
EL6	RCEO00000000	33	386	28,046	59.43	5,272,664	20	5,146	<i>S. siderea</i>
EL84	RCEN00000000	25	221	51,500	59.46	5,178,755	17	5,037	<i>C. abbreviata</i>
EL85	RCEM00000000	27	319	39,739	59.41	5,216,376	19	5,133	<i>C. abbreviata</i>
EL95	RCEL00000000	25	469	24,414	59.39	5,160,448	18	5,238	<i>A. palmata</i>
EL96	RCEK00000000	23	576	18,069	59.32	5,209,705	19	5,321	<i>A. palmata</i>
EL97	RCEJ00000000	25	737	13,342	59.25	5,163,374	22	5,439	<i>A. palmata</i>
EL98	RCEI00000000	29	549	19,346	59.42	5,177,503	24	5,294	<i>A. palmata</i>
EL108	RCEG00000000	45	35	681,033	59.46	5,226,639	39	4,912	<i>S. bournoni</i>
EL109	RCEH00000000	27	424	26,182	59.36	5,274,065	22	5,209	<i>S. bournoni</i>
EL110	RCEF00000000	31	274	39,152	59.43	5,217,630	24	5,075	<i>S. bournoni</i>
EL113	RCEE00000000	25	251	52,631	59.45	5,193,348	19	5,064	<i>A. palmata</i>
EL114	RCED00000000	31	230	47,281	59.45	5,195,401	19	5,054	<i>A. palmata</i>
EL115	RCEC00000000	27	293	42,433	59.38	5,342,905	22	5,108	<i>A. palmata</i>
EL117	RCEB00000000	27	524	20,600	59.34	5,169,891	19	5,302	<i>A. palmata</i>
EL118	RCEA00000000	23	552	18,973	59.30	5,162,774	20	5,292	<i>A. palmata</i>
EL120	RCDZ00000000	23	522	22,383	59.31	5,228,290	21	5,292	<i>A. palmata</i>
EL121	RCDY00000000	23	497	20,434	59.31	5,224,848	18	5,264	<i>A. palmata</i>
EL122	RCDX00000000	23	546	19,553	59.33	5,210,136	20	5,285	<i>A. palmata</i>
KS1	RCDW00000000	31	509	20,965	59.43	5,176,180	29	5,247	<i>C. abbreviata</i>
KS5	RCDV00000000	29	493	20,749	59.41	5,189,506	25	5,266	<i>C. abbreviata</i>
KS9	RCDU00000000	25	372	29,264	59.40	5,165,351	19	5,152	<i>C. abbreviata</i>
KS12	RCDT00000000	31	299	39,028	59.29	5,312,456	21	5,107	<i>C. abbreviata</i>
KS16	RCDS00000000	35	262	42,604	59.46	5,193,145	24	5,037	<i>C. abbreviata</i>
KS23	RCDR00000000	31	194	57,502	59.60	5,172,913	22	4,921	<i>C. abbreviata</i>
KS25	RCDQ00000000	29	213	51,510	59.46	5,207,877	21	5,050	<i>C. abbreviata</i>
KS40	RCDP00000000	29	468	22,926	59.44	5,226,754	25	5,236	<i>C. abbreviata</i>
KS45	RCD00000000	29	177	34,539	59.38	5,267,350	22	5,011	<i>C. abbreviata</i>
KS65	RCDN00000000	33	301	36,304	59.43	5,211,114	25	5,095	<i>C. abbreviata</i>
WWI31	RCDM00000000	27	324	34,866	59.86	5,127,259	16	5,009	Wastewater influent
PDL100 <sup>c</sup>	RCDL00000000	27	494	22,127	59.20	4,963,330	15	5,089	<i>A. palmata</i>
ATCC 13880 <sup>d</sup>	RWJO00000000	33	661	13,688	59.54	5,044,366	31	5,238	Pond water

<sup>a</sup> All isolates, except the ATCC isolate, were collected in the Florida Keys National Marine Sanctuary (USA).

<sup>b</sup> Hybrid assemblies.

<sup>c</sup> Deposited to ATCC in 2002.

<sup>d</sup> Deposited to ATCC in 1961.

USA). Adapter sequences and low-quality bases were removed from Illumina reads with TrimGalore! version 0.4.0 (options, -paired and -retain\_unpaired) (12). The processed reads were assembled *de novo* with Velvet version 1.2.10 (options -scaffolding, no; -exp\_cov, 80; -cov\_cutoff, 10; -min\_contig\_lgth, 500) (13) using the k-mer sizes listed in Table 1. Hybrid assemblies were constructed with MaSuRCA version 3.2.1 (options, default) (14) using the mean insert sizes and insert size standard deviations calculated with BWA (15). Assembly metrics were determined using QUAST version 5.0.2 (options, default) (16). All genomes were annotated using the National Center for Biotechnology Information (NCBI) Prokaryotic Genome Annotation Pipeline (PGAP) (17).

Table 1 shows summaries of the 38 draft genome assemblies. The availability of these genomes will aid in more comprehensive analyses of *S. marcescens* and acroporid serratiosis.

**Data availability.** This whole-genome shotgun project has been deposited in GenBank under the accession numbers listed in Table 1. The raw sequence reads were deposited in the Sequence Read Archive under the BioProject accession numbers [PRJNA494152](https://www.ncbi.nlm.nih.gov/bioproject/PRJNA494152) (nonhybrid assemblies) and [PRJNA438529](https://www.ncbi.nlm.nih.gov/bioproject/PRJNA438529) (hybrid assemblies).

## ACKNOWLEDGMENT

This project was funded by NSF-NIH Ecology of Infectious Disease program grants EF1015342 (awarded to Erin K. Lipp and John P. Wares) and EF1015032 (awarded to Kathryn P. Sutherland).

## REFERENCES

- Grimont PAD, Grimont F. 1978. The genus *Serratia*. *Annu Rev Microbiol* 32:221–248. <https://doi.org/10.1146/annurev.mi.32.100178.001253>.
- Hejazi A, Falkiner FR. 1997. *Serratia marcescens*. *J Med Microbiol* 46: 903–912. <https://doi.org/10.1099/00222615-46-11-903>.
- Bell JV, King EG, Hamalle RJ. 1974. Interactions between bollworms, a braconid parasite, and the bacterium *Serratia marcescens*. *Ann Entomol Soc Am* 67:712–714. <https://doi.org/10.1093/aesa/67.4.712>.
- Lysenko O. 1985. Non-sporeforming bacteria pathogenic to insects: incidence and mechanisms. *Annu Rev Microbiol* 39:673–695. <https://doi.org/10.1146/annurev.mi.39.100185.003325>.
- Burritt NL, Foss NJ, Neeno-Eckwall EC, Church JO, Hilger AM, Hildebrand JA, Warshauer DM, Perna NT, Burritt JB. 2016. Sepsis and hemocyte loss in honey bees (*Apis mellifera*) infected with *Serratia marcescens* strain sicaria. *PLoS One* 11:e0167752. <https://doi.org/10.1371/journal.pone.0167752>.
- Bruton BD, Mitchell F, Fletcher J, Pair SD, Wayadande A, Melcher U, Brady J, Bextine B, Popham TW. 2003. *Serratia marcescens*, a phloem-colonizing, squash bug-transmitted bacterium: causal agent of cucurbit yellow vine disease. *Plant Dis* 87:937–944. <https://doi.org/10.1094/PDIS.2003.87.8.937>.
- Patterson KL, Porter JW, Ritchie KB, Polson SW, Mueller E, Peters EC, Santavy DL, Smith GW. 2002. The etiology of white pox, a lethal disease of the Caribbean elkhorn coral, *Acropora palmata*. *Proc Natl Acad Sci U S A* 99:8725–8730. <https://doi.org/10.1073/pnas.092260099>.
- Sutherland KP, Shaban S, Joyner JL, Porter JW, Lipp EK. 2011. Human pathogen shown to cause disease in the threatened elkhorn coral *Acropora palmata*. *PLoS One* 6:e23468. <https://doi.org/10.1371/journal.pone.0023468>.
- Sutherland KP, Porter JW, Turner JW, Thomas BJ, Looney EE, Luna TP, Meyers MK, Futch JC, Lipp EK. 2010. Human sewage identified as likely source of white pox disease of the threatened Caribbean elkhorn coral, *Acropora palmata*. *Environ Microbiol* 12:1122–1131. <https://doi.org/10.1111/j.1462-2920.2010.02152.x>.
- Joyner JL, Sutherland KP, Kemp DW, Berry B, Griffin A, Porter JW, Amador MHB, Noren HKG, Lipp EK. 2015. Systematic analysis of white pox disease in *Acropora palmata* of the Florida Keys and the role of *Serratia marcescens*. *Appl Environ Microbiol* 81:4451–4457. <https://doi.org/10.1128/AEM.00116-15>.
- Wilson K. 2001. Preparation of genomic DNA from bacteria. *Curr Protoc Mol Biol* 56:2.4.1–2.4.5. <https://doi.org/10.1002/0471142727.mb0204s56>.
- Krueger F. 2015. Trim galore. A wrapper tool around Cutadapt and FastQC to consistently apply quality and adapter trimming to FastQ files. [http://www.bioinformatics.babraham.ac.uk/projects/trim\\_galore/](http://www.bioinformatics.babraham.ac.uk/projects/trim_galore/).
- Zerbino D, Birney E. 2008. Velvet: algorithms for de novo short read assembly using de Bruijn graphs. *Genome Res* 18:821–829. <https://doi.org/10.1101/gr.074492.107>.
- Zimin AV, Marçais G, Puiu D, Roberts M, Salzberg SL, Yorke JA. 2013. The MaSuRCA genome assembler. *Bioinformatics* 29:2669–2677. <https://doi.org/10.1093/bioinformatics/btt476>.
- Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* 25:1754–1760. <https://doi.org/10.1093/bioinformatics/btp324>.
- Gurevich A, Saveliev V, Vyahhi N, Tesler G. 2013. QUILT: quality assessment tool for genome assemblies. *Bioinformatics* 29:1072–1075. <https://doi.org/10.1093/bioinformatics/btt086>.
- Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44: 6614–6624. <https://doi.org/10.1093/nar/gkw569>.