



# Draft Genome Sequence of Clinical Isolate *Alcaligenaceae* sp. Strain 429

Michael J. Zilliox,<sup>a</sup> Paul C. Schreckenberger,<sup>b,†</sup>  Catherine Putonti<sup>c,d,e,f</sup>

<sup>a</sup>Department of Public Health Sciences, Loyola University Chicago, Maywood, Illinois, USA

<sup>b</sup>Department of Pathology, Loyola University Chicago, Maywood, Illinois, USA

<sup>c</sup>Department of Computer Science, Loyola University Chicago, Chicago, Illinois, USA

<sup>d</sup>Bioinformatics Program, Loyola University Chicago, Chicago, Illinois, USA

<sup>e</sup>Department of Biology, Loyola University Chicago, Chicago, Illinois, USA

<sup>f</sup>Department of Microbiology and Immunology, Loyola University Chicago, Maywood, Illinois, USA

**ABSTRACT** Here, we present the 3.53-Mb genome for *Alcaligenaceae* sp. strain 429, isolated from a patient with unknown etiology. While the 16S rRNA gene most closely resembles *Paenalcaligenes* species, average amino acid identity (AAI) analysis did not meet the threshold to classify our strain as a species of this family.

The bacterial isolate was received from Elmhurst Hospital in Elmhurst, IL, in 2012 for identification. No patient data are available. Prior to sequencing, identification was performed using a matrix-assisted laser desorption ionization (MALDI) Biotyper (Bruker Scientific, Billerica, MA, USA), which classified the strain as an *Alcaligenes* sp., whereas the Vitek MS system (bioMérieux, Marcy-l'Étoile, France) found no match. However, 16S rRNA gene sequencing performed by our group indicated that the isolate was more closely related to another member of the *Alcaligenaceae* family, *Paenalcaligenes suwonensis*. *Paenalcaligenes* species have been isolated from both the environment (1) and patients (2, 3), and only one genome has been sequenced for the genus to date (*P. hominis*). Furthermore, it is rarely associated with clinical symptoms (2, 3). In contrast, infections attributed to *Alcaligenes faecalis* have been reported, although they are rare and nosocomial (4–8). To resolve the taxonomy of this clinical isolate, we sequenced the genome.

The isolate was originally saved in Brucella broth with 10% glycerol and frozen at –80°C. The frozen sample was then subcultured onto blood agar plates and incubated at 35°C in 5% CO<sub>2</sub> for 24 to 48 hours. Genomic DNA was extracted from bacterial cells using a validated mixture of lysozyme and mutanolysin and the DNeasy blood and tissue kit (Qiagen, Valencia, CA) (9). All steps were performed in a UV-irradiated, HEPA-filtered PCR workstation to minimize potential contaminants. The library was constructed using the Nextera XT kit (Illumina, San Diego, CA) following the manufacturer's protocol and sequenced on an Illumina MiSeq platform using a 500-cycle v2 kit (250 × 2 bp) rendering 1,336,945 paired-end reads.

The raw reads first were trimmed using Sickle v1.33 (<https://github.com/najoshi/sickle>), specifying that trimmed reads of ≤30 nucleotides be removed from consideration (“-l 30”), and then assembled with SPAdes v3.11.1 using the “careful” option (10). Genome coverage was calculated for the contigs using the BBDMap tool v37 (<http://sourceforge.net/projects/bbmap/>). Contigs with a length less than 600 bp or a coverage less than 60 (1 standard deviation from the mean coverage) were removed from further consideration. The coverage was recalculated using BBDMap as 98.64×. The remaining 391 contigs were queried via megablast against the nonredundant/nucleotide (nr/nt) database to confirm the genus and species of the isolate; all hits were to sequences of

**Citation** Zilliox MJ, Schreckenberger PC, Putonti C. 2019. Draft genome sequence of clinical isolate *Alcaligenaceae* sp. strain 429. *Microbiol Resour Announc* 8:e00439-19. <https://doi.org/10.1128/MRA.00439-19>.

**Editor** Frank J. Stewart, Georgia Institute of Technology

**Copyright** © 2019 Zilliox et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Catherine Putonti, [cputonti@luc.edu](mailto:cputonti@luc.edu).

† Deceased.

**Received** 12 April 2019

**Accepted** 22 April 2019

**Published** 9 May 2019

strains belonging to the genera *Paenalcaligenes* and *Alcaligenes*. The contigs have a total length of 3,539,013 nucleotides with a GC content of 50.1% and an  $N_{50}$  value of 14,837 bp. The genome was annotated using the NCBI Prokaryotic Genome Annotation Pipeline (11). Three rRNA (1 5S, 1 16S, and 1 23S) gene sequences, 48 tRNA gene sequences, and 3,422 protein coding genes were identified.

The 16S rRNA sequence from the assembled genome was compared to NCBI's 16S rRNA database via blastn, identifying the best hit to the *P. suwonensis* strain ABC02-12 sequence (GenBank accession number [NR\\_133804](https://doi.org/10.1093/nar/gky359); query coverage, 93%; identity, 99.16%). Average amino acid identity (AAI), calculated using the AAI-profiler online server (12), identified the nearest neighbor as *Alcaligenes faecalis* (72%). The number of coding regions shared between our isolate and *Paenalcaligenes* and *Alcaligenes* strains, however, did not exceed 56%. This finding supports recent evidence suggesting that the *Alcaligenes* genus has an open pangenome (13). Thus, at this time, no genus designation can be made.

**Data availability.** The draft whole-genome project for *Alcaligenaceae* sp. strain 429 has been deposited at DDBJ/EMBL/GenBank under accession number [SRSN00000000](https://doi.org/10.1093/nar/gky359). Raw sequence reads are deposited at DDBJ/EMBL/GenBank under accession number [SRR8862657](https://doi.org/10.1093/nar/gky359).

## ACKNOWLEDGMENTS

We acknowledge Kathleen McKinley for bacterial culture and isolation and Gina Kuffel for whole-genome sequencing of the isolate.

## REFERENCES

- Moon J-Y, Lim J-M, Ahn J-H, Weon H-Y, Kwon S-W, Kim S-J. 2014. *Paenalcaligenes suwonensis* sp. nov., isolated from spent mushroom compost. *Int J Syst Evol Microbiol* 64:882–886. <https://doi.org/10.1099/ijs.0.058412-0>.
- Mukhopadhyay R, Joaquin J, Hogue R, Kilaru A, Jospin G, Mars K, Eisen JA, Chaturvedi V. 2017. Complete genome sequence of a *Paenalcaligenes hominis* strain isolated from a paraplegic patient with neurogenic bladder using single-molecule real-time sequencing technology. *Genome Announc* 5:e00252-17. <https://doi.org/10.1128/genomeA.00252-17>.
- Kämpfer P, Falsen E, Langer S, Lodders N, Busse H-J. 2010. *Paenalcaligenes hominis* gen. nov., sp. nov., a new member of the family *Alcaligenaceae*. *Int J Syst Evol Microbiol* 60:1537–1542. <https://doi.org/10.1099/ijs.0.016576-0>.
- Bizet J, Bizet C. 1997. Strains of *Alcaligenes faecalis* from clinical material. *J Infect* 35:167–169. [https://doi.org/10.1016/S0163-4453\(97\)91710-2](https://doi.org/10.1016/S0163-4453(97)91710-2).
- Ashwath ML, Katner HP. 2005. Pancreatic abscess secondary to *Alcaligenes faecalis*. *Am J Med Sci* 329:54–55. <https://doi.org/10.1097/0000441-200501000-00011>.
- Pal SS, Panigrahi PK, Roy R, Nandi K, Das S. 2013. Endophthalmitis caused by *Alcaligenes faecalis*: a case series. *Ocul Immunol Inflamm* 21:446–448. <https://doi.org/10.3109/09273948.2013.817592>.
- Tena D, Fernández C, Lago MR. 2015. *Alcaligenes faecalis*: an unusual cause of skin and soft tissue infection. *Jpn J Infect Dis* 68:128–130. <https://doi.org/10.7883/yoken.JJID.2014.164>.
- Chu AS, Harkness J. 2017. *Alcaligenes faecalis* cellulitis after a dog bite: case report and literature review. *Pediatr Emerg Care* 33:497–498. <https://doi.org/10.1097/PEC.0000000000000645>.
- Yuan S, Cohen DB, Ravel J, Abdo Z, Forney LJ. 2012. Evaluation of methods for the extraction and purification of DNA from the human microbiome. *PLoS One* 7:e33865. <https://doi.org/10.1371/journal.pone.0033865>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
- Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>.
- Medlar AJ, Törönen P, Holm L. 2018. AAI-profiler: fast proteome-wide exploratory analysis reveals taxonomic identity, misclassification and contamination. *Nucleic Acids Res* 46:W479–W485. <https://doi.org/10.1093/nar/gky359>.
- Basharat Z, Yasmin A, He T, Tong Y. 2018. Genome sequencing and analysis of *Alcaligenes faecalis* subsp. phenolicus MB207. *Sci Rep* 8:3616. <https://doi.org/10.1038/s41598-018-21919-4>.