



Complete Genome Sequence of Hypervirulent *Streptococcus pyogenes emm3* Strain 1838

Benfang Lei,^a Anthony R. Flores,^b Carl Yeoman,^c Mengyao Liu^a

^aDepartment of Microbiology and Immunology, Montana State University, Bozeman, Montana, USA

^bDivision of Infectious Diseases, Department of Pediatrics, Center for Antimicrobial Resistance and Microbial Genomics, McGovern Medical School, University of Texas Health Science Center at Houston, Houston, Texas, USA

^cDepartment of Animal & Range Sciences, Montana State University, Bozeman, Montana, USA

ABSTRACT We report the complete genome sequence for *Streptococcus pyogenes* strain 1838 (type *emm3*) isolated from a patient with toxic shock syndrome. The strain lacked the *speK*- and *sla*-encoding prophage frequently encountered among *emm3* strains and possessed an Arg66His mutation in CovR of the 2-component virulence regulatory system CovRS.

Streptococcus pyogenes (group A *Streptococcus* [GAS]) strain 1838 was isolated from a patient with toxic shock syndrome in 2011 by the Streptococcus Laboratory at the Centers for Disease Control and Prevention (CDC) (1). The strain was included in a study to compare contemporary pharyngeal and invasive *S. pyogenes* isolates in their capacity to acquire *covRS* mutations using a mouse model of skin infection (2). CovRS (also known as CrsRS) is a 2-component regulatory system known to regulate expression of multiple GAS virulence factors (3–5). Naturally occurring CovRS mutations enhance expression of virulence genes and simultaneously down-regulate the protease *speB*, which leads to hypervirulence (6–8). Among 6 SpeB-positive *emm3* GAS strains, mice infected with strain 1838 demonstrated increased mortality in an *in vivo* subcutaneous infection assay for selection of CovRS mutants (2). Like *emm3* strain MGAS315 (9), strain 1838 can invade the vascular system in a mouse model of pulmonary infection.

To understand the basis for hypervirulence, we sequenced the genome of strain 1838 using reads generated with both the PacBio RS II system and the Illumina MiSeq (300-bp, paired-end) instrument by the W.M. Keck Foundation Biotechnology Resource Laboratory at Yale University and Otogenetics Corporation, respectively. For DNA extraction, bacteria were streaked from a vial of the frozen isolate from the CDC on a Todd-Hewitt broth/2% yeast extract agar plate and incubated for 8 h at 37°C in 5% CO₂. Genomic DNA from the bacteria was extracted using the MasterPure Gram-positive DNA purification kit (Lucigen/Epicentre, catalog number MGP04100) following the manufacturer's protocol with the following modifications: bacteria from the plate were washed with 1-ml phosphate-buffered saline (PBS) 3 times and treated with 0.2-μg proteinase K in 1-ml PBS at 37°C for 2 h; the bacteria were pelleted by centrifugation, washed with 1-ml PBS 3 times, and resuspended in 150-μl Tris-EDTA buffer; and the bacterial suspension was mixed first with 10 μl of 1.0-mg/ml PlyC (10) and then with 150-μl lysis buffer and a 175-μl precipitation solution of the MasterPure kit. PacBio sequencing generated 132,929 reads with an average read length of 9,788 bp, and Illumina sequencing generated 14,839,914 reads and 1,869,829,164 bp. Hybrid assembly (i.e., long- and short-read sequences) using the SPAdes assembler (v 3.12.0) (11) and default parameters yielded a complete genome with greater than 1000× coverage. The complete genome was polished using Pilon (v 1.22) (12). The polished complete

Citation Lei B, Flores AR, Yeoman C, Liu M. 2019. Complete genome sequence of hypervirulent *Streptococcus pyogenes emm3* strain 1838. Microbiol Resour Announc 8:e01494-18. <https://doi.org/10.1128/MRA.01494-18>.

Editor David A. Baltus, University of Arizona

Copyright © 2019 Lei et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Benfang Lei, blei@montana.edu.

Received 30 October 2018

Accepted 3 December 2018

Published 10 January 2019

genome was annotated using the Prokaryotic Genome Annotation Pipeline at the National Center for Biotechnology Information (13).

In comparison with the genome of the *emm3* strain MGAS315 (14), the strain 1838 genome shows a chromosomal inversion known to occur in approximately one-quarter of GAS genomes, presumably from a recombination event at homologous copies of *comX/sigX* (15). The inversion was identified using whole-genome alignments to completed serotype M3 GAS genomes with progressiveMauve (16). The inversion was verified by aligning both short and long reads to the completed 1838 genome and thus is unlikely to be a misassembly. Analysis of the strain 1838 genome with PHASTER (17) found 4 intact prophages, encoding *speC*, *speA*, and *spd1*. From the mid-1980s through in the early 2000s, virtually all *emm3* strains had a 315.4-like prophage encoding *speK* and *sla* (14). However, like many *emm3* strains from approximately 2007 to 2009 (18), strain 1838 lacked the *speK*- and *sla*-encoding prophage. Strain 1838 does have 315.3-, 315.5-, and 315.6-like prophages. Polymorphisms relative to MGAS315 were identified with NUCmer (v 3.1) (19) and by mapping Illumina short-read sequences from strain 1838 to MGAS315 with a custom pipeline as described by Long et al. (20). Excluding the prophage sequences, there were 189 single-nucleotide polymorphisms (SNPs) and 13 insertions/deletions (indels) in strain 1838 relative to MGAS315. Based on the chromosomal SNP data, strain 1838 appears to be closely related to *emm3* strains that were responsible for a dramatic upsurge of M3 invasive infections in the United Kingdom in 2008 and 2009 (18) and consistent with currently circulating *emm3* strains in the United States (21).

The 189 chromosomal SNPs between MGAS315 and strain 1838 contain 39 missense mutations, including one in each of *covR*, *covS*, and *ropB*. The *ropB* and *covS* SNPs are consistent with known polymorphisms in MGAS315 (8, 22), and the SNP in *covR* results in an arginine-to-histidine amino acid change at position 66 (Arg66His) in strain 1838 CovR. No other mutations were found in the known virulence genes and regulators. The Arg66His mutation is near the phosphorylation sites of CovR (aspartate at position 53, D53, and threonine at position 65, T65) (23), which may affect the phosphorylation of CovR and lead to enhanced virulence gene expression and hypervirulence of strain 1838.

Data availability. The genome sequence of GAS strain 1838 has been deposited in GenBank with the accession number [CP029694](https://doi.org/10.1093/cid/ciw248). Raw sequences were deposited in the NCBI SRA database under BioProject number [PRJNA473837](https://doi.org/10.1093/cid/ciw248).

ACKNOWLEDGMENTS

This work was supported in part by grants AI095704 and GM110732 from the National Institutes of Health and the Montana State Agricultural Experimental Station.

We thank Bernard Beall and Chris A. Van Beneden at the National Center for Immunization and Respiratory Diseases, Centers for Disease Control and Prevention, for providing strain 1838.

REFERENCES

- Nelson GE, Pondo T, Toews KA, Farley MM, Lindegren ML, Lynfield R, Aragon D, Zansky SM, Watt JP, Cieslak PR, Angeles K, Harrison LH, Petit S, Beall B, Van Beneden CA. 2016. Epidemiology of invasive group A streptococcal infections in the United States, 2005–2012. *Clin Infect Dis* 63:478–486. <https://doi.org/10.1093/cid/ciw248>.
- Feng W, Liu M, Chen DG, Yiu R, Fang FC, Lei B. 2016. Contemporary pharyngeal and invasive *emm1* and invasive *emm12* group A *Streptococcus* isolates exhibit similar *in vivo* selection for CovRS mutants in mice. *PLoS One* 11:e0162742. <https://doi.org/10.1371/journal.pone.0162742>.
- Levin JC, Wessels MR. 1998. Identification of *csrR/csrS*, a genetic locus that regulates hyaluronic acid capsule synthesis in group A *Streptococcus*. *Mol Microbiol* 30:209–219.
- Federle MJ, McIver KS, Scott JR. 1999. A response regulator that represses transcription of several virulence operons in the group A *Streptococcus*. *J Bacteriol* 181:3649–3657.
- Treviño J, Perez N, Ramirez-Peña E, Liu Z, Shelburne SA, III, Musser JM, Sumbly P. 2009. CovS simultaneously activates and inhibits the CovR-mediated repression of distinct subsets of group A *Streptococcus* virulence factor-encoding genes. *Infect Immun* 77:3141–3149. <https://doi.org/10.1128/IAI.01560-08>.
- Sumbly P, Whitney AR, Graviss EA, DeLeo FR, Musser JM. 2006. Genome-wide analysis of group A streptococci reveals a mutation that modulates global phenotype and disease specificity. *PLoS Pathog* 2:41–49. <https://doi.org/10.1371/journal.ppat.0020005>.
- Li J, Zhu H, Feng W, Liu M, Song Y, Zhang X, Zhou Y, Bei W, Lei B. 2013. Regulation of inhibition of neutrophil infiltration by the two-component regulatory system CovRS in subcutaneous murine infection with group A *Streptococcus*. *Infect Immun* 81:974–983. <https://doi.org/10.1128/IAI.01218-12>.
- Stetzner ZW, Li D, Feng W, Liu M, Liu G, Wiley J, Lei B. 2015. Serotype M3 and M28 group A streptococci have distinct capacities to evade neutro-

- phil and TNF- α responses and to invade soft tissues. *PLoS One* 10: e0129417. <https://doi.org/10.1371/journal.pone.0129417>.
9. Lei B, Minor D, Feng W, Liu M. 2018. Hypervirulent group A *Streptococcus* of genotype *emm3* invades the vascular system in pulmonary infection of mice. *Infect Immun* 86:e00080-18. <https://doi.org/10.1128/IAI.00080-18>.
 10. Shen Y, Köller T, Kreikemeyer B, Nelson DC. 2013. Rapid degradation of *Streptococcus pyogenes* biofilms by PlyC, a bacteriophage-encoded endolysin. *J Antimicrob Chemother* 68:1818–1824. <https://doi.org/10.1093/jac/dkt104>.
 11. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
 12. Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, Cuomo CA, Zeng Q, Wortman J, Young SK, Earl AM. 2014. Pilon: an integrated tool for comprehensive microbial variant detection and genome assembly improvement. *PLoS One* 9:e112963. <https://doi.org/10.1371/journal.pone.0112963>.
 13. Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44: 6614–6624. <https://doi.org/10.1093/nar/gkw569>.
 14. Beres SB, Sylva GL, Barbian KD, Lei B, Hoff JS, Mammarella ND, Liu M-Y, Smoot JC, Porcella SF, Parkins LD, Campbell DS, Smith TM, McCormick JK, Leung DY, Schlievert PM, Musser JM. 2002. Genome sequence of a serotype M3 strain of group A *Streptococcus*: phage-encoded toxins, the high-virulence phenotype, and clone emergence. *Proc Natl Acad Sci USA* 99:10078–10083. <https://doi.org/10.1073/pnas.152298499>.
 15. Bessen DE, McShan WM, Nguyen SV, Shetty A, Agrawal S, Tettelin H. 2015. Molecular epidemiology and genomics of group A *Streptococcus*. *Infect Genet Evol* 33:393–418. <https://doi.org/10.1016/j.meegid.2014.10.011>.
 16. Darling AE, Mau B, Perna NT. 2010. progressiveMauve: multiple genome alignment with gene gain, loss and rearrangement. *PLoS One* 5:e11147. <https://doi.org/10.1371/journal.pone.0011147>.
 17. Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang Y, Wishart DS. 2016. PHASTER: a better, faster version of the PHAST phage search tool. *Nucleic Acids Res* 44:W16–W21. <https://doi.org/10.1093/nar/gkw387>.
 18. Al-Shahib A, Underwood A, Afshar B, Turner CE, Lamagni T, Sriskandan S, Efstratiou A. 2016. Emergence of a novel lineage containing a prophage in *emm*/M3 group A *Streptococcus* associated with upsurge in invasive disease in the UK. *Microb Genom* 2:e000059. <https://doi.org/10.1099/mgen.0.000059>.
 19. Kurtz S, Phillippy A, Delcher AL, Smoot M, Shumway M, Antonescu C, Salzberg SL. 2004. Versatile and open software for comparing large genomes. *Genome Biol* 5:R12. <https://doi.org/10.1186/gb-2004-5-2-r12>.
 20. Long SW, Olsen RJ, Eagar TN, Beres SB, Zhao P, Davis JJ, Brettin T, Xia F, Musser JM. 2017. Population genomic analysis of 1,777 extended-spectrum beta-lactamase-producing *Klebsiella pneumoniae* isolates, Houston, Texas: unexpected abundance of clonal group 307. *mBio* 8:e00489-17. <https://doi.org/10.1128/mBio.00489-17>.
 21. Chochua S, Metcalf BJ, Li Z, Rivers J, Mathis S, Jackson D, Gertz RE, Jr, Srinivasan V, Lynfield R, Van Beneden C, McGee L, Beall B. 2017. Population and whole genome sequence based characterization of invasive group A *Streptococci* recovered in the United States during 2015. *mBio* 8:e01422-17. <https://doi.org/10.1128/mBio.01422-17>.
 22. Kappeler KV, Anbalagan S, Dmitriev AV, McDowell EJ, Neely MN, Chaussee MS. 2009. A naturally occurring Rgg variant in serotype M3 *Streptococcus pyogenes* does not activate *speB* expression due to altered specificity of DNA binding. *Infect Immun* 77:5411–5417. <https://doi.org/10.1128/IAI.00373-09>.
 23. Horstmann N, Saldaña M, Sahasrabhojane P, Yao H, Su X, Thompson E, Koller A, Shelburne SA, III. 2014. Dual-site phosphorylation of the control of virulence regulator impacts group A streptococcal global gene expression and pathogenesis. *PLoS Pathog* 10:e1004088. <https://doi.org/10.1371/journal.ppat.1004088>.