



Complete Genome Sequence of Genotype Psittacine Beak and Feather Disease Virus, a Strain Identified from Budgerigars in China

Rujia Cheng,^{a,b} Yaqing Mao,^{a,b} Qiuchen Li,^{a,b} Shijie Xie,^a Yingju Xia,^a Haojie Sun,^a Kai Niu,^{a,b} Shijing Sun,^a Jie Li,^c Yu Feng,^a Xiaowei Peng,^a Hui Jiang,^a Liangquan Zhu,^a Xuezheng Fan,^a Zhanye Lin,^d Jiabo Ding,^a Yuming Qin,^a Guanlong Xu^a

^aNational Reference Laboratory for Animal Brucellosis, China Institute of Veterinary Drug Control, Beijing, China

^bCollege of Veterinary Medicine, Shandong Agricultural University, Tai'an, Shandong, China

^cDepartment of Population Medicine and Diagnostic Sciences, College of Veterinary Medicine, Cornell University, Ithaca, New York, USA

^dAnimal Husbandry and Veterinary Bureau, Ministry of Agriculture and Rural Affairs, Beijing, China

ABSTRACT Psittacine beak and feather disease virus (PBFVDV) has been reported in many countries, such as Australia, Poland, the United States, South Africa, etc. In this study, the complete genome of a PBFVDV isolate was determined and characterized from budgerigars in China.

Psittacine beak and feather disease virus (PBFVDV) is a nonenveloped virus with a diameter of 14 to 16 nm and belongs to the genus *Circovirus* in the family *Circoviridae*. The genome of PBFVDV is a 2-kb circular single-stranded DNA molecule possessing two major open reading frames (ORFs) encoding a replication-associated protein (Rep) and the capsid protein (Cap) (1, 2). Psittacine beak and feather disease (Pbfd) is characterized by immune suppression, loss of feathers, weight loss, and the development of morphologically abnormal feathers, and it sometimes leads to sudden death in young birds (3, 4).

PBFVDV is distributed worldwide with a wide range of hosts, including more than 60 species of psittacine birds (5). PBFVDV has been reported in many countries, such as Australia, Poland, the United States, South Africa, etc. (6–8). In this study, the full-genome sequence of PBFVDV isolated from budgerigars in Shandong Province, China, was determined and analyzed. Clinical signs of feather loss were observed in budgerigars at a breeding facility. The feather samples were collected and homogenized, and total DNA was extracted from the feather samples using a QIAamp minikit (Qiagen, Hilden, Germany). A primer pair (forward, 1,368 bp-TTGGGTCCTCCTATCGGGATC-1,392 bp; reverse, 1,868 bp-AGACACCGTTTTACAACCAATAG-1,843 bp) targeting 500 bp of the capsid gene was designed using Primer Premier version 5.0 (Canada). The DNA samples were screened using PCR, and the PCR products were sequenced using Sanger sequencing by the TsingKe Biological Technology Company (Beijing, China). All of the clinical samples were positive for the identical sequence; this indicated that these birds were infected with PBFVDV. To further characterize the pathogen, the full genome of one psittacine beak and feather disease virus isolate in the positive samples (named PBFVDV-GD) was sequenced and analyzed. DNA isolated from the samples was directly amplified using a second pair of primers to cover the full genome (forward, 1 bp-GTT ATGTAGTCAGAATTCCAAATTA-25 bp; reverse, 2,000 bp-GTTAGCTAAAATGGAAATGAGG CCCAC-1,973 bp). The resulting PCR product (2 kb) was sequenced using Sanger sequencing in both directions and walking primers. The final genome was assembled and analyzed with the Lasergene software version 7.0 (DNASTar, USA) (9).

The genome of the PBFVDV-GD isolate was found to be 2,000 bp long, with 52% GC content, and contains two ORFs which encode Rep (289 amino acids [aa]) and Cap

Citation Cheng R, Mao Y, Li Q, Xie S, Xia Y, Sun H, Niu K, Sun S, Li J, Feng Y, Peng X, Jiang H, Zhu L, Fan X, Lin Z, Ding J, Qin Y, Xu G. 2019. Complete genome sequence of genotype psittacine beak and feather disease virus, a strain identified from budgerigars in China. *Microbiol Resour Announc* 8:e00040-19. <https://doi.org/10.1128/MRA.00040-19>.

Editor Jelle Matthijnsens, KU Leuven

Copyright © 2019 Cheng et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Yuming Qin, Qinyuming73@163.com, or Guanlong Xu, Xuguanlongw@163.com.

R.C., Y.M., Q.L., and S.X. contributed equally to this work.

Received 10 January 2019

Accepted 1 March 2019

Published 16 May 2019

(237 aa). Our data showed that PBFDV-GD shares about 77% to 90% similarity with the PBFDV genomes deposited in GenBank from Australia, South Africa, and New Zealand, suggesting that the PBFDV circulating in China is different from those PBFDV strains found in Australia, South Africa, and New Zealand.

Given the extensive international bird trade, PBFDV has spread worldwide and poses a threat to psittacine birds. The PBFDV strain identified in the diseased budgerigars indicates that a different lineage of PBFDV is circulating in China. Although PBFDV was identified in the diseased budgerigars, it remains to be confirmed whether PBFDV was the sole cause of the observed disease. Further active surveillance is needed to determine the prevalence of PBFDV in China.

Data availability. The complete genome sequence reported here was submitted to GenBank under the accession number [MK120438](https://www.ncbi.nlm.nih.gov/nuclseq/MK120438).

ACKNOWLEDGMENT

This study was supported by the National Natural Science Foundation of Youth Fund (grant 31802201).

REFERENCES

- Ritchie BW, Niagro FD, Latimer KS, Lukert PD, Steffens WL, III, Rakich PM, Pritchard N. 1990. Ultrastructural, protein composition, and antigenic comparison of psittacine beak and feather disease virus purified from four genera of psittacine birds. *J Wildlife Dis* 26:196–203. <https://doi.org/10.7589/0090-3558-26.2.196>.
- Heath L, Williamson AL, Rybicki EP. 2006. The capsid protein of beak and feather disease virus binds to the viral DNA and is responsible for transporting the replication-associated protein into the nucleus. *J Virol* 80:7219–7225. <https://doi.org/10.1128/JVI.02559-05>.
- Ritchie BW, Niagro FD, Lukert PD, Steffens WL, III, Latimer KS. 1989. Characterization of a new virus from cockatoos with psittacine beak and feather disease. *Virology* 171:83–88. [https://doi.org/10.1016/0042-6822\(89\)90513-8](https://doi.org/10.1016/0042-6822(89)90513-8).
- Latimer KS, Rakich PM, Steffens WL, Kircher IM, Ritchie BW, Niagro FD, Lukert PD. 1991. A novel DNA virus associated with feather inclusions in psittacine beak and feather disease. *Vet Pathol* 28:300–304. <https://doi.org/10.1177/030098589102800406>.
- Varsani A, de Villiers GK, Regnard GL, Bragg RR, Kondiah K, Hitzeroth II, Rybicki EP. 2010. A unique isolate of beak and feather disease virus isolated from budgerigars (*Melopsittacus undulatus*) in South Africa. *Arch Virol* 155:435–439. <https://doi.org/10.1007/s00705-010-0589-0>.
- Ha HJ, Anderson IL, Alley MR, Springett BP, Gartrell BD. 2007. The prevalence of beak and feather disease virus infection in wild populations of parrots and cockatoos in New Zealand. *N Z Vet J* 55:235–238. <https://doi.org/10.1080/00480169.2007.36774>.
- Kondiah K, Albertyn J, Bragg RR. 2006. Genetic diversity of the Rep gene of beak and feather disease virus in South Africa. *Arch Virol* 151:2539–2545. <https://doi.org/10.1007/s00705-006-0800-5>.
- Sarker S, Das S, Ghorashi SA, Forwood JK, Raidal SR. 2014. Molecular characterization of genome sequences of beak and feather disease virus from the Australian twenty-eight parrot (*Barnardius zonarius semitorquatus*). *Genome Announc* 2:e01255-14. <https://doi.org/10.1128/genomeA.01255-14>.
- Jin JQ, Sun YB. 2018. AutoSeqMan: batch assembly of contigs for Sanger sequences. *Zool Res* 39:123–126. <https://doi.org/10.24272/j.issn.2095-8137.2018.027>.