



Draft Genome Sequence of *Nitrosomonas* sp. Strain APG5, a Betaproteobacterial Ammonia-Oxidizing Bacterium Isolated from Beach Sand

Hidetoshi Urakawa,^a Jorie Skutas,^b Jose V. Lopez^b

^aDepartment of Marine and Ecological Sciences, Florida Gulf Coast University, Fort Myers, Florida, USA

^bDepartment of Biological Sciences, Halmos College of Natural Sciences and Oceanography, Nova Southeastern University, Dania Beach, Florida, USA

ABSTRACT *Nitrosomonas* sp. strain APG5 (=NCIMB 14870 = ATCC TSA-116) was isolated from dry beach sand collected from a supralittoral zone of the northwest coast of the United States. The draft genome sequence revealed that it represents a new species of the cluster 6 *Nitrosomonas* spp. that is closely related to *Nitrosomonas ureae* and *Nitrosomonas oligotropha*.

Nitrosomonas is a genus of nitrifying bacteria belonging to the class *Betaproteobacteria*. Together with ammonia-oxidizing archaea, ammonia-oxidizing bacteria (AOB) play important roles in both natural and artificial environments. However, AOB are sensitive to environmental stressors, such as hydrocarbon contamination (1), and serve as valuable ecological indicators (2).

In 2008, *Nitrosomonas* sp. strain APG5 was isolated from a dry sand sample collected from Edmonds Ferry Dock Beach, located in the Puget Sound, WA, which is historically and routinely contaminated by small oil spills due to heavy ship transportation, including a state-run ferry system (3). Five grams of sand collected in a sterilized 50-ml centrifuge tube was incubated at 20°C with 10 ml of autoclaved ultrapure water supplemented with ammonium chloride to 100 μM as a final concentration. After the first indication of nitrite production (4), 1 ml of sample was transferred into 9 ml of APG medium, and the pure culture was obtained by a serial dilution technique (5). Initial phylogenetic analysis based on the 16S rRNA gene revealed that APG5 belongs to the cluster 6a *Nitrosomonas* spp. (6). To determine the genome sequence of APG5, cells were cultured accordingly (5), and genomic DNA was prepared using a modified phenol-chloroform extraction method, as previously described (7). A draft genome sequence was obtained using the Illumina MiSeq platform and Nextera XT sample preparation kit v2 with 2 × 250-bp paired-end reads (3,338,462 reads). Genome assembly quality was evaluated using Quast (version 4.6.3) (8), and reads were assembled using Unicycler (version 0.4.6.0) (9) on Galaxy (10) using default settings. The draft genome was initially annotated with the Rapid Annotations using Subsystems Technology (RAST) server (version 2.0) (11) and SEED Viewer (12) to confirm some house-keeping genes and key functional genes. The NCBI Prokaryotic Genome Annotation Pipeline (GeneMarkS+ version 4.4) (13) was used in the last stage of genome annotation.

The assembled draft genome sequence comprised 3.75 Mbp at 222-fold coverage and consisted of 239 contigs with an average size of 12,702 bp and N_{50} length of 56,724 bp. The G+C content was 43.5%, and one plasmid was found (pAPG501, 14,708 bp). The draft genome contains 3,172 protein-coding DNA sequences, 40 tRNA genes, and a single 16S-23S-5S rRNA operon.

The two-way average nucleotide identity values (14) of strain APG5 with *Nitrosomonas ureae* Nm10 (GenBank accession number CP013341) (15), *Nitrosomonas* sp.

Citation Urakawa H, Skutas J, Lopez JV. 2019. Draft genome sequence of *Nitrosomonas* sp. strain APG5, a betaproteobacterial ammonia-oxidizing bacterium isolated from beach sand. *Microbiol Resour Announc* 8:e01573-18. <https://doi.org/10.1128/MRA.01573-18>.

Editor J. Cameron Thrash, University of Southern California

Copyright © 2019 Urakawa et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Hidetoshi Urakawa, hurakawa@fgcu.edu.

Received 17 November 2018

Accepted 1 May 2019

Published 23 May 2019

strain AL212 (GenBank accession number [CP002552](#)) (16), and *Nitrosomonas* sp. strain Is79 (GenBank accession number [CP002876](#)) (17) were 82.2%, 81.9%, and 77.9%, respectively, which were lower than the average nucleotide identity (ANI) value of 95%, which corresponds to a 70% species level cutoff determined by DNA-DNA hybridization (18). The results evidently indicate that these genomes do not belong to the same species.

Genes involved in ammonia oxidation, including those encoding ammonia-mono-oxygenase, hydroxylamine dehydrogenase, and c-type cytochromes, were identified, as well as that encoding nitrosocyanin. As in other AOB, genes encoding nitrite reductase (*nirK*) and urease were present. The APG5 genome encodes 95 chemotaxis and flagellum-associated proteins, based on the subsystem information in the SEED Viewer. Concomitantly, genes encoding catalase and superoxide dismutase (Fe) were found for protection from reactive oxygen species. The plasmid contained a gene used to stabilize plasmid function and some ecologically significant genes. For example, a gene encoding the universal stress protein UspA, which modulates the expression of a variety of genes that help to cope with stress (19), was found in the plasmid. In addition, a sodium-proton exchanger (which maintains the homeostasis of pH and sodium), ATP-dependent exonuclease SbcCD, antitoxin Phd family protein, and addiction module toxin RelE were found in the same plasmid. These genes may increase the fitness of APG5 in supralittoral beach sand, in which microorganisms need to cope with long periods of exposure to air, heat, cold, low nutrients, and freshwater exposure through precipitation (20). Further genome annotation and genome comparisons with other *Nitrosomonas* species will provide additional insights into the ecological adaptation of this bacterium.

Data availability. The *Nitrosomonas* sp. APG5 whole-genome shotgun (WGS) project has the project accession number [PXXU00000000](#). This version of the project (01) has the accession number [PXXU01000000](#) and consists of the sequences [PXXU01000001](#) to [PXXU01000239](#). The sequencing reads (under Sequence Read Archive number [SRP170980](#)) can be accessed through BioProject number [PRJNA438189](#) and BioSample number [SAMN08707727](#). Data and additional information are publicly available through the Gulf of Mexico Research Initiative Information & Data Cooperative (GRIIDC) at <https://doi.org/10.7266/n7-fnva-0v52>.

ACKNOWLEDGMENTS

We thank David A. Stahl at the University of Washington and Willm Martens-Habbena at the University of Florida for their early contributions to this study. We also thank Megan E. Feeney (Florida Gulf Coast University) for her laboratory assistance.

A part of this research was supported by NSF Division of Environmental Biology grant DEB-1664052 to H.U. A part of this work was also supported via funding provided by the Office of Research and Graduate Studies of Florida Gulf Coast University (to H.U.) and funding from the Gulf of Mexico Research Initiative through the Alabama Center for Ecological Resilience, administered by the Dauphin Island Sea Lab (to H.U.).

J.V.L. is part of the DEEPEND consortium ([DEEPENDconsortium.org](https://deependconsortium.org)).

REFERENCES

- Urakawa H, Rajan S, Feeney ME, Sobecky PA, Mortazavi B. 2019. Ecological response of nitrification to oil spills and its impact on the nitrogen cycle. *Environ Microbiol* 21:18–33. <https://doi.org/10.1111/1462-2920.14391>.
- Urakawa H, Bernhard AE. 2017. Wetland management using microbial indicators. *Ecol Eng* 108:456–476. <https://doi.org/10.1016/j.ecoleng.2017.07.022>.
- Washington State Department of Ecology. 2018. 30 years of spill prevention, preparedness, and response. Publication 18-08-012. Spill Prevention, Preparedness, and Response Program, Washington State Department of Ecology, Olympia, WA.
- Martens-Habbena W, Berube PM, Urakawa H, de la Torre JR, Stahl DA. 2009. Ammonia oxidation kinetics determine niche separation of nitrifying Archaea and Bacteria. *Nature* 461:976–979. <https://doi.org/10.1038/nature08465>.
- Urakawa H, Garcia JC, Nielsen JL, Le VQ, Kozłowski JA, Stein LY, Lim CK, Pommerening-Roser A, Martens-Habbena W, Stahl DA, Klotz MG. 2015. *Nitrosospira lacus* sp. nov., a psychrotolerant, ammonia-oxidizing bacterium from sandy lake sediment. *Int J Syst Evol Microbiol* 65:242–250. <https://doi.org/10.1099/ijs.0.070789-0>.
- Purkhold U, Pommerening-Roser A, Juretschko S, Schmid MC, Koops HP, Wagner M. 2000. Phylogeny of all recognized species of ammonia oxidizers based on comparative 16S rRNA and *amoA* sequence analysis: implications for molecular diversity surveys. *Appl Environ Microbiol* 66:5368–5382. <https://doi.org/10.1128/AEM.66.12.5368-5382.2000>.
- Urakawa H, Martens-Habbena W, Stahl DA. 2010. High abundance of

- ammonia-oxidizing *Archaea* in coastal waters, determined using a modified DNA extraction method. *Appl Environ Microbiol* 76:2129–2135. <https://doi.org/10.1128/AEM.02692-09>.
8. Gurevich A, Saveliev V, Vyahhi N, Tesler G. 2013. QUAST: quality assessment tool for genome assemblies. *Bioinformatics* 29:1072–1075. <https://doi.org/10.1093/bioinformatics/btt086>.
 9. Wick RR, Judd LM, Gorrie CL, Holt KE. 2017. Unicycler: resolving bacterial genome assemblies from short and long sequencing reads. *PLoS Comput Biol* 13:e1005595. <https://doi.org/10.1371/journal.pcbi.1005595>.
 10. Afgan E, Baker D, Batut B, van den Beek M, Bouvier D, Čech M, Chilton J, Clements D, Coraor N, Grüning BA, Guerler A, Hillman-Jackson J, Hiltemann S, Jalili V, Rasche H, Soranzo N, Goecks J, Taylor J, Nekrutenko A, Blankenberg D. 2018. The Galaxy platform for accessible, reproducible and collaborative biomedical analyses: 2018 update. *Nucleic Acids Res* 46:W537–W544. <https://doi.org/10.1093/nar/gky379>.
 11. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75. <https://doi.org/10.1186/1471-2164-9-75>.
 12. Overbeek R, Begley T, Butler RM, Choudhuri JV, Chuang H-Y, Cohoon M, de Crécy-Lagard V, Diaz N, Disz T, Edwards R, Fonstein M, Frank ED, Gerdes S, Glass EM, Goesmann A, Hanson A, Iwata-Reuyl D, Jensen R, Jamshidi N, Krause L, Kubal M, Larsen N, Linke B, McHardy AC, Meyer F, Neuweger H, Olsen G, Olson R, Osterman A, Portnoy V, Pusch GD, Rodionov DA, Rückert C, Steiner J, Stevens R, Thiele I, Vassieva O, Ye Y, Zagnitko O, Vonstein V. 2005. The subsystems approach to genome annotation and its use in the project to annotate 1000 genomes. *Nucleic Acids Res* 33:5691–5702. <https://doi.org/10.1093/nar/gki866>.
 13. Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>.
 14. Goris J, Konstantinidis KT, Klappenbach JA, Coenye T, Vandamme P, Tiedje JM. 2007. DNA-DNA hybridization values and their relationship to whole-genome sequence similarities. *Int J Syst Evol Microbiol* 57:81–91. <https://doi.org/10.1099/ijs.0.64483-0>.
 15. Kozłowski JA, Kits KD, Stein LY. 2016. Complete genome sequence of *Nitrosomonas ureae* strain Nm10, an oligotrophic group 6a nitrosomonad. *Genome Announc* 4:e00094-16. <https://doi.org/10.1128/genomeA.00094-16>.
 16. Yuichi S, Norton JM, Bollmann A, Klotz MG, Stein LY, Laanbroek HJ, Arp DJ, Goodwin LA, Chertkov O, Held B, Bruce D, Detter JC, Detter JC, Tapia R, Han CS. 2011. Genome sequence of *Nitrosomonas* sp. strain AL212, an ammonia-oxidizing bacterium sensitive to high levels of ammonia. *J Bacteriol* 193:5047–5048. <https://doi.org/10.1128/JB.05521-11>.
 17. Bollmann A, Sedlacek CJ, Norton J, Laanbroek HJ, Suwa Y, Stein LY, Klotz MG, Arp D, Sayavedra-Soto L, Lu M. 2013. Complete genome sequence of *Nitrosomonas* sp. Is79, an ammonia oxidizing bacterium adapted to low ammonium concentrations. *Stand Genomic Sci* 7:469–482.
 18. Konstantinidis KT, Tiedje JM. 2005. Genomic insights that advance the species definition for prokaryotes. *Proc Natl Acad Sci U S A* 102:2567–2572. <https://doi.org/10.1073/pnas.0409727102>.
 19. Kvint K, Nachin L, Diez A, Nyström T. 2003. The bacterial universal stress protein: function and regulation. *Curr Opin Microbiol* 6:140–145. [https://doi.org/10.1016/S1369-5274\(03\)00025-0](https://doi.org/10.1016/S1369-5274(03)00025-0).
 20. McLachlan A. 1983. Sandy beach ecology—a review, p 321–380. *In* McLachlan A, Erasmus T (ed), *Sandy beaches as ecosystems*. Springer, Dordrecht, the Netherlands.