



# Complete Genome Sequence of *Luteibacter pinisoli* MAH-14

David A. Baltrus,<sup>a,b</sup> Meara Clark,<sup>a</sup> Patrik Inderbitzin,<sup>c</sup> Daniela Pignatta,<sup>c</sup> Victoria Knight-Connoni,<sup>c</sup> A. Elizabeth Arnold<sup>a,d</sup>

<sup>a</sup>School of Plant Sciences, University of Arizona, Tucson, Arizona, USA

<sup>b</sup>School of Animal and Comparative Biomedical Sciences, University of Arizona, Tucson, Arizona, USA

<sup>c</sup>Indigo Agriculture, Charlestown, Massachusetts, USA

<sup>d</sup>Department of Ecology and Evolutionary Biology, University of Arizona, Tucson, Arizona, USA

**ABSTRACT** Diverse strains of *Luteibacter* (*Gammaproteobacteria*) have been isolated from a variety of environments, most frequently in association with both plants and fungi. Motivated by the lack of genomic information for strains throughout the genus *Luteibacter*, we report here a complete genome sequence for *Luteibacter pinisoli* strain MAH-14.

Strains classified within the genus *Luteibacter* (*Gammaproteobacteria*) are yellow-pigmented members of microbiomes found in close association with plants (1–4). A subset of strains have also been characterized as endohyphal bacteria within a diverse set of ascomycete fungi (5–8). These bacterial/fungal complexes modulate diverse fungal phenotypes, including the potential for some fungi to act as plant growth promoters (5–8). To increase the amount of genomic information for strains in this genus, we have sequenced a species of *Luteibacter* originally isolated from rhizosphere soil of *Pinus koraiensis* (2). Here, we report the complete genome sequence of *Luteibacter pinisoli* strain MAH-14.

An isolate of strain MAH-14 (KACC 19298) was acquired from the Korean Agricultural Culture Collection (KACC) and streaked to single colonies onto lysogeny broth (LB) agar plates. Plates were incubated at 27°C for 3 days until single colonies were apparent. A single colony was transferred to LB liquid medium and frozen in glycerol as part of the Baltrus lab culture collection (DBL1436). A sample of this frozen stock was independently streaked onto LB agar plates for each DNA extraction. Single colonies were then picked and transferred to 2 ml of LB broth and grown overnight at 27°C in a shaking incubator at 220 rpm. Genomic DNA used for Illumina sequencing was isolated from a 2-ml overnight culture via the Wizard kit (Promega, Madison, WI) following the manufacturer's protocols. Genomic DNA used for Nanopore sequencing was isolated from a separate 2-ml overnight culture via a Nanobind kit (Circulomics, Baltimore, MD) following the manufacturer's protocols. RNase was added per the manufacturer's protocols for each of the genomic isolations.

Genomic DNA was sequenced by SNPsaurus (Eugene, OR) using an Illumina HiSeq 4000 instrument and following their standard workflow for library preparation and read trimming. This workflow uses a Nextera tagmentation kit for library generation, followed by sequencing that generated 150-bp paired-end reads, followed by trimming of adaptors from reads with the computational suite BBDuk (BBMap version 38.41) (9). This workflow generated a total of 2,707,288 trimmed paired reads and 817.6 Mbp of sequence (~177× coverage).

A separate isolation of genomic DNA was sequenced by the Baltrus lab via an Oxford Nanopore MinION device with 1 μg of DNA prepared using the LSK-109 kit without shearing. Reads were called during sequencing using Guppy version 2.0.10, and only the reads which passed the initial quality check (QC) during the run (and which were

**Citation** Baltrus DA, Clark M, Inderbitzin P, Pignatta D, Knight-Connoni V, Arnold AE. 2019. Complete genome sequence of *Luteibacter pinisoli* MAH-14. *Microbiol Resour Announc* 8:e00774-19. <https://doi.org/10.1128/MRA.00774-19>.

**Editor** Julie C. Dunning Hotopp, University of Maryland School of Medicine

**Copyright** © 2019 Baltrus et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to David A. Baltrus, [baltrus@email.arizona.edu](mailto:baltrus@email.arizona.edu).

**Received** 28 June 2019

**Accepted** 1 July 2019

**Published** 18 July 2019

therefore placed into the “fast5\_pass” folder) were used. Sequencing on the MinION device generated 4,000 reads for a total of 42.8 Mbp of sequence (~10× coverage).

Hybrid assembly of all read types was performed using Unicycler version 0.4.8 (10) and resulted in a single chromosome of 4,610,686 bp of sequence with a 66.1% GC content. Unicycler also specified that this chromosomal assembly was circular. This chromosomal sequence was annotated by the NCBI Prokaryotic Genome Annotation Pipeline (PGAP) version 4.8 (11), and it is predicted to contain 4,175 genes representing 4,081 predicted protein coding sequences, 2 complete sets of rRNAs (5S, 16S, and 23S rRNAs), 51 tRNAs, and 4 noncoding RNAs (ncRNAs). Default parameters were used for all software.

**Data availability.** This genome project is indexed at GenBank under BioProject accession number [PRJNA548583](https://www.ncbi.nlm.nih.gov/bioproject/PRJNA548583). The complete genome sequence for *Luteibacter pinisoli* can be found under GenBank accession number [CP041046](https://www.ncbi.nlm.nih.gov/genbank/CP041046), and the reported genome is the first version, CP041046.1. Trimmed Illumina reads can be found under SRA run accession number [SRR9305630](https://www.ncbi.nlm.nih.gov/sra/SRR9305630). Fast5 files from the MinION run, which passed the initial QC during the MinION run and were therefore placed into the “fast5\_pass” folder, can be found under SRA run accession number [SRR9305631](https://www.ncbi.nlm.nih.gov/sra/SRR9305631). A log file generated by Unicycler for an assembly of this genome can be found at figshare (<https://doi.org/10.6084/m9.figshare.8330105.v1>).

## ACKNOWLEDGMENTS

We thank Nick Loman, Jennifer Gardy, Josh Quick, Jared Simpson, Matt Loose, John Tyson, Lauren Cowley, and the whole PoreCamp Vancouver crew for providing a great environment to learn about Nanopore sequencing.

Indigo Agriculture provided funds to D.A.B. for sequencing of this genome.

## REFERENCES

- Johansen JE, Binnerup SJ, Kroer N, Mølbak L. 2005. *Luteibacter rhizovicinus* gen. nov., sp. nov., a yellow-pigmented gammaproteobacterium isolated from the rhizosphere of barley (*Hordeum vulgare* L.). *Int J Syst Evol Microbiol* 55:2285–2291. <https://doi.org/10.1099/ijs.0.63497-0>.
- Akter S, Huq MA. 2018. *Luteibacter pinisoli* sp. nov., a casein degrading bacterium isolated from rhizospheric soil of *Pinus koraiensis*. *Arch Microbiol* 200:1017–1023. <https://doi.org/10.1007/s00203-018-1515-1>.
- Lasa AV, Mašínová T, Baldrian P, Fernández-López M. 2019. Bacteria from the endosphere and rhizosphere of *Quercus* spp. use mainly cell wall-associated enzymes to decompose organic matter. *PLoS One* 14: e0214422. <https://doi.org/10.1371/journal.pone.0214422>.
- Aamot HU, Hofgaard IS, Lysøe E. 2017. Complete genome sequence of *Luteibacter rhizovicinus* strain LJ96T, isolated from the rhizosphere of barley (*Hordeum vulgare* L.) in Denmark. *Genomics Data* 11:104. <https://doi.org/10.1016/j.gdata.2016.12.012>.
- Arendt KR, Hockett KL, Araldi-Brondolo SJ, Baltrus DA, Arnold AE. 2016. Isolation of endohyphal bacteria from foliar Ascomycota and *in vitro* establishment of their symbiotic associations. *Appl Environ Microbiol* 82:2943–2949. <https://doi.org/10.1128/AEM.00452-16>.
- Baltrus DA, Dougherty K, Arendt KR, Huntemann M, Clum A, Pillay M, Palaniappan K, Varghese N, Mikhailova N, Stamatis D, Reddy TBK, Ngan CY, Daum C, Shapiro N, Markowitz V, Ivanova N, Kyrpides N, Woyke T, Arnold AE. 2017. Absence of genome reduction in diverse, facultative endohyphal bacteria. *Microb Genom* 3:e000101. <https://doi.org/10.1099/mgen.0.000101>.
- Hoffman MT, Gunatilaka MK, Wijeratne K, Gunatilaka L, Elizabeth Arnold A. 2013. Endohyphal bacterium enhances production of indole-3-acetic acid by a foliar fungal endophyte. *PLoS One* 8:e73132. <https://doi.org/10.1371/journal.pone.0073132>.
- Araldi-Brondolo SJ, Spraker J, Shaffer JP, Woytenko EH, Baltrus DA, Gallery RE, Arnold AE. 2017. Bacterial endosymbionts: master modulators of fungal phenotypes. *Microbiol Spectr* 5. <https://doi.org/10.1128/microbiolspec.FUNK-0056-2016>.
- Bushnell B. BBMap. <https://sourceforge.net/projects/bbmap/>.
- Wick RR, Judd LM, Gorrie CL, Holt KE. 2017. Unicycler: resolving bacterial genome assemblies from short and long sequencing reads. *PLoS Comput Biol* 13:e1005595. <https://doi.org/10.1371/journal.pcbi.1005595>.
- Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>.