



Draft Genome Sequence of *Massilia* sp. Strain MC02, Isolated from a Sandy Loam Maize Soil

Rachel Raths,^a Vincent Peta,^a Heike Bücking^a

^aBiology and Microbiology Department, South Dakota State University, Brookings, South Dakota, USA

ABSTRACT From farmed corn soil in California, we isolated and sequenced a new member of the genus *Massilia*, *Massilia* sp. strain MC02. *Massilia* sp. MC02 has an assembled draft genome of 5,023,356 bp with a total of 4,790 protein-encoding genes and 3,028 predicted proteins, 47 tRNA genes, and 2 rRNA operons.

The first species within the genus *Massilia* was isolated from clinical samples (1). Since then, *Massilia* species have been isolated from plant tissues (2), water (3), air (4), ice cores (5), and soils (6). *Massilia* spp. have been shown to be abundant in the plant rhizosphere and to colonize roots (7, 8). Some species have plant growth-promoting capabilities, such as the production of indole-3-acetic acid (9) or siderophores, and are involved in soil carbon and nitrogen cycling (10). *Massilia* is the most species-rich genus of the *Oxalobacteraceae* family and consists mainly of Gram-negative, aerobic, non-spore-forming, motile rods.

Massilia sp. strain MC02 was isolated from a maize rhizosphere sample from a sandy loam soil in California on 22 May 2015. The geographical coordinates are 37.6058, -120.7478. Soil was added to phosphate-buffered saline, and dilutions were plated on Reasoner's 2A (R2A) agar plates and incubated at 30°C for 1 to 2 days, followed by 20°C for 1 to 2 days. Sequential colony streaks were performed on R2A agar to acquire pure colonies. Genomic DNA was extracted from a freshly grown R2A broth culture using the AllPrep bacterial DNA/RNA/protein kit (Qiagen, Inc., Germantown, MD) following the kit protocol. The genomic library was prepared with Illumina Nextera technology (San Diego, CA), size selected to an average fragment length of 475 bp, and sequenced using Illumina NextSeq paired-end v2 chemistry on v2.5 flow cells at 150 bp per read. A target coverage of 20× was used, and the genome was assembled using SPAdes v3.11.0 (11). Default parameters were used for all software unless otherwise specified. We obtained 1,566,408 total reads, with an average read length of 148 bp. The total read length was 231,096,882 bp, with 275 contigs, an N_{50} value of 28,267 bp (range, 1,074 to 119,695 bp), and an L_{50} value of 49. The genome length was 5,023,356 bp, with a GC content of 66.2%. Assembly quality assessment using BUSCO (12) revealed a measured completeness of 95%. Gene prediction and annotation using PATRIC v3.5.27 (13) resulted in a total of 4,790 protein-coding sequences consisting of 1,762 hypothetical proteins and 3,028 proteins with functional assignments, 47 tRNA genes, and 2 rRNA operons. The Microbial Genomes Atlas (MiGA) (14) revealed that the closest related strain from the NCBI database is *Massilia armeniaca* ZMN-3 (GenBank accession number [NZ_CP028324](https://doi.org/10.1128/MRA.00410-19)), with an amino acid identity (AAI) of 67.37%. Based on the MiGA results, MC02 belongs to the *Massilia* genus and was designated *Massilia* sp. strain MC02.

Using Galaxy (15), several genes with putative plant growth-promoting characteristics were identified, such as a nitrate reductase gene (*napA*), several phosphatase genes (*ppk*, *phoA*, *phoB*, *phoD*, and *phoR*), and biotin biosynthesis genes (*bioA*, *bioB*, *bioD*, and *bioF*). Using RAST 2.0 (16), we identified 48 putative virulence genes, including 34 genes indicating a resistance to antibiotics and toxic compounds, 14

Citation Raths R, Peta V, Bücking H. 2019. Draft genome sequence of *Massilia* sp. strain MC02, isolated from a sandy loam maize soil. Microbiol Resour Announc 8:e00410-19. <https://doi.org/10.1128/MRA.00410-19>.

Editor David A. Baltrus, University of Arizona

Copyright © 2019 Raths et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Heike Bücking, heike.buecking@sdsu.edu.

R.R. and V.P. contributed equally to this work.

Received 5 April 2019

Accepted 14 July 2019

Published 8 August 2019

genes putatively involved in invasion and intracellular resistance, and 26 genes responsible for flagellar motility. Invasion and flagellum genes are essential for attaching and entering plant cells (17), suggesting that MCO2 is a plant endophyte.

Data availability. The complete genome sequence has been deposited in NCBI/EBI/GenBank under BioProject number [PRJNA529270](https://www.ncbi.nlm.nih.gov/bioproject/PRJNA529270), BioSample number [SAMN11263498](https://www.ncbi.nlm.nih.gov/biosample/SAMN11263498), GenBank accession number [SPVF00000000](https://www.ncbi.nlm.nih.gov/nuccore/SPVF00000000), and SRA accession number [SRX6098478](https://www.ncbi.nlm.nih.gov/sra/SRX6098478).

ACKNOWLEDGMENTS

This project was funded by Novozymes North America.

We thank Novozymes North America for providing the novel isolate and for whole-genome sequencing and Timothy Lilburn (Novozymes North America), Deborah Springer (Novozymes North America), and Alex Soupir (South Dakota State University) for technical guidance and support throughout this project. We also acknowledge support by the SDSU Agricultural Experiment Station.

REFERENCES

- Scola BL, Birtles RJ, Mallet M-N, Raoult D. 1998. *Massilia timonae* gen. nov., sp. nov., isolated from blood of an immunocompromised patient with cerebellar lesions. *J Clin Microbiol* 36:2847–2852.
- Chimwamurombe PM, Grönemeyer JL, Reinhold-Hurek B. 2016. Isolation and characterization of culturable seed-associated bacterial endophytes from gnotobiotically grown Marama bean seedlings. *FEMS Microbiol Ecol* 92:11.
- Lu JY, Tian Z, Yu JW, Yang M, Zhang Y. 2018. Distribution and abundance of antibiotic resistance genes in sand settling reservoirs and drinking water treatment plants across the Yellow River, China. *Water* 10:12. <https://doi.org/10.3390/w10030246>.
- Orthova I, Kämpfer P, Glaeser SP, Kaden R, Busse HJ. 2015. *Massilia norwichensis* sp. nov., isolated from an air sample. *Int J Syst Evol Microbiol* 65:56–64. <https://doi.org/10.1099/ijvs.0.068296-0>.
- Guo BX, Liu YQ, Gu ZQ, Shen L, Liu KS, Wang NL, Xing TT, Liu HC, Zhou YG, Li JL. 2016. *Massilia psychrophila* sp. nov., isolated from an ice core. *Int J Syst Evol Microbiol* 66:4088–4093. <https://doi.org/10.1099/ijsem.0.001315>.
- Ren MN, Li XY, Zhang YQ, Jin Y, Li SQ, Huang HD. 2018. *Massilia armeniaca* sp. nov., isolated from desert soil. *Int J Syst Evol Microbiol* 68:2319–2324. <https://doi.org/10.1099/ijsem.0.002836>.
- Maya O, Yitzhak H, Dror M. 2012. Ecology of root colonizing *Massilia* (*Oxalobacteraceae*). *PLoS One* 7:e40117. <https://doi.org/10.1371/journal.pone.0040117>.
- Li X, Rui J, Mao Y, Yannarell A, Mackie R. 2014. Dynamics of the bacterial community structure in the rhizosphere of a maize cultivar. *Soil Biol Biochem* 68:392–401. <https://doi.org/10.1016/j.soilbio.2013.10.017>.
- Kuffner M, De Maria S, Puschenreiter M, Fallmann K, Wieshammer G, Gorfer M, Strauss J, Rivelli AR, Sessitsch A. 2010. Culturable bacteria from Zn- and Cd-accumulating *Salix caprea* with differential effects on plant growth and heavy metal availability. *J Appl Microbiol* 108:1471–1484. <https://doi.org/10.1111/j.1365-2672.2010.04670.x>.
- Hryniewicz K, Baum C, Leinweber P. 2010. Density, metabolic activity, and identity of cultivable rhizosphere bacteria on *Salix viminalis* in disturbed arable and landfill soils. *J Plant Nutr Soil Sci* 173:747–756. <https://doi.org/10.1002/jpln.200900286>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
- Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. 2015. BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics* 31:3210–3212. <https://doi.org/10.1093/bioinformatics/btv351>.
- Wattam AR, Davis JJ, Assaf R, Boisvert S, Brettin T, Bun C, Conrad N, Dietrich EM, Disz T, Gabbard JL, Gerdes S, Henry CS, Kenyon RW, Machi D, Mao C, Nordberg EK, Olsen GJ, Murphy-Olson DE, Olson R, Overbeek R, Parrello B, Pusch GD, Shukla M, Vonstein V, Warren A, Xia FF, Yoo H, Stevens RL. 2017. Improvements to PATRIC, the all-bacterial bioinformatics database and analysis resource center. *Nucleic Acids Res* 45:D535–D542. <https://doi.org/10.1093/nar/gkw1017>.
- Rodriguez R. 2018. The Microbial Genomes Atlas (MiGA) webserver: taxonomic and gene diversity analysis of archaea and bacteria at the whole genome level. *Nucleic Acids Res* 46:282–288. <https://doi.org/10.1093/nar/gky467>.
- Afgan E, Baker D, Batut B, van den Beek M, Bouvier D, Čech M, Chilton J, Clements D, Coraor N, Grüning BA, Guerler A, Hillman-Jackson J, Hiltmann S, Jalili V, Rasche H, Soranzo N, Goecks J, Taylor J, Nekrutenko A, Blankenberg D. 2018. The Galaxy platform for accessible, reproducible and collaborative biomedical analyses: 2018 update. *Nucleic Acids Res* 46:W537–W544. <https://doi.org/10.1093/nar/gky379>.
- Overbeek R, Begley T, Butler RM, Choudhuri JV, Chuang HY, Cohoon M, de Crecy-Lagard V, Diaz N, Disz T, Edwards R, Fonstein M, Frank ED, Gerdes S, Glass EM, Goessmann A, Hanson A, Iwata-Reuyl D, Jensen R, Jamshidi N, Krause L, Kubal M, Larsen N, Linke B, McHardy AC, Meyer F, Neuweger H, Olsen G, Olson R, Osterman A, Portnoy V, Pusch GD, Rodionov DA, Ruckert C, Steiner J, Stevens R, Thiele I, Vassieva O, Ye Y, Zagnitko O, Vonstein V. 2005. The subsystems approach to genome annotation and its use in the project to annotate 1000 genomes. *Nucleic Acids Res* 33:5691–5702. <https://doi.org/10.1093/nar/gki866>.
- Czaban J, Gajda A, Wróblewska B. 2007. The motility of bacteria from rhizosphere and different zones of winter wheat roots. *Polish J Environ Stud* 16:301–308.