



Elucidation of the Genome of *Bradyrhizobium* sp. Strain USDA 3456, a Historic Agricultural Diazotroph from Cowpea (*Vigna unguiculata*)

Richard Allen White III,^{a,b,c} Jeffrey S. Norman,^d Emily E. Mclachlan,^a Joseph P. Dunham,^e Aaron Garoutte,^d Maren L. Friesen^{a,d,f}

^aDepartment of Plant Pathology, Washington State University, Pullman, Washington, USA

^bRAW Molecular Systems (RMS), LLC, Spokane, Washington, USA

^cAustralian Centre for Astrobiology, University of New South Wales, Sydney, NSW, Australia

^dDepartment of Plant Biology, Michigan State University, East Lansing, Michigan, USA

^eSeqOnce Biosciences, Inc., Pasadena, California, USA

^fDepartment of Crop and Soil Sciences, Washington State University, Pullman, Washington, USA

ABSTRACT *Bradyrhizobium* sp. strain USDA 3456 is a historic strain from the United States Department of Agriculture (USDA) Agricultural Research Service (ARS) National *Rhizobium* Germplasm Collection isolated from *Vigna unguiculata* (cowpea) in 1966. Strain USDA 3456 has been utilized in global agricultural applications, including improving soil nitrogen fertility. The draft genome sequence here provides a genetic reference of a novel diazotroph.

The rhizosphere microbiome is one of the most dynamic interfaces on Earth (1–3), with diazotrophs fixing biologically required nitrogen (4, 5). Some diazotrophs can perform symbiotic nitrogen fixation (SNF) by forming root nodules with compatible plant hosts typically in the family Fabaceae, where SNF is selected for by the plant, resulting in fitness alignment (4, 5), or they can fix nitrogen while free-living in soil, aquatic, or other habitats (6).

Bradyrhizobium sp. strain USDA 3456 was isolated from a *Vigna unguiculata* (cowpea) nodule from Wisconsin in 1966 (7). Field trials with USDA 3456 in peanut (*Arachis hypogaea*) suggest that it possesses high phosphorus solubilization and moderate indole acetic acid (IAA) production (7, 8). Inoculation of USDA 3456 in peanut crops over two 75-day field seasons resulted in nitrogenase activity of 18 to 19 $\mu\text{mol C}_2\text{H}_4/\text{g}^{-1}$ (nodule dry weight)/h (8). USDA 3456 can be coinoculated with other plant growth-promoting bacteria, resulting in a synergistic effect on crop production (8).

A lyophilized culture of *Bradyrhizobium* strain USDA 3456 was obtained from the National Rhizobium Germplasm Collection (NRGC) (<https://data.nal.usda.gov/dataset/usda-ars-national-rhizobium-germplasm-collection>). A single colony was inoculated in AG broth culture (25 ml) at 30°C at 200 rpm to obtain biomass for DNA extraction (9).

DNA was extracted using the MasterPure DNA extraction kit (Epicentre, Madison, WI, USA) following the manufacturer's protocols. A SeqOnce RhinoSeq kit was used to prepare libraries (<https://seqonce.com/rhinoseq/>). Libraries were quantified and sequenced on a HiSeq 4000 instrument in a 150-bp paired-end read format at the Michigan State University Research Technology Support Facility (RTSF).

Default parameters were used for all software unless otherwise specified, but versions of software are provided. Illumina sequencing data were quality filtered and decontaminated using ATLAS (version 1.0) (10). The resulting cleaned reads from ATLAS

Citation White RA, III, Norman JS, Mclachlan EE, Dunham JP, Garoutte A, Friesen ML. 2019. Elucidation of the genome of *Bradyrhizobium* sp. strain USDA 3456, a historic agricultural diazotroph from cowpea (*Vigna unguiculata*). *Microbiol Resour Announc* 8:e00812-19. <https://doi.org/10.1128/MRA.00812-19>.

Editor J. Cameron Thrash, University of Southern California

Copyright © 2019 White et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Maren L. Friesen, m.friesen@wsu.edu.

Received 14 July 2019

Accepted 29 July 2019

Published 15 August 2019

(1,815,290 paired-end reads) were then assembled with Unicycler (version 0.4.7) using default Illumina assembly parameters (11).

The Unicycler final genome assembly for strain USDA 3456 is 107 contigs, with a genome size of 9,771,557 bp, a G+C content of 63.57%, and an N_{50} value of 324,457 bp. We estimated completeness and contamination of the draft genome using CheckM (version 1.0.12); the genome was 100% complete with 0% contamination (12). Eight contigs that were <200 bp were removed for submission to GenBank, including the version discussed here. All assemblies are publicly available at Open Science Framework (OSF) for the genome. Prokka (version 1.13.3) with the -rfam flag annotated the genome to obtain rRNAs and transfer-messenger RNAs (tmRNAs) (13). The annotation from Prokka predicted 49 tRNAs, 1 tmRNA, 30 noncoding RNAs (misc_RNAs), 1 copy of 5S-16S-23S operons, 0 CRISPRs, and 9,142 protein-coding genes.

Bradyrhizobium sp. strain USDA 3456 is a versatile diazotroph that has been used in agricultural applications for greater than half a century (7). We provide this high-quality draft genome sequence as a template for further metabolic engineering and synthetic biology applications for sustainable agriculture (14, 15).

Data availability. This whole-genome shotgun project has been deposited at DDBJ/ENA/GenBank under the accession number [VIDU00000000](https://doi.org/10.1093/bioinformatics/btu153). The version described in this paper is the first version, VIDU01000000. Raw data, contigs, and annotations for this genome can be found at OSF (<https://osf.io/5tx6c/>), and code used to generate the assembly can be found at www.github.com/friesenlab/Bradyrhizobium_USDA3456.

ACKNOWLEDGMENTS

We acknowledge support from NSF IOS 1331218 to M.L.F. and NSF DEB 1547024 to M.L.F. and J.S.N.

REFERENCES

- White RA, III, Rivas-Ubach A, Borkum MI, Köberl M, Bilbao A, Colby SM, Hoyt DW, Bingol K, Kim YM, Wendler JP, Hixson KK, Jansson C. 2017. The state of rhizospheric science in the era of multi-omics: a practical guide to omics technologies. *Rhizosphere* 3:212–221. <https://doi.org/10.1016/j.rhisph.2017.05.003>.
- Berendsen RL, Pieterse CM, Bakker PA. 2012. The rhizosphere microbiome and plant health. *Trends Plant Sci* 17:478–486. <https://doi.org/10.1016/j.tplants.2012.04.001>.
- White IIR, Borkum MI, Rivas-Ubach A, Bilbao A, Wendler JP, Colby SM, Köberl M, Jansson C. 2017. From data to knowledge: the future of multi-omics data analysis for the rhizosphere. *Rhizosphere* 3:222–229. <https://doi.org/10.1016/j.rhisph.2017.05.001>.
- Masson-Boivin C, Giraud E, Perret X, Batut J. 2009. Establishing nitrogen-fixing symbiosis with legumes: how many rhizobium recipes? *Trends Microbiol* 17:458–466. <https://doi.org/10.1016/j.tim.2009.07.004>.
- Friesen ML. 2012. Widespread fitness alignment in the legume–rhizobium symbiosis. *New Phytol* 194:1096–1111. <https://doi.org/10.1111/j.1469-8137.2012.04099.x>.
- Smercina DN, Evans SE, Friesen ML, Tiemann LK. 2019. To fix or not to fix: controls on free-living nitrogen fixation in the rhizosphere. *Appl Environ Microbiol* 85:e02546-18. <https://doi.org/10.1128/AEM.02546-18>.
- van Berkum P, Elia P, Song Q, Eardly BD. 2012. Development and application of a multilocus sequence analysis method for the identification of genotypes within genus *Bradyrhizobium* and for establishing nodule occupancy of soybean (*Glycine max* L. Merr). *Mol Plant Microbe Interact* 25:321–330. <https://doi.org/10.1094/MPMI-09-11-0241>.
- Badawi F, Biomy AMM, Desoky AH. 2011. Peanut plant growth and yield as influenced by co-inoculation with *Bradyrhizobium* and some rhizobium microorganisms under sandy loam soil conditions. *Ann Agric Sci* 56:17–25. <https://doi.org/10.1016/j.aosas.2011.05.005>.
- Sadowsky MJ, Tully RE, Cregan PB, Keyser HH. 1987. Genetic diversity in *Bradyrhizobium japonicum* serogroup 123 and its relation to genotype-specific nodulation. *Appl Environ Microbiol* 53:2624–2630.
- White III RA, Brown J, Colby S, Overall CC, Lee J, Zucker J, Glaesemann KR, Jansson C, Jansson JK. 2017. ATLAS (Automatic Tool for Local Assembly Structures)—a comprehensive infrastructure for assembly, annotation, and genomic binning of metagenomic and metatranscriptomic data. *PeerJ Preprints* 5:e2843v1. <https://peerj.com/preprints/2843/>.
- Wick RR, Judd LM, Gorrie CL, Holt KE. 2017. Unicycler: resolving bacterial genome assemblies from short and long sequencing reads. *PLoS Comput Biol* 13:e1005595. <https://doi.org/10.1371/journal.pcbi.1005595>.
- Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. 2015. CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res* 25:1043–1055. <https://doi.org/10.1101/gr.186072.114>.
- Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068–2069. <https://doi.org/10.1093/bioinformatics/btu153>.
- Ahkami AH, White IIR, Handakumbura PP, Jansson C. 2017. Rhizosphere engineering: enhancing sustainable plant ecosystem productivity. *Rhizosphere* 3:233–243. <https://doi.org/10.1016/j.rhisph.2017.04.012>.
- Busby PE, Soman C, Wagner MR, Friesen ML, Kremer J, Bennett A, Morsy M, Eisen JA, Leach JE, Dangl JL. 2017. Research priorities for harnessing plant microbiomes in sustainable agriculture. *PLoS Biol* 15:e2001793. <https://doi.org/10.1371/journal.pbio.2001793>.