



First Detection of a Novel Reassortant Avian Influenza A(H5N6) Clade 2.3.2.1c Virus, Isolated from a Wild Bird in China

Tao Zhang,^a  Ruiyun Li,^b Gaodong Zhu,^c Jianyu Chang,^d Bing Xu^a

^aMinistry of Education Key Laboratory for Earth System Modeling, Department of Earth System Science, Tsinghua University, Beijing, China

^bMRC Centre for Global Infectious Disease Analysis, Department of Infectious Disease Epidemiology, School of Public Health, Faculty of Medicine, Imperial College London, London, United Kingdom

^cSuichuan Forestry Bureau, Jiangxi, China

^dCollege of Veterinary Medicine, China Agricultural University, Beijing, China

ABSTRACT We report the first isolation of a reassortant clade 2.3.2.1c avian influenza A(H5N6) virus isolated from a wild bird sample in Jiangxi, China, in 2016. Sequence analyses indicated that this virus most likely evolved from Eurasia-derived H5N1 and H6N6 viruses through frequent interactions at the wild-domestic bird interface.

Avian influenza virus (AIV) is a single-stranded segmented negative-sense RNA virus classified in the *Alphainfluenzavirus* genus in the family *Orthomyxoviridae* (1). The hemagglutinin (HA) gene of highly pathogenic avian influenza (HPAI) A(H5) viruses has undergone continuous evolution, generating emerging reassorted subtypes and clades (2). Among these variants, two relatively new clades, i.e., 2.3.4.4 and 2.3.2.1c, predominantly shaped the current viral evolution and transcontinental expansions (3, 4). It is remarkable that clade 2.3.4.4 involves multiple subtypes, including H5N1, H5N2, H5N6, and H5N8 (2). In contrast, all clade 2.3.2.1c viruses, including the novel Sanmenxia clade 2.3.2.1c-like H5N1 virus isolated in a wild bird, belong to H5N1 (5).

Here, we provide a report on the first detection and nearly complete genome sequence of a novel reassortant clade 2.3.2.1c H5N6 sample isolated from a wild bird in China. A total of 488 tracheal and cloacal swab samples were collected during a routine bird survey in Suichuan County, Jiangxi Province, which were preserved in a sample solution in the fridge (4°C) and subsequently shipped to a laboratory and stored frozen at –80°C. From these samples, 89 and 85 samples were taken from *Streptopelia decaocto* and yellow-legged button quail, respectively. Virus isolation using these specimens was conducted in 9- to 11-day-old specific-pathogen-free embryonated

TABLE 1 Nucleotide sequencing identities between the novel clade 2.3.2.1c H5N6 virus and nearest homologs in the GenBank database

Gene	Virus	Accession no.	Subtype	Identity (%)
HA	A/bar-headed goose/China/F/2015	MK641386	H5N1	99.21
	A/bar-headed goose/China/70/2015	MK641378	H5N1	99.21
	A/bar-headed goose/China/133/2015	MK641362	H5N1	99.21
	A/whooper swan/Shanxi/17L/2015	KP715064	H5N1	99.21
NA	A/chicken/Jiangxi/NCDZT1123/2014	KP090441	H5N6	99.37
PB2	A/chicken/Jiangsu/2477/2014	KP762511	H5N1	99.53
PB1	A/pigeon/Zhejiang/112090/2014	KU042689	H5N1	99.82
PA	A/chicken/Jiangxi/NCDZT1123/2014	KP090438	H5N6	99.24
NP	A/chicken/Jiangxi/NCDZT1123/2014	KP090440	H5N6	99.30
M	A/duck/Guangzhou/021/2014	KX094409	H5N6	99.51
NS	A/chicken/Jiangsu/2477/2014	KP762516	H5N1	99.77

Citation Zhang T, Li R, Zhu G, Chang J, Xu B. 2019. First detection of a novel reassortant avian influenza A(H5N6) clade 2.3.2.1c virus, isolated from a wild bird in China. *Microbiol Resour Announc* 8:e00797-19. <https://doi.org/10.1128/MRA.00797-19>.

Editor Simon Roux, DOE Joint Genome Institute

Copyright © 2019 Zhang et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Jianyu Chang, changjianyu@cau.edu.cn, or Bing Xu, bingxu@tsinghua.edu.cn.

T.Z. and R.L. contributed equally to this work.

Received 2 July 2019

Accepted 25 July 2019

Published 5 September 2019




FIG 1 Phylogenetic analysis of Jiangxi clade 2.3.2.1c H5N6 virus. (A) Phylogeny of the HA gene was inferred using the maximum likelihood method with 1,000 bootstrap replicates. Sequence alignment and the inference of phylogeny were conducted using MEGA v6.06. Clades were classified in accordance with the published clade designation. The Jiangxi clade 2.3.2.1c H5N6 virus and clade 2.3.2.1 viruses are highlighted by a dot and colored rectangle, respectively. (B) Phylogenetic relationship of the divergent clade 2.3.2.1 viruses. (C to I) Phylogeny of the NA (C), PB2 (D), PB1 (E), PA (F), NP (G), M (H), and NS (I) genes.

chicken eggs. The viral RNAs were extracted from allantoic fluid of 13 samples with hemagglutination activity using an RNeasy minikit (Qiagen, Hilden, Germany). The SuperScript III reverse transcriptase (RT) PCR kit (Invitrogen, USA) was used for the reverse transcription.

The subtype of each of the 13 positive samples was first determined using PCR of a marker gene (6, 7). Seven out of 13 samples were identified as H5N6 strains. All segments of the H5N6 strains were amplified by using a Phusion high-fidelity PCR system (New England BioLabs, Ipswich, MA, USA), adhering to the manufacturer's guide (8). Sequencing of each segment was subsequently performed as individual amplicons using the Applied Biosystems automated 3730xl DNA analyzer. Among seven H5N6 strains, one named *A/Streptopelia decaocto/Jiangxi/G6/2016* (H5N6) was a novel reassortant strain. The coding region of each segment of this strain is base pairs 1 to 1776 (HA), base pairs 1 to 1431 (NA), base pairs 1 to 2341 (PB2), base pairs 1 to 2274 (PB1), base pairs 1 to 2233 (PA), base pairs 1 to 1565 (NP), base pairs 1 to 1027 (M), and base pairs 1 to 875 (NS). The GC contents were calculated using DNASTar v7.1.0 and were

Downloaded from <http://mra.asm.org/> on September 16, 2019 by guest

40.82% (HA), 43.16% (NA), 44.86% (PB2), 4.14% (PB1), 43.93% (PA), 47.35% (NP), 47.03% (M), and 44.34% (NS).

The amino acid sequence at the HA cleavage site is RERRRKR/GL, which is characteristic of high pathogenicity in poultry. Additionally, HA had Q222 and G224 (H3 numbering) at the receptor binding site associated with an adaptation to avian-like receptors (9). NA had an 11-amino-acid deletion (positions 58 to 68) in the stalk region, which may be associated with viral adaptation to terrestrial poultry after being introduced from water birds (10). However, there were no mutations for E627K and D701N in PB2 or at positions 26, 27, 30, 31, and 34 in the M2 protein, indicating an inefficiency to replicate in mammals (11) and sensitivity to amantadine (12).

A BLAST search in the GenBank database showed that all the eight genes of Jiangxi H5N6 virus had high nucleotide identities with viruses isolated from terrestrial poultry in China 2014 to 2015 (Table 1). These high nucleotide identities of viruses isolated from wild birds and terrestrial poultry are suggestive of frequent interactions and viral circulation between two species. Specifically, the HA gene shared 99.12% nucleotide identity with the HA gene from four H5N1 viruses. NA, PA, and NP were closely related to those of A/chicken/Jiangxi/NCDZT1123/2014 (H5N6), with identities of 99.37%, 99.24%, and 99.30%, respectively. The PB2 and NS genes shared the closest nucleotide similarity (>99.50%) with the PB2 and NS genes of A/chicken/Jiangsu/2477/2014 (H5N1). For the PB1 and M genes, the highest nucleotide identities were with those of A/pigeon/Zhejiang/112090/2014 (99.85%) and A/duck/Guangzhou/021/2014 (99.51%), respectively. Notably, phylogenetic analysis revealed that this Jiangxi H5N6 virus was a novel reassortant strain, in which the HA gene was found to belong to clade 2.3.2.1c (Fig. 1).

This detection suggests continuous reassortment and generation of emerging variants. Further investigation and active surveillance are required to detect new AIV variants.

Data availability. The genome sequence of Jiangxi clade 2.3.2.1c H5N6 virus has been deposited in GenBank with the accession numbers [MN165550](#) to [MN165557](#).

ACKNOWLEDGMENTS

This work was supported by the National Research Program of the Ministry of Science and Technology of the People's Republic of China (grant 2016YFA0600104) and by donations from Delos Living LLC and the Cyrus Tang Foundation to Tsinghua University.

We sincerely acknowledge the authors and submitting laboratories of the sequences deposited in the GenBank database, on which the phylogenetic analysis of this research is based.

We do not have any financial or other relationships that may pose conflicts of interest.

REFERENCES

- Lefkowitz EJ, Dempsey DM, Hendrickson RC, Orton RJ, Siddell SG, Smith DB. 2018. Virus taxonomy: the database of the International Committee on Taxonomy of Viruses (ICTV). *Nucleic Acids Res* 46:D708–D717. <https://doi.org/10.1093/nar/gkx932>.
- Smith GJ, Donis RO, World Health Organization/World Organisation for Animal Health/Food and Agriculture Organization (WHO/OIE/FAO) H5 Evolution Working Group. 2015. Nomenclature updates resulting from the evolution of avian influenza A(H5) virus clades 2.1.3.2a, 2.2.1, and 2.3.4 during 2013–2014. *Influenza Other Respir Viruses* 9:271–276. <https://doi.org/10.1111/irv.12324>.
- Global Consortium for H5N8 and Related Influenza Viruses. 2016. Role for migratory wild birds in the global spread of avian influenza H5N8. *Science* 14:213–217. <https://doi.org/10.1126/science.aaf8852>.
- Naguib MM, Kinne J, Chen H, Chan KH, Joseph S, Wong PC, Wernery R, Beer M, Wernery U, Harder TC. 2015. Outbreaks of highly pathogenic avian influenza H5N1 clade 2.3.2.1c in hunting falcons and kept wild birds in Dubai implicate intercontinental virus spread. *J Gen Virol* 96:3212–3222. <https://doi.org/10.1099/jgv.0.000274>.
- Bi Y, Chen J, Zhang Z, Li M, Cai T, Sharshov K, Susloparov I, Shestopalov A, Wong G, He Y, Xing Z, Sun J, Liu D, Liu Y, Liu L, Liu W, Lei F, Shi W, Gao GF. 2016. Highly pathogenic avian influenza H5N1 clade 2.3.2.1c virus in migratory birds, 2014–2015. *Virol Sin* 31:300–305. <https://doi.org/10.1007/s12250-016-3750-4>.
- Tsukamoto K, Ashizawa T, Nakanishi K, Kaji N, Suzuki K, Shishido M, Okamatsu M, Mase M. 2009. Use of reverse transcriptase PCR to subtype N1 to N9 neuraminidase genes of avian influenza viruses. *J Clin Microbiol* 47:2301–2303. <https://doi.org/10.1128/JCM.02366-08>.
- Lee MS, Chang PC, Shien JH, Cheng MC, Shieh HK. 2001. Identification and subtyping of avian influenza viruses by reverse transcription-PCR. *J Virol Methods* 97:13–22. [https://doi.org/10.1016/S0166-0934\(01\)00301-9](https://doi.org/10.1016/S0166-0934(01)00301-9).
- Hoffmann E, Stech J, Guan Y, Webster RG, Perez DR. 2001. Universal primer set for the full-length amplification of all influenza A viruses. *Arch Virol* 146:2275–2289. <https://doi.org/10.1007/s007050170002>.
- Guan Y, Poon LLM, Cheung CY, Ellis TM, Lim W, Lipatov AS, Chan KH, Sturm-Ramirez KM, Cheung CL, Leung YHC, Yuen KY, Webster RG, Peiris

- J. 2004. H5N1 influenza: a protean pandemic threat. *Proc Natl Acad Sci U S A* 101:8156–8161. <https://doi.org/10.1073/pnas.0402443101>.
10. Zhou H, Yu Z, Hu Y, Tu J, Zou W, Peng Y, Zhu J, Li Y, Zhang A, Yu Z, Ye Z, Chen H, Jin M. 2009. The special neuraminidase stalk-motif responsible for increased virulence and pathogenesis of H5N1 influenza A virus. *PLoS One* 4:e6277. <https://doi.org/10.1371/journal.pone.0006277>.
11. Li Z, Chen H, Jiao P, Deng G, Tian G, Li Y, Hoffmann E, Webster RG, Matsuoka Y, Yu K. 2005. Molecular basis of replication of duck H5N1 influenza viruses in a mammalian mouse model. *J Virol* 79:12058–12064. <https://doi.org/10.1128/JVI.79.18.12058-12064.2005>.
12. Furuse Y, Suzuki A, Oshitani H. 2009. Large-scale sequence analysis of M gene of influenza A viruses from different species: mechanisms for emergence and spread of amantadine resistance. *Antimicrob Agents Chemother* 53:4457–4463. <https://doi.org/10.1128/AAC.00650-09>.