




Genome Sequence of *Pseudomonas* sp. Strain ST1, Isolated from Olive (*Olea europaea* L.) Knot Galls in Croatia

Gabriela Vuletin Selak,^a Marina Raboteg,^a Pascale Fournier,^b Audrey Dubost,^b Danis Abrouk,^b Katja Žanić,^a Slavko Perica,^a Philippe Normand,^b  Petar Pujić^b

^aInstitute for Adriatic Crops and Karst Reclamation, Split, Croatia

^bEcologie Microbienne, Centre National de la Recherche Scientifique UMR 5557, Université de Lyon, Université Claude Bernard Lyon I, INRA, UMR1418, Villeurbanne, France

ABSTRACT We report the genome sequence of a *Pseudomonas* sp. strain isolated from olive knot galls. The genome size is 6.101 Mbp with a G+C content of 58%. A total of 6,137 coding DNA sequences (CDS) were predicted, including 52 tRNA and 4 rRNA genes.

Olive knot is one of the most important diseases of the olive crop and is present in all olive-growing regions. *Pseudomonas savastanoi* pv. *savastanoi* is the causal agent of olive knot disease, which results in tumorous overgrowths (knots). This olive knot pathogen bacterium can survive and multiply on aerial plant surfaces as well as in knots. The earlier studies of *Pseudomonas savastanoi* pv. *savastanoi* virulence implicated the type III secretion system, phytohormones, and quorum sensing (QS) as being involved in the disease process (1, 2). The pathogen can be dispersed both within the plant and to surrounding plants in wind-blown rain, by insects, and by human activities, entering the plant through wounds. Populations of *P. savastanoi* are normally associated with nonpathogenic bacteria, both epiphytically and endophytically (1, 3, 4). More specifically, the disease progression and knot volume were increased by coinoculation of *Pseudomonas savastanoi* pv. *savastanoi* with nonpathogenic bacteria (1, 4, 5).

Pseudomonas cells were isolated from olive knots from olive plants grown in the central region of Dalmatia, Croatia (43°30'19.6"N, 16°29'55.0"E). Olive knots were harvested from a plant trunk using a sterile scalpel, immediately surface sterilized using 75% ethanol, and sliced using a sterile scalpel, and the slices were placed on the surface of King's medium agar plates (6). Plates were kept in the dark at 25°C for 48 hours. Cells from a single fluorescent colony identified under UV light on an agar plate were transferred into 10 ml of liquid lysogeny broth (LB) medium in a 50-ml Falcon tube and grown at 28°C for 24 hours with shaking. Genomic DNA was isolated from bacterial cells using a microbial DNA kit (reference number 740235; Macherey-Nagel, Hoerd, France). The 16S rRNA gene was amplified with PCR using 20 ng genomic DNA, 0.2 mM deoxynucleoside triphosphates (dNTPs), 50 nM each com1 (5'CAGCAGCCGCGTAA TAC) and com2 (5'CCGTCAATTCCTTGAGTTT) primers, and 2.5 U *Taq* polymerase (Invitrogen, France). The amplified DNA product was sequenced using Sanger sequencing at Biofidal, Vaulx-en-Velin, France. Sequence comparison using a BLASTN search with the NCBI database showed that the strain belongs to a group of the genus *Pseudomonas*. The *Pseudomonas* sp. strain ST1 genome bank was made using a Nextera XT DNA library prep kit and protocol (Illumina, Évry, France) and sequenced using Illumina MiSeq technology with a paired-end 2 × 300-bp run (Biofidal). Quality controls were made with FastQC (7) and Trimmomatic (8). We obtained a total of 8,334,104 reads with 416× coverage, an N_{50} value of 0.082 Mbp, and a G+C content of 58%. Default parameters were used for Unicycler assembly with a minimum contig length of 200 bp. Genome assembly was performed using Unicycler version 0.4.3, and annotation was

Citation Vuletin Selak G, Raboteg M, Fournier P, Dubost A, Abrouk D, Žanić K, Perica S, Normand P, Pujić P. 2019. Genome sequence of *Pseudomonas* sp. strain ST1, isolated from olive (*Olea europaea* L.) knot galls in Croatia. *Microbiol Resour Announc* 8:e00986-19. <https://doi.org/10.1128/MRA.00986-19>.

Editor David A. Baltrus, University of Arizona

Copyright © 2019 Vuletin Selak et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Gabriela Vuletin Selak, gabriela.vuletin.selak@krs.hr, or Petar Pujić, petar.pujic@univ-lyon1.fr.

Received 13 August 2019

Accepted 20 October 2019

Published 14 November 2019

done with the MicroScope platform version 3.10.0 (9, 10) using the Rapid Annotations using Subsystems Technology (RAST) (11) and PATRIC (12) Web servers and Prokka software (13). The genome of *Pseudomonas* sp. strain ST1 has 6,070,031 bp assembled in 318 contigs. The genome has 6,019 predicted genes. The public version of the *Pseudomonas* sp. strain ST1 genome sequence at GenBank was annotated using the Prokaryotic Genome Annotation Pipeline (PGAP) (14). The version described in this paper is the first version.

Data availability. The complete genome sequence described here has been deposited in NCBI/GenBank under BioProject number [PRJNA555035](https://ncbi.nlm.nih.gov/bioproject/PRJNA555035), BioSample number [SAMN12289196](https://ncbi.nlm.nih.gov/biosample/SAMN12289196), accession number [VKOF00000000](https://ncbi.nlm.nih.gov/nuccore/VKOF00000000), and SRA number [SRX6799169](https://ncbi.nlm.nih.gov/sra/SRX6799169).

ACKNOWLEDGMENTS

This work was supported by the Split-Dalmatia County grant for the project “Agents, transmitters and bio-control of olive knot disease in the olive groves of the Split-Dalmatia County” and by the Unity Through Knowledge Fund, collaboration grant 2017 (contract number 10/17) within the Research Cooperability Program. M.R. is a Ph.D. fellow supported by a French government scholarship in 2019/2020 announced by the Ministry of Science and Education of Republic of Croatia, Agency for Mobility and EU Programmes and Embassy of the French Republic to the Republic of Croatia. The LABGeM (CEA/Genoscope and CNRS UMR8030), the France Génomique, and French Bioinformatics Institute National Infrastructures (funded as part of an Investissement d’Avenir program managed by the Agence Nationale pour la Recherche, contracts ANR-10-INBS-09 and ANR-11-INBS-0013) are acknowledged for support within the MicroScope annotation platform.

We thank Corinne Sannaire and platform Genomique Environnementale PGE, UMR CNRS 5557 for technical support.

We declare no conflict of interest.

REFERENCES

- Hosni T, Moretti C, Devescovi G, Suarez-Moreno ZR, Fatmi MB, Guarnaccia C, Pongor S, Onofri A, Buonauro R, Venturi V. 2011. Sharing of quorum-sensing signals and role of interspecies communities in a bacterial plant disease. *ISME J* 5:1857–1870. <https://doi.org/10.1038/ismej.2011.65>.
- Pérez-Martínez I, Rodríguez-Moreno L, Lambertsen L, Matas IM, Murillo J, Tegli S, Jiménez AJ, Ramos C. 2010. Fate of a *Pseudomonas savastanoi* pv. *savastanoi* type III secretion system mutant in olive plants (*Olea europaea* L.). *Appl Environ Microbiol* 76:3611–3619. <https://doi.org/10.1128/AEM.00133-10>.
- Vuletin Selak G, Raboteg M, Dubost A, Abrouk D, Žanić K, Normand P, Pujic P. 2019. Whole-genome sequence of a *Pantoea* sp. strain isolated from an olive (*Olea europaea* L.) knot. *Microbiol Resour Announc* 8:e00978-19. <https://doi.org/10.1128/MRA.00978-19>.
- Marchi G, Sisto A, Cimmino A, Andolfi A, Cipriani MG, Evidente A, Surico G. 2006. Interaction between *Pseudomonas savastanoi* pv. *savastanoi* and *Pantoea agglomerans* in olive knots. *Plant Pathol* 55:614–624. <https://doi.org/10.1111/j.1365-3059.2006.01449.x>.
- Passos da Silva D, Castaneda-Ojeda MP, Moretti C, Buonauro R, Ramos C, Venturi V. 2014. Bacterial multispecies studies and microbiome analysis of a plant disease. *Microbiology* 160:556–566. <https://doi.org/10.1099/mic.0.074468-0>.
- King EO, Ward MK, Raney DE. 1954. Two simple media for the demonstration of pyocyanin and fluorescin. *J Lab Clin Med* 44:301–307.
- Andrews S. 2010. FastQC: a quality control tool for high throughput sequencing data. <http://www.bioinformatics.babraham.ac.uk/projects/fastqc>.
- Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
- Wick RR, Judd LM, Gorrie CL, Holt KE. 2017. Unicycler: resolving bacterial genome assemblies from short and long sequencing reads. *PLoS Comput Biol* 13:e1005595. <https://doi.org/10.1371/journal.pcbi.1005595>.
- Vallenet D, Calteau A, Cruveiller S, Gachet M, Lajus A, Josso A, Mercier J, Renaux A, Rollin J, Rouy Z, Roche D, Scarpelli C, Médigue C. 2017. MicroScope in 2017: an expanding and evolving integrated resource for community expertise of microbial genomes. *Nucleic Acids Res* 45:D517–D528. <https://doi.org/10.1093/nar/gkw1101>.
- Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75. <https://doi.org/10.1186/1471-2164-9-75>.
- Wattam AR, Abraham D, Dalay O, Disz TL, Driscoll T, Gabbard JL, Gillespie JJ, Gough R, Hix D, Kenyon R, Machi D, Mao C, Nordberg EK, Olson R, Overbeek R, Pusch GD, Shukla M, Schulman J, Stevens RL, Sullivan DE, Vonstein V, Warren A, Will R, Wilson MJC, Yoo HS, Zhang C, Zhang Y, Sobral BW. 2014. PATRIC, the bacterial bioinformatics database and analysis resource. *Nucleic Acids Res* 42:D581–D591. <https://doi.org/10.1093/nar/gkt1099>.
- Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068–2069. <https://doi.org/10.1093/bioinformatics/btu153>.
- Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>.