



First Complete Genome Sequence of *Cucurbit Aphid-Borne Yellows Virus* from Pumpkin in the United States

Vivek Khanal,^a Akhtar Ali^a

^aDepartment of Biological Science, The University of Tulsa, Tulsa, Oklahoma, USA

ABSTRACT *Cucurbit aphid-borne yellows virus* (CABYV) was first described in France in 1992 and since then has been reported in various parts of the world, including the United States. Here, we present the first complete genome sequence of a CABYV isolate (BL4) that was collected from pumpkin during the 2017 growing season in Oklahoma.

Cucurbits are economically important cash crops in the United States and are mostly grown in the southern states, including Texas and Florida, which are the two major watermelon-producing states (1). CABYV belongs to the genus *Polevirus* in the family *Luteoviridae* (2) and was first reported in France in 1992 (3). The virus causes yellowing and thickening of the older leaves in cucurbit plants and is often mistakenly attributed as a nutrient deficiency. Although the major veins of younger leaves would remain green after the infection, plant yield may be reduced (3). The virus is transmitted primarily by *Aphis gossypii* Glover and *Myzus persicae* Sulzer, and the transmission could be circulative, persistent, and nonpropagative (4, 5). CABYV has been reported from cucurbit crops across different climatic regions of the world such as temperate, Mediterranean, and subtropical (6), and no mechanical transmission has been reported (7). The main constraint for the management of diseases caused by members of *Luteoviridae* is that no effective strategy exists to cure plants after virus infection (8).

It has been nearly two and half decades since the first report of CABYV in the United States (9); however, to our knowledge, no complete genome sequence of any CABYV isolate from the United States has been reported so far. In this work, we report the first complete genome sequence of a CABYV isolate collected from a grower's field in Oklahoma.

Previously, we reported CABYV for the first time from commercial cucurbit fields in Blaine County in Oklahoma (10). One of the dot-immunobinding assay (DIBA)-positive samples (10) against the CABYV antibody (designated as CABYV isolate BL4) was used in this work.

Total RNA was extracted from the CABYV-infected leaf tissues of pumpkin (11). Seven pairs of overlapping primers were designed (Table 1) and synthesized commercially (IDT Technologies, USA) from the previous CABYV isolates available from GenBank. All seven genome fragments were amplified by reverse transcription-PCR (RT-PCR) with the respective primer pairs using total RNA as the template, as described previously (11). Both 5' end and 3' end rapid amplification of cDNA ends (RACE) was performed using a commercial kit (TaKaRa Bio, Inc., Japan). Expected PCR products were analyzed and confirmed on 1% agarose gels and cleaned with Exosap-IT (Affymetrix). Purified PCR products were directly sequenced in both directions using an Applied Biosystems 3130 instrument.

Nucleotide sequences of each genome fragment were confirmed and assembled using DNASTAR Lasergene 13 (Madison, WI) software. Sequences of all fragments were aligned using both Clustal W v2.1 (12) and Muscle v3.8.31 (13) alignment tools

Citation Khanal V, Ali A. 2019. First complete genome sequence of *Cucurbit aphid-borne yellows virus* from pumpkin in the United States. *Microbiol Resour Announc* 8:e01448-18. <https://doi.org/10.1128/MRA.01448-18>.

Editor Jelle Matthijnsens, KU Leuven

Copyright © 2019 Khanal and Ali. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Akhtar Ali, akhtar-ali@utulsa.edu.

Received 7 November 2018

Accepted 10 January 2019

Published 7 February 2019

TABLE 1 Primers used in RT-PCR to amplify the complete genome of *Cucurbit aphid-borne yellow virus*

Primer name	Primer sequence	Product size (nucleotides)
CABYVF1	ACAAAAGATACGAGCGGGTGA	665
CABYVR1	GCAAGTCGCCAAAAATCCAA	
CABYV F2	AGAGAGTCTGCTCCACGTGA	1,099
CABYV R2	GAGACTGTGCGTCTCCACTT	
CABYV F3	CGACCGCCGAAACAACCG	1,168
CABYVR3	TGCTCATCCGTAGACAAGCC	
CABYVF4	AACAAGCGCGAGATAGCGTT	998
CABYVR4	CGCCTCCCTGCATTTTGATT	
CABYVCP5	ATGAATACGGCCGCGCTAGAAATC	600
CABYVCP5	CTATTTCCGGTTCTGGACCTGGCA	
CABYVF6	AACGGATCTTCTCGGTTGC	1,320
CABYVR6	CACCTTCTCGTCTTCTCGTC	
CABYV F7	CACAGCCCACCCGGACTATA	544
CABYVR7	ACACCGAAACGCCAGGGG	

by joining overlapping sequences to generate the complete genome sequence of CABYV.

After the alignment, the complete genome sequence of the CABYV BL4 isolate is 5,679 nucleotides long (G+C content, 49.8%). Nucleotide BLAST searches showed that the CABYV BL4 isolate shares 85% to 97% nucleotide and 81% to 96% amino acid sequence identities with the available CABYV complete genomes in the GenBank database. The highest nucleotide (97%) and amino acid (96%) sequence similarities were observed with a Chinese isolate (GenBank accession no. [EU000535](https://doi.org/10.1093/phyto/14.10.1400)) (14).

Data availability. The complete genome sequence of the CABYV BL4 isolate was deposited in GenBank under accession no. [MK055337](https://doi.org/10.1093/phyto/14.10.1400).

ACKNOWLEDGMENTS

This publication was supported by the U.S. Department of Agriculture (USDA) Agricultural Marketing Service through grant no. 15SCBGPOK0019. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the USDA.

REFERENCES

- Ali A, Abdalla O, Bruton B, Fish W, Sikora E, Zhang S, Taylor M. 2012. Occurrence of viruses infecting watermelon, other cucurbits, and weeds in the parts of southern United States. *Plant Health Prog* <https://doi.org/10.1094/PHP-2012-0824-01-RS>.
- King A, Adams M, Carstens E, Lefkowitz E. 2012. Virus taxonomy: ninth report of the International Committee on Taxonomy of Viruses. Academic Press, London, United Kingdom.
- Lecoq H, Bourdin D, Wipf-Scheibel C, Bon M, Lot H, Lemaire O, Herrbach E. 1992. A new yellowing disease of cucurbits caused by a *Luteovirus*, *Cucurbit aphid-borne yellows virus*. *Plant Pathol* 41:749–761. <https://doi.org/10.1111/j.1365-3059.1992.tb02559.x>.
- Dogimont C, Slama S, Martin J, Lecoq H, Pitrat M. 1996. Sources of resistance to *Cucurbit aphid-borne yellows Luteovirus* in a melon germ plasm collection. *Plant Dis* 80:1379–1382. <https://doi.org/10.1094/PD-80-1379>.
- Gray S, Cilia M, Ghanim M. 2014. Circulative, “nonpropagative” virus transmission: an orchestra of virus-, insect-, and plant-derived instruments. *Adv Virus Res* 89:141–199. <https://doi.org/10.1016/B978-0-12-800172-1.00004-5>.
- Lecoq H. 1999. Epidemiology of Cucurbit aphid-borne yellows virus, p 243–248. In Smith HG, Baker H (ed). *The Luteoviridae*. CABI Publishing, Wallingford, United Kingdom.
- D’Arcy CJ, Domier LL. 2005. *Luteoviridae*, p 891–900. In Fauquet CM, Mayo MA, Maniloff J, Desselberger U, Ball LA (ed). *Virus taxonomy*, VIIIth report of the ICTV. Elsevier/Academic Press, London, United Kingdom.
- Michelle H, Veronique B. 2018. Targeted disruption of aphid transmission: a vision for the management of crop diseases caused by *Luteoviridae* members. *Curr Opin Virol* 33:24–32. <https://doi.org/10.1016/j.coviro.2018.07.007>.
- Lemaire OJ, Gubler WD, Valencia J, Lecoq H, Falk BW. 1993. First report of *Cucurbit aphid borne yellows Luteovirus* in the United States. *Plant Dis* 77:1169. <https://doi.org/10.1094/PD-77-1169B>.
- Khanal V, Ali A. 2018. First report of *Cucurbit aphid-borne yellows virus* infecting *Cucurbita pepo* in Oklahoma. *Plant Dis* 102:1046. <https://doi.org/10.1094/PDIS-10-17-1675-PDN>.
- Ali A, Mohammad O, Khattab A. 2012. Distribution of viruses infecting cucurbit crops and isolation of potential new virus-like sequences from weeds in Oklahoma. *Plant Dis* 96:243–248. <https://doi.org/10.1094/PDIS-05-11-0419>.
- Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, McWilliam H, Remmert M, Söding J, Thompson JD, Higgins DG. 2014. Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. *Mol Syst Biol* 7:539. <https://doi.org/10.1038/msb.2011.75>.
- Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res* 32:1792–1797. <https://doi.org/10.1093/nar/gkh340>.
- Xiang HY, Shang QX, Han CG, Li DW, Yu JL. 2008. Complete sequence analysis reveals two distinct *Poleroviruses* infecting cucurbits in China. *Arch Virol* 153:1155–1160. <https://doi.org/10.1007/s00705-008-0083-0>.