



Complete Genome Sequences of *Mycobacterium tuberculosis* Isolates Subjected to 200 Continuous Passages

 Lucio Vera-Cabrera,^a Carmen A. Molina-Torres,^a Jose Marco Lopez-Ortiz,^a Jorge Castro-Garza,^b Carlos Cordova-Fletes,^c Jorge Ocampo-Candiani^a

^aLaboratorio Interdisciplinario de Investigación Dermatológica, Servicio de Dermatología, Hospital Universitario “Dr. José Eleuterio González,” Universidad Autónoma de Nuevo León, Monterrey, NL, México

^bCentro de Investigación Biomédica del Noreste, Instituto Mexicano del Seguro Social, Monterrey, NL, México

^cDepartamento de Bioquímica, Facultad de Medicina, Universidad Autónoma de Nuevo León, Monterrey, NL, México

ABSTRACT We report 6 draft genome sequences corresponding to *Mycobacterium tuberculosis* H37Rv and *M. tuberculosis* DR689, a Beijing isolate, plus their counterparts subjected to 200 continuous passages in Middlebrook 7H9 broth, either alone or with ox bile.

Tuberculosis is still a public health problem worldwide (1). In Mexico, the number of new pulmonary and meningeal tuberculosis cases has increased in the last several years (2). New drugs and immunoprophylactic media are needed to control this disease. Considering that continuous passaging produces important biological changes in microorganisms (3, 4), we decided to subculture *Mycobacterium tuberculosis* isolates. Although *M. tuberculosis* and *M. bovis* belong to the same bacterial complex, they have important biological differences (5).

In our project, we subcultured isolates of different lineages of *M. tuberculosis* (6), including MTBH37Rv and a Beijing isolate, DR689. The only isolate that has already been sequenced is strain H37Rv, which constitutes the reference genome sequence used worldwide (GenBank accession number [NC_000962.3](https://www.ncbi.nlm.nih.gov/nuccore/NC_000962.3)). This strain was donated to J.C.-G. in 1998 by Fred Quinn, when he was a research microbiologist at the Tuberculosis and Mycobacteriology Branch, Division of AIDS, STD, and TB Laboratory Research, Centers for Disease Control and Prevention (Atlanta, GA). It was cryopreserved at -70°C until we started the subculture. Both strains MTBH37Rv and MTBDR689 were subjected to 200 continuous passages on M7H9-Tween 80-oleic acid-albumin-dextrose-catalase (OADC); because Calmette and Guérin used ox bile to better suspend the bacteria, we processed some samples with and some without bile. We use the following nomenclature to refer to our cultures: *M. tuberculosis* H37Rv parental strain (MTBH37RvP), without bile (MTBH37Rv-P200) and with bile (MTBH37Rv-P200B), and *M. tuberculosis* DR689 parental strain (MTBDR689P) without bile (MTBDR689-P200) and with bile (MTBDR689P-200B).

DNA was extracted from the colonies grown on Lowenstein-Jensen agar using the cetyltrimethylammonium bromide (CTAB)-NaCl technique with modifications described previously (6). Subsequently, high-throughput sequencing was carried out using the Illumina MiSeq genome analyzer (San Diego, CA, USA). Sequencing libraries were prepared using the NuGEN Ultralow v2 kit or the Nextera XT v2 kit following the manufacturer's recommended protocol. The libraries were sequenced on a NextSeq 500 instrument using midoutput v2 chemistry (2×150 bp) or on a MiSeq instrument using V2 chemistry (2×250 bp); the fastq reads obtained were mapped against the *M. tuberculosis* H37Rv reference genome sequence reported in GenBank (accession number [NC_000962.3](https://www.ncbi.nlm.nih.gov/nuccore/NC_000962.3)) using the BWA-MEM aligner (7) from the Sequencher v5.4.6 suite

Citation Vera-Cabrera L, Molina-Torres CA, Lopez-Ortiz JM, Castro-Garza J, Cordova-Fletes C, Ocampo-Candiani J. 2020. Complete genome sequences of *Mycobacterium tuberculosis* isolates subjected to 200 continuous passages. *Microbiol Resour Announc* 9:e00356-20. <https://doi.org/10.1128/MRA.00356-20>.

Editor Christina A. Cuomo, Broad Institute

Copyright © 2020 Vera-Cabrera et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Lucio Vera-Cabrera, drluciovera@yahoo.com.

Received 9 April 2020

Accepted 2 June 2020

Published 25 June 2020

TABLE 1 Draft genome characteristics and accession numbers of *M. tuberculosis* H37RvP, *M. tuberculosis* DR689P, and their counterparts, subjected to 200 continuous passages with or without bile

Strain/isolate	GenBank accession no.	SRA accession no.	Genome size (Mbp)	G+C content (%)	No. of contigs	N ₅₀ value (bp)	No. of reads	Coverage (x)	No. of CDSs ^a	% coding proteins	No. of tRNAs	No. of rRNAs
<i>M. tuberculosis</i> H37RvP	VRNC000000000	SRX8374041	4.3	65.55	103	142,434	1,690,242	162	5,489	98.2	42	1
<i>M. tuberculosis</i> H37Rv-P200	VRND000000000	SRX8374042	4.3	65.55	155	64,842	7,851,072	162	4,378	97.1	45	1
<i>M. tuberculosis</i> H37Rv-P200B	VRNE000000000	SRX8374043	4.3	65.43	205	57,433	3,899,892	109.5	4,622	93.5	46	1
<i>M. tuberculosis</i> DR689P	VRNF000000000	SRX8374038	4.3	65.59	115	105,204	1,373,090	34	4,464	95.8	49	1
<i>M. tuberculosis</i> DR689-P200	VRNG000000000	SRX8374039	4.3	65.44	230	45,562	2,040,850	58	4,523	94.2	45	1
<i>M. tuberculosis</i> DR689-P200B	VRNH000000000	SRX8374040	4.3	65.56	160	64,269	5,448,990	81	4,359	96.4	47	1

^aCDSs, coding DNA sequences.

(Gene Codes, Ann Arbor, MI). Default parameters were used except where otherwise noted.

The genomes were assembled into contigs using SPAdes Genome Assembler v3.9.0 and deposited in GenBank. Annotations were made using the single Prokaryotic Genome Annotation Pipeline (PGAP) of the NCBI (8).

Data availability. This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under the following accession numbers: [VRNC000000000](https://www.ncbi.nlm.nih.gov/nuccore/VRNC000000000) (*Mycobacterium tuberculosis* H37RvP), [VRND000000000](https://www.ncbi.nlm.nih.gov/nuccore/VRND000000000) (*Mycobacterium tuberculosis* H37Rv-P200), [VRNE000000000](https://www.ncbi.nlm.nih.gov/nuccore/VRNE000000000) (*Mycobacterium tuberculosis* H37Rv-P200B), [VRNF000000000](https://www.ncbi.nlm.nih.gov/nuccore/VRNF000000000) (*Mycobacterium tuberculosis* DR689P), [VRNG000000000](https://www.ncbi.nlm.nih.gov/nuccore/VRNG000000000) (*Mycobacterium tuberculosis* DR689-P200), and [VRNH000000000](https://www.ncbi.nlm.nih.gov/nuccore/VRNH000000000) (*Mycobacterium tuberculosis* DR689-P200B). All these sequences belong to BioProject accession number [PRJNA559369](https://www.ncbi.nlm.nih.gov/bioproject/PRJNA559369). The characteristics of the draft genomes are summarized in Table 1.

REFERENCES

- World Health Organization. 2018. Global tuberculosis report 2018. World Health Organization. <https://apps.who.int/iris/handle/10665/274453>.
- Bello-López JM, León-García G, Rojas-Bernabé A, Fernández-Sánchez V, García-Hernández O, Mancilla Ramírez J, Ibáñez-Cervantes G. 2019. Morbidity trends and risk of tuberculosis: Mexico 2007–2017. *Can Respir J* 2019:8295261. <https://doi.org/10.1155/2019/8295261>.
- Moreira D, Santarém N, Loureiro I, Tavares J, Silva AM, Amorim AM, Ouaisi A, Cordeiro-da-Silva A, Silvestre R. 2012. Impact of continuous axenic cultivation in *Leishmania infantum* virulence. *PLoS Negl Trop Dis* 6:e1469. <https://doi.org/10.1371/journal.pntd.0001469>.
- Songe MM, Thoen E, Evensen Ø, Skaar I. 2014. In vitro passages impact on virulence of *Saprolegnia parasitica* to Atlantic salmon, *Salmo salar* L. parr. *J Fish Dis* 37:825–834. <https://doi.org/10.1111/jfd.12175>.
- Malone KM, Rue-Albrecht K, Magee DA, Conlon K, Schubert OT, Nalpas NC, Browne JA, Smyth A, Gormley E, Aebersold R, MacHugh DE, Gordon SV. 2018. Comparative 'omics analyses differentiate *Mycobacterium tuberculosis* and *Mycobacterium bovis* and reveal distinct macrophage responses to infection with the human and bovine tubercle bacilli. *Microb Genom* 4:e000163. <https://doi.org/10.1099/mgen.0.000163>.
- Molina-Torres CA, Castro-Garza J, Ocampo-Candiani J, Monot M, Cole ST, Vera-Cabrera L. 2010. Effect of serial subculturing on the genetic composition and cytotoxic activity of *Mycobacterium tuberculosis*. *J Med Microbiol* 59:384–391. <https://doi.org/10.1099/jmm.0.015966-0>.
- Li H. 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. arXiv 1303.3997 [q-bio.GN]. <http://arxiv.org/abs/1303.3997>.
- Haft DH, DiCuccio M, Badretdin A, Brover V, Chetvernin V, O'Neill K, Li W, Chitsaz F, Derbyshire MK, Gonzales NR, Gwadz M, Lu F, Marchler GH, Song JS, Thanki N, Yamashita RA, Zheng C, Thibaud-Nissen F, Geer LY, Marchler-Bauer A, Pruitt KD. 2018. RefSeq: an update on prokaryotic genome annotation and curation. *Nucleic Acids Res* 46:D851–D860. <https://doi.org/10.1093/nar/gkx1068>.