



Nearly Complete Genome Sequence of *Brugia pahangi* FR3

John Mattick,^a Silvia Libro,^b Benjamin C. Sparklin,^a  Matthew Chung,^a  Robin E. Bromley,^a Suvarna Nadendla,^a Xuechu Zhao,^a Sandra Ott,^a Lisa Sadzewicz,^a Luke J. Tallon,^a Michelle L. Michalski,^c  Jeremy M. Foster,^b  Julie C. Dunning Hotopp^{a,d,e}

^aInstitute for Genome Science, University of Maryland School of Medicine, Baltimore, Maryland, USA

^bDivision of Protein Expression and Modification, New England Biolabs, Ipswich, Massachusetts, USA

^cDepartment of Biology and Microbiology, University of Wisconsin Oshkosh, Oshkosh, Wisconsin, USA

^dDepartment of Microbiology and Immunology, University of Maryland School of Medicine, Baltimore, Maryland, USA

^eGreenebaum Cancer Center, University of Maryland School of Medicine, Baltimore, Maryland, USA

ABSTRACT *Brugia pahangi* is a zoonotic parasite that is closely related to human-infecting filarial nematodes. Here, we report the nearly complete genome of *Brugia pahangi*, including assemblies of four autosomes and an X chromosome, with only seven gaps. The Y chromosome is still not completely assembled.

We sequenced the zoonotic parasitic roundworm *Brugia pahangi* FR3, which was originally obtained from a green leaf monkey in Kuala Lumpur and is distributed by the NIAID Filariasis Research Reagent Resource Center, known as FR3 (1). The 107,643,863 Illumina HiSeq 2500 paired-end 150-bp reads were generated from KAPA Hyper libraries using *Brugia pahangi* FR3 genomic DNA acquired from BEI Resources. Genomic DNA for PacBio RS II sequencing was obtained from 16 adult male and 16 adult female frozen worms obtained from FR3, which were homogenized in Qiagen G2 buffer with RNase A, and purified using Qiagen gravity-flow Genomic-tips with 80 U of proteinase K and DNA precipitation by centrifugation in the presence of 20 μ g of glycogen. PacBio RS II data (P6C4 chemistry; 3,267,281 reads; average read length, 8,695 bp; N_{50} , 25.4 kbp; maximum read length, 139 kbp) were generated from Sage Blue Pippin size-selected (>15 kbp) SMRTbell v1.0 libraries constructed using Covaris gTUBE-fragmented DNA. For Oxford Nanopore Technologies (ONT) sequencing, 185 adult female worms acquired from TRS Labs (Athens, GA, USA) were ground in liquid nitrogen, and DNA was extracted with a single phenol-chloroform DNA extraction, with spooling from an ethanol precipitation. Rapid libraries were constructed (SQK-RAD004) four times, from 4 μ g, 1.8 μ g, 0.9 μ g, and 0.2 μ g DNA. The latter three libraries were sequenced with an R9.4 MinION flow cell (FLO-MIN106), replacing the loading beads with water. The 4- μ g library was modified using 1.5 μ l of DNA fragmentation mixture and 3.5 μ l of 10 mM Tris-Cl (pH 8.0) and 0.02% Triton X-100 (2) and was sequenced with a different R9.4 MinION flow cell, compared with the previous runs. This collectively resulted in 727,358 reads, which were called using Guppy v3.1.5 (average read length, 10,109 bp; N_{50} , 22 kbp; maximum read length, 828 kbp).

Illumina reads after quality control and trimming with FastQC v0.11.7 (3) and Trimmomatic v0.38 (4), respectively, and untrimmed PacBio and MinION reads were assembled with Canu v1.8 (5) multiple times but using different subsets of MinION experiments (excluding and including the 4- μ g sample) and varying the estimated genome size as 90 Mbp and 100 Mbp. Default parameters were used except as otherwise noted. These assemblies were manually examined, including unique joins leading to a manual merger of assemblies. This manually merged assembly was indel corrected using Illumina reads and Pilon v1.22 (mindepth, 5; K, 85; minmq, 0; minqual, 35; fix indels) (6). Contigs were ordered and oriented according to the *Brugia*

Citation Mattick J, Libro S, Sparklin BC, Chung M, Bromley RE, Nadendla S, Zhao X, Ott S, Sadzewicz L, Tallon LJ, Michalski ML, Foster JM, Dunning Hotopp JC. 2020. Nearly complete genome sequence of *Brugia pahangi* FR3. Microbiol Resour Announc 9:e00479-20. <https://doi.org/10.1128/MRA.00479-20>.

Editor Christina A. Cuomo, Broad Institute

Copyright © 2020 Mattick et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Julie C. Dunning Hotopp, jdhotopp@som.umaryland.edu.

Received 29 April 2020

Accepted 11 June 2020

Published 2 July 2020




FIG 1 Synteny between *Brugia pahangi* and *Brugia malayi* chromosomes. The current *Brugia pahangi* genome was aligned to the *Brugia malayi* V4 genome (7) using NUCmer v3.1 (default options). The resulting delta file was converted into tabular form using the show-coords function with the option -qITHb. Contigs >4.5% of the length of the matching *B. malayi* chromosome were identified in R (v. 3.4.4 [9]; libraries: RColorBrewer, ggplot2, data.table) and plotted with ggplots by chromosome. (A) Chromosome 1; (B) chromosome 2; (C) chromosome 3; (D) chromosome 4; (E) chromosome X. This nearly complete *B. pahangi* genome spans most of the *B. malayi* genome with some rearrangements. The orthologous regions of the genomes vary in percent identity but typically show >90% identity, with more nucleotide variation at the ends of chromosomes, relative to the middle of the same chromosomes, using the *B. malayi* chromosomes as references.

malayi reference genome (7), using NUCmer v3.1 to identify matches between the two genomes and to order *B. pahangi* contigs based on their *B. malayi* counterparts (Fig. 1). The genome spans 80.8 Mbp with seven gaps (N_{50} , 11.2 Mbp; L_{50} , 3), has a GC content of 29% with four autosomes and an X chromosome, and was deemed 96.4% complete with BUSCO v3 (8), with 206× Illumina, 182× PacBio, and 72× ONT sequencing depths. The remaining 142 contigs, spanning 16 Mbp, likely contain the mitochondrial genome, haplotypes, and pseudoautosomal and Y chromosome fragments.

Data availability. This genome has been deposited in GenBank under accession number [JAAVKF0100000000](https://www.ncbi.nlm.nih.gov/nuccore/JAAVKF0100000000). The version described in this paper is the first version, [JAAVKF0100000000](https://www.ncbi.nlm.nih.gov/nuccore/JAAVKF0100000000). The raw data have been deposited in the SRA under the accession numbers [SRX4135331](https://www.ncbi.nlm.nih.gov/sra/SRX4135331), [SRX4135330](https://www.ncbi.nlm.nih.gov/sra/SRX4135330), [SRX4135329](https://www.ncbi.nlm.nih.gov/sra/SRX4135329), [SRX4135328](https://www.ncbi.nlm.nih.gov/sra/SRX4135328), [SRX4135327](https://www.ncbi.nlm.nih.gov/sra/SRX4135327), [SRX4135326](https://www.ncbi.nlm.nih.gov/sra/SRX4135326), [SRX4135325](https://www.ncbi.nlm.nih.gov/sra/SRX4135325), [SRX4135324](https://www.ncbi.nlm.nih.gov/sra/SRX4135324), [SRX4135323](https://www.ncbi.nlm.nih.gov/sra/SRX4135323), [SRX4135322](https://www.ncbi.nlm.nih.gov/sra/SRX4135322), [SRX4135321](https://www.ncbi.nlm.nih.gov/sra/SRX4135321), [SRX4135320](https://www.ncbi.nlm.nih.gov/sra/SRX4135320), [SRX4135319](https://www.ncbi.nlm.nih.gov/sra/SRX4135319), [SRX4135318](https://www.ncbi.nlm.nih.gov/sra/SRX4135318), [SRX4135317](https://www.ncbi.nlm.nih.gov/sra/SRX4135317), [SRX4135316](https://www.ncbi.nlm.nih.gov/sra/SRX4135316), [SRX4135315](https://www.ncbi.nlm.nih.gov/sra/SRX4135315), [SRX4135314](https://www.ncbi.nlm.nih.gov/sra/SRX4135314), [SRX4135313](https://www.ncbi.nlm.nih.gov/sra/SRX4135313), and [SRX4135312](https://www.ncbi.nlm.nih.gov/sra/SRX4135312) (PacBio), [SRX7658407](https://www.ncbi.nlm.nih.gov/sra/SRX7658407), [SRX7658383](https://www.ncbi.nlm.nih.gov/sra/SRX7658383), [SRX7658378](https://www.ncbi.nlm.nih.gov/sra/SRX7658378), [SRX7658377](https://www.ncbi.nlm.nih.gov/sra/SRX7658377), [SRX7658352](https://www.ncbi.nlm.nih.gov/sra/SRX7658352), [SRX7658349](https://www.ncbi.nlm.nih.gov/sra/SRX7658349), [SRX7658341](https://www.ncbi.nlm.nih.gov/sra/SRX7658341), [SRX7658327](https://www.ncbi.nlm.nih.gov/sra/SRX7658327), [SRX7658323](https://www.ncbi.nlm.nih.gov/sra/SRX7658323), [SRX7658322](https://www.ncbi.nlm.nih.gov/sra/SRX7658322), [SRX7658317](https://www.ncbi.nlm.nih.gov/sra/SRX7658317), and [SRR10997235](https://www.ncbi.nlm.nih.gov/sra/SRR10997235) (Illumina), and [SRR11565851](https://www.ncbi.nlm.nih.gov/sra/SRR11565851), [SRR11472020](https://www.ncbi.nlm.nih.gov/sra/SRR11472020), [SRR11565849](https://www.ncbi.nlm.nih.gov/sra/SRR11565849), and [SRR11565826](https://www.ncbi.nlm.nih.gov/sra/SRR11565826) (MinION).

ACKNOWLEDGMENTS

This project was funded by federal funds from the National Institute of Allergy and Infectious Diseases, National Institutes of Health, under grant U19AI110820.

Genomic DNA from adult female *Brugia pahangi* FR3 (NR-46542) and L3-stage (NR-42499) worms, and whole adult male (NR-48906) and female (NR-48904) worms were provided by the NIH-NIAID Filariasis Research Reagent Resource Center for distribution by BEI Resources, NIAID, NIH.

REFERENCES

- Michalski ML, Griffiths KG, Williams SA, Kaplan RM, Moorhead AR. 2011. The NIH-NIAID Filariasis Research Reagent Resource Center. *PLoS Negl Trop Dis* 5:e1261. <https://doi.org/10.1371/journal.pntd.0001261>.
- Tyson JR, O'Neil NJ, Jain M, Olsen HE, Hieter P, Snutch TP. 2018. MinION-based long-read sequencing and assembly extends the *Caenorhabditis elegans* reference genome. *Genome Res* 28:266–274. <https://doi.org/10.1101/gr.221184.117>.
- Andrews S. 2013. FastQC. <https://www.bioinformatics.babraham.ac.uk/projects/fastqc>.
- Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
- Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM. 2017. Canu: scalable and accurate long-read assembly via adaptive k-mer weighting and repeat separation. *Genome Res* 27:722–736. <https://doi.org/10.1101/gr.215087.116>.
- Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, Cuomo CA, Zeng Q, Wortman J, Young SK, Earl AM. 2014. Pilon: an integrated tool for comprehensive microbial variant detection and genome assembly improvement. *PLoS One* 9:e112963. <https://doi.org/10.1371/journal.pone.0112963>.
- Tracey A, Foster JM, Paulini M, Grote A, Mattick J, Tsai Y-C, Chung M, Cotton JA, Clark TA, Geber A, Holroyd N, Korlach J, Libro S, Lustigman S, Michalski ML, Rogers MB, Twaddle A, Dunning Hotopp JC, Berriman M, Ghedin E. 2020. Nearly complete genome sequence of *Brugia malayi* strain FR3. *Microbiol Resour Announc* 9:e00154-20. <https://doi.org/10.1128/MRA.00154-20>.
- Seppy M, Manni M, Zdobnov EM. 2019. BUSCO: assessing genome assembly and annotation completeness. *Methods Mol Biol* 1962:227–245. https://doi.org/10.1007/978-1-4939-9173-0_14.
- Ihaka R, Gentleman R. R: a language for data analysis and graphics. *J Comput Graph Stat* 1996;5:299–314. <https://doi.org/10.1080/10618600.1996.10474713>.